

***Borniopsis mortoni* sp. n. (Heterodonta, Galeommatoidea, Galeommatidae sensu lato), a new bivalve commensal with a synaptid sea cucumber from Japan**

Ryutaro Goto¹, Hiroshi Ishikawa²

¹ Museum of Zoology and Department of Ecology and Evolutionary Biology, University of Michigan, 1109 Geddes Avenue, Ann Arbor, Michigan 48109-1079, USA ² 965-1 Kawachi-ko, Uwajima, Ehime 798-0075, Japan

Corresponding author: Ryutaro Goto (gotoryutaro@gmail.com)

Academic editor: R. C. Willan | Received 13 February 2016 | Accepted 29 July 2016 | Published 7 September 2016

<http://zoobank.org/7967128B-EDED-4C19-A14E-680F3109F676>

Citation: Goto R, Ishikawa H (2016) *Borniopsis mortoni* sp. n. (Heterodonta, Galeommatoidea, Galeommatidae sensu lato), a new bivalve commensal with a synaptid sea cucumber from Japan. ZooKeys 615: 33–45. doi: 10.3897/zookeys.615.8125

Abstract

The Galeommatoidea is a bivalve superfamily that exhibits high species diversity in shallow waters. Many members of this superfamily are associated commensally with burrowing marine invertebrates in benthic sediments. The genus *Borniopsis* is known only from eastern Asia and exhibits high host diversity (e.g., mantis shrimps, crabs, holothurians, sipunculans and echiurans). A new species, *Borniopsis mortoni* sp. n., is described from mud flats at the mouth of the Souzu River, southwestern Shikoku Island, Japan. This species has elongate-ovate shells covered by a tan to dark brown periostracum, and lives attached by both its foot and byssal threads to the body surface of the synaptid sea cucumber *Patinapta ooplax*. Several individuals of *B. mortoni* are often found on the same host, but sometimes more than 10 individuals can occur together. *Borniopsis mortoni* is one of the smallest species in this genus. Probably, its small body size is an adaptation to the mode of life in a narrow host burrow. Until now, only two other *Borniopsis* species were known to have commensal associations with synaptids. Thus, this is the third example of a synaptid-associated species from this genus. In addition, we briefly review the galeommatoideans commensal with apodid sea cucumbers.

Keywords

Apodida, Bivalvia, *Borniopsis*, *Byssobornia*, commensalism, Galeommatoidea, host shift, Holothuroidea, *Pseudopythina*, symbiosis, Synaptidae

Introduction

The Galeommatoidea is a superfamily of small bivalves that exhibits tremendous diversity in the intertidal zone (Bouchet et al. 2002, Paulay 2003, Lützen and Nielsen 2005). Many members of this superfamily are commensals associated in highly specific relationships with benthic invertebrates that burrow in soft sediments (Boss 1965, Morton and Scott 1989, Li et al. 2012, Goto et al. 2012). Most commensal galeommatoideans live attached directly onto the host's body surface or the walls of its burrow (Boss 1965, Morton and Scott 1989, Goto et al. 2012).

The genus *Borniopsis* was established for *B. tsurumaru* Habe, 1959 as the type species with a second species, *B. ariakensis* Habe, 1959, from Japan (Habe 1959). Morton and Scott (1989) described four *Pseudopythina* (*Borniopsis*) species from Hong Kong. After this publication, *Borniopsis* was frequently synonymized with *Pseudopythina* (e.g., Lützen et al. 2004, Goto et al. 2012). However, Huber (2015) noticed that these East Asian *Pseudopythina* species are distinct from the type species *P. macandrewi* (P. Fischer, 1867), which occurs only in the eastern Atlantic Ocean, in dentition, seminal receptacles, and demibranchs and thus he transferred them to *Borniopsis* together with some species of *Byssobornia*, *Squillaconcha* and *Kellia* (Huber, 2015), although *B. fujitaniana* (Yokoyama, 1927) was recently reassigned to the genus *Tellimya* (Goto et al., 2016). As a result, at least nine species currently belong to this genus – *B. tsurumaru*, *B. ariakensis*, *B. macrophthalmensis* (Morton & Scott, 1989), *B. maipoensis* (Morton & Scott, 1989), *B. nodosa* (Morton & Scott, 1989), *B. ochetostomae* (Morton & Scott, 1989), *B. subsinuata* (Lischke, 1871), *B. yamakawai* (Yokoyama, 1922) and *B. sagamiensis* (Habe, 1961) (Huber 2015). All of these species are known only from eastern Asia. Those *Borniopsis* species, for which the biology is known, are host-specific commensals associated with burrowing invertebrates (mantis shrimps, crabs, holothurians, sipunculans, echinurans and probably tanaids) (Kuroda 1937, Morton 1972, 1988, Morton and Scott 1989, Goto and Kato 2012, Goto et al. 2012). Host animals are different among the *Borniopsis* species, except for one species pair (*B. tsurumaru* and *B. ariakensis*), suggesting that this group diversified by repeated host shifts among various invertebrates in eastern Asia. This assumption is partially confirmed by molecular phylogeny (Goto et al. 2012).

In this study, we describe a new species of *Borniopsis*, which was collected from the synaptid sea cucumber *Patinapta ooplax* (von Marenzeller, 1881) on mud flats at the mouth of Souzu River, southwestern Shikoku Island, Japan. *Patinapta ooplax* is a small earthworm-like holothurian that burrows in muddy sediments in the intertidal zone. We compared its morphology and host associations with the other *Borniopsis* species. In addition, we reviewed galeommatoideans associated with apodid sea cucumbers.

Materials and methods

An undescribed species of *Borniopsis* was found attached to the synaptid sea cucumber *Patinapta ooplax* in the mud flats at the mouth of the Souzu River, Ainan-cho,

Ehime Prefecture, southwestern Shikoku Island, Japan (32°57'N, 132°33'E) on 20 May 2000. We collected the specimens of this bivalve species during spring low tides in 2000, 2001, 2003, 2006, 2007, 2009, 2012, and 2013. Seven specimens collected on 11 March 2012 and two specimens collected on 15 March 2013 were preserved in 100% ethanol and brought back to the laboratory and observed under a binocular dissecting microscope for description. One specimen collected on 12 April 2009 was bleached to remove the periostracum for observation of shell surface sculpture. We deposited the holotype and two paratypes in the Museum of Zoology, University of Michigan (UMMZ), and two paratypes in National Museum of Nature and Science, Tokyo (NSMT). We also observed specimens of *B. tsurumaru* (SBMNH 149526), *B. ariakensis* (SBMNH35056), *B. ochetostomae* (SBMNH 149525) and *B. maipoensis* (SBMNH 35126) loaned from the Santa Barbara Natural History Museum. All the loaned specimens were collected from the mud flats of Hong Kong.

Systematics

Superfamily Galeommatoidea J.E. Gray, 1840

Family Galeommatidae *sensu lato* (Ponder, 1998)

Genus *Borniopsis* Habe, 1959

***Borniopsis mortoni* sp. n.**

<http://zoobank.org/1782E002-A7F1-42D1-A1EC-0824A89B5A3E>

New Japanese name: Himoikarinamako-yadorigai

Figs 1–4

Material examined. Holotype (Figs 1, 2): UMMZ 305035 (SL 4.1 mm, SH 2.8 mm). Paratype 1 (Fig. 3): UMMZ 305036-1 (SL 3.9 mm, SH 2.7 mm), paratype 2: UMMZ 30536-2 (SL 2.4 mm, SH 1.8 mm), paratype 3: NSMT-Mo 78968 (SL 3.7 mm, SH 2.7 mm) and paratype 4 (Fig. 1B): NSMT-Mo 78969 (SL 3.3 mm, SH 2.3 mm). Non-type specimens: four individuals of *B. mortoni* (SL 2.4, 3.3, 3.8, 3.3 mm). All specimens were collected in the mud flats at the mouth of the Souzu River, Ainan-cho, Ehime Prefecture, southwestern Shikoku Island, Japan (32°57'N, 132°33'E). Comparative species (Fig. 4): *B. tsurumaru*, SBMNH 149526; *B. ariakensis*, SBMNH 35056; *B. ochetostomae*, SBMNH149525; and *B. maipoensis*, SBMNH 35126.

Type locality and habitat. Mud flats at the mouth of the Souzu River, Ainan-cho, Ehime, southwestern Shikoku Island, Japan (32°57'N, 132°33'E).

Description. Shell (Figs 1–3): Shell small (up to 4.1 mm), thin, slightly inflated, shape elongate-ovate and equivalve; inequilateral, longer anteriorly. Anterior and posterior margins rounded, ventral margins slightly rounded. Umbo small. Beak small, prosogyrate, situated 2/3 of way toward posterior. Each valve covered by tan to dark brown periostracum with black deposits, often eroded around beaks (Figs 2, 3). Shell surface underneath periostracum smooth and whitish with pearly luster (Fig. 1B).



Figure 1. *Borniopsis mortoni* and its host. **A** A crawling individual of *B. mortoni*. **B** Right shell valve of Paratype 4 (NSMT Mo 78969) bleached to remove the periostracum for observation of shell surface sculpture **C, D** *Patinapta ooplax* with multiple individuals of *B. mortoni* attached **E, F** *P. ooplax* in situ in mud flats with *B. mortoni* attached (arrowed). Photo credits: H. Ishikawa: **A–C, E, F**; Y. Hamamura: **D**. Scale bar: 1 mm.

Sculpture consisting of fine, dense and a few strong, widely spaced commarginal growth striae visible even on periostracum and very faint dense radial striae only visible underneath periostracum (Fig. 1B). Hinge of each valve consisting of a single stout cardinal tooth in front of umbo and well-developed oblique internal ligament posterior to umbo (Figs 2F, G; 3E, F). Soft parts (Figs 1, 2): Mantle not reflected,




Figure 2. Holotype of *Borniopsis mortoni* (UMMZ 305035) (SL 4.1 mm). **A, C, E** Right valve **B, D** Left valve **F, G** Hinge structure of left and right valves. Abbreviations: aa, anterior adductor muscle; ac, anterior cardinal tooth; f, foot; gn, gonad; il, internal ligament; lid, inner demibranch of left side; pa, posterior adductor muscle; rid, inner demibranch of right side; st, stomach. Scale bars 1 mm. Photo credits: R. Goto: **A–G**.

without prominent tentacles, edges narrowly extend beyond margin of shell, with numerous short papillae regularly arranged. Both anterior and posterior adductor muscles elongate-ovate, subequal and situated in the middle of dorsal and ventral margin. Ctenidia with gill axis nearly vertical, flat, consisting of single demibranch with both ascending and descending lamellae, joined anteriorly to inner and outer




Figure 3. Paratype 1 of *Borniopsis mortoni* (UMMZ 305035) (SL 2.4 mm). **A, C** Right valve **B, D** Left valve **E, F** Hinge structure of right and left valves. Abbreviations: ac, anterior cardinal tooth; il, internal ligament. Scale bars 0.5 mm. Photo credits: R. Goto: **A–F**.

labial palps. Labial palps leaf-shaped. Foot slender, of moderate size, with small heel; byssal glands located just in front of heel. Gonads situated from middle to posterior in visceral mass just below umbo. Stomach and digestive gland large, occupying anterior part of visceral mass.

Distribution. Only known from the type locality.

Host. *Patinapta ooplax* (Echinodermata: Holothuroidea: Apodida: Synaptidae) (Fig. 1C–F).

Host association. *Borniopsis mortoni* attaches to the body surface of *P. ooplax* by both its foot and byssal threads (Fig. 1C–F). Individuals were patchily distributed in the mud flats. Within the particular patches we sampled, approximately 70% of synaptids served as hosts for *B. mortoni*. Several *B. mortoni* often occurred on one the same host. At the maximum, more than 10 individuals were attached to a single host. Two

P. ooplax infested by *B. mortoni* also harbored the endoparasitic eulimid gastropod *Hypermastus lacteus* (A. Adams, 1864).

Etymology. The specific name is dedicated to Dr. Brian Morton who has made great contributions to marine biology, marine ecology and malacology. He discovered many interesting commensal galeommatoid species from Hong Kong, some of which now belong to the genus *Borniopsis*.

Remarks. The genus *Borniopsis* has been variously assigned to the Kelliidae (Morton and Scott 1989), Lasaeidae (Bieler et al. 2010), and the subfamily Montacutinae within Galeommatidae *sensu lato* (Huber 2015). However, these family- or subfamily-level groupings are ill-defined when a range of characters and taxa are considered (Ponder 1998). Indeed, the molecular analyses conducted by Goto et al. (2012) showed that each of these groups is actually polyphyletic. In this study, we assigned the genus *Borniopsis* tentatively to Galeommatidae *sensu lato*, which Ponder (1998) defined by the same diagnosis that was applied to the superfamily Galeommatoidae, as did Huber (2015). Further taxonomic assignment of this genus within Galeommatoidae (or Galeommatidae *sensu lato*) should be delayed until its family-level (or subfamily-level) classification is revised.

As with *B. mortoni*, both *B. tsurumaru* and *B. ariakensis* have a symbiotic relationship with synaptid sea cucumbers (Morton 1988, Morton and Scott 1989, Lützen et al. 2004, Kai and Henmi 2008). However, the particular host species differ between them – *Patinapta ooplax* (host solely for *B. mortoni*) and *Protankyra bidentata* (Woodward & Barrett, 1858) (host for both *B. tsurumaru* and *B. ariakensis*) (Morton and Scott 1989, Lützen et al. 2004). *Borniopsis mortoni* always attaches directly onto the body surface of the host (this study), whereas *B. tsurumaru* can attach to the body surface of the host, or the wall of the host's burrow, or to the carapace of commensal crabs living in the same burrows (Morton 1988, Morton and Scott 1989, Lützen et al. 2004, Kai and Henmi 2008, Goto et al. 2012). Furthermore, the number of bivalves per host is much higher in *B. mortoni* (several to more than 10) than *B. tsurumaru* and *B. ariakensis* (usually one) (Lützen et al. 2004, Goto, Ishikawa, and Hamamura, personal observations).

The shells of *B. tsurumaru* and *B. ariakensis* are much larger (up to 11–12 mm in SL) than those of *B. mortoni* (up to 4.1 mm) (Morton and Scott 1989) (Fig. 4A, B). Probably, this corresponds with the size of the host because *P. bidentata* is much larger than *P. ooplax*. The shells of *B. tsurumaru* and *B. ariakensis* are thicker and more inflated than those of *B. mortoni* (Lützen et al. 2004; this study). In addition, the shells of *B. mortoni* are always covered by a dark brown periostracum, whereas those of *B. tsurumaru* and *B. ariakensis* are often whitish, although some are dark brown. The umbones of *B. tsurumaru* and *B. ariakensis* are more protruding than those of *B. mortoni* (Fig. 4A, B). A molecular analysis is needed to understand whether these three synaptid-associated species are monophyletic or not. In addition, morphological variation of *B. tsurumaru* and *B. ariakensis* is apparently continuous (Goto, Ishikawa and Hamamura, pers. obs.) so molecular testing should be employed to investigate whether they can be distinguished genetically or not.




Figure 4. **A** *Borniopsis tsurumaru* (SBMNH 149526) **B** *B. ariakensis* (SBMNH 35056) **C** *B. ochetostomae* (SBMNH 149525) and **D** *B. maipoensis* (SBMNH 35126). Photo credits: R. Goto: **A–C**; P. Valentich-Scott: **D**. Scale bars 1 mm.

The present new species also closely resembles *B. ochetostomae* and *B. maipoensis* in having an elongate ovate shell covered by a brownish periostracum (Fig. 4C, D). However, *B. ochetostomae* is much larger (up to 11 mm) than *B. mortoni* and its beak is located more centrally than that of *B. mortoni* (Morton and Scott 1989, Jespersen et al. 2002, this study) (Fig. 4C). On the other hand, *B. maipoensis* is rather smaller (up to 3 mm) and more rounded than *B. mortoni* (Morton and Scott 1989, this study) (Fig. 4D). In addition, *B. maipoensis* has two distinct papillae on the dorsal surface of the foot (Morton and Scott 1989), whereas we did not observe such papillae on *B. mortoni* (Fig. 1A). And lastly, the hosts for these three species are different – *B. mortoni* (holothurian hosts), *B. ochetostomae* (echiuran hosts) and *B. maipoensis* (probably tanaid hosts) (Morton and Scott 1989, this study).

Key to *Borniopsis*

- 1 Beaks subcentral 2
- Beaks near posterior end 3
- 2 Shell outline elliptical to subelliptical..... 4

–	Shell outline trigonal to subtrigonal.....	5
3	Shell outline subovate, without nodules on inner surface of anterior valve margin.....	6
–	Shell outline quadrate, with nodules on inner surface of anterior valve margin.....	<i>B. nodosa</i>
4	SH/SL 0.60–0.66 (Morton and Scott 1989).....	<i>B. ochetostomae</i>
–	SH/SL 0.73 (this study).....	<i>B. yamakawai</i>
5	Periostracum brownish, max. SL < 4 mm.....	<i>B. macrophthalmensis</i>
–	Periostracum whitish, max. SL > 4 mm.....	7
6	Umbo slightly protruding.....	8
–	Umbo not protruding, max. SL < 5 mm.....	<i>B. mortoni</i>
7	SH/SL 0.80 (Morton, 1972).....	<i>B. subsinuata</i>
–	SH/SL 0.59–0.64 (Habe, 1961).....	<i>B. sagamiensis</i>
8	With two distinctive papillae on dorsal surface of foot, max. SL < 3 mm.....	<i>B. maipoensis</i>
–	Without two distinctive papillae on dorsal surface of foot, max. SL > 3 mm...	9
9	SH/SL 0.73 (this study).....	<i>B. tsurumaru</i>
–	SH/SL 0.66 (this study).....	<i>B. ariakensis</i>

Discussion

Apodid sea cucumbers, including members of the families Synaptidae and Chiridotidae, are one of the major hosts for galeommatoideans (Boss 1965, Morton and Scott 1989, Kato 1998, Middelfart and Craig 2004) (Table 1). Including *B. mortoni*, at least 13 species are known to have commensal associations with apodid holothurians (Table 1). They can be separated into the following four groups: the first group includes *Anisodevonia*, *Austrodevonia*, *Devonia* and *Entovalva*, which have reduced shells covered by well-developed mantle lobes (Kawahara 1942, Kato 1998, Middelfart and Craig 2004); the second group includes *Borniopsis* covered in this study (Morton 1988, Morton and Scott 1989); the third groups includes *Montacuta* (Bateson 1923, Fox 1979); and the forth group includes *Scintillona* (Morton 1957, Ó Foighil and Gibson 1984). All of these four groups live attached to the apodid's body surface, except for *Entovalva*, which lives inside the host's oesophagus (Spärk 1931, Kato 1998). Molecular phylogenies suggested that associations with apodids have evolved repeatedly in the Galeommatoidea (Goto et al. 2012).

Borniopsis mortoni lives commensally with the apodid sea cucumber *P. ooplax* on the temperate coast of western Shikoku Island. On the other hand, another galeommatoidean species, *Anisodevonia ohshimai*, lives attached to *P. ooplax* on the subtropical coast of the Ryukyu Islands, southwestern Japan (Kawahara 1942, Kato 1998). Extensive sampling for *A. ohshimai* has been undertaken in the Ryukyu Islands (Kawahara 1942, Kato 1998, Kosuge 2001). However, *B. mortoni* has never been found there. Thus, *B. mortoni* probably does not occur on the subtropical coast of the Ryukyu Islands and may be restricted to more temperate coasts. An alternative hypothesis is that *P. ooplax* used by

Table 1. Galeommatoidan bivalves commensal with apodid sea cucumbers.

Species	Host	Distribution	References
<i>Anisodevonia obshimai</i> (Kawahara, 1942)	<i>Patinapta ooplax</i>	NW Pacific	Kawahara 1942; Kato 1998
<i>Austrodevonia sharnae</i> Middelfart & Craig, 2004	<i>Taeniogyrus australianus</i>	NW Pacific	Middelfart and Craig 2004
<i>Devonia perrieri</i> (Malard, 1903)	<i>Leptosynapta inhaerens</i>	NE Atlantic	Malard 1903
<i>Devonia semperi</i> (Oshima, 1930)	<i>Protankyra bidentata</i>	NW Pacific	Oshima 1930
<i>Devonia</i> sp.	<i>Protankyra similis</i>	NW Pacific	Semper 1868
<i>Entovalva amboinensis</i> (Spärck, 1931)	<i>Patinapta laevis</i>	NW Pacific	Spärck 1931
<i>Borniopsis tsurumaru</i> Habe, 1959	<i>Protankyra bidentata</i>	NW Pacific	Morton and Scott 1989
<i>Borniopsis ariakensis</i> Habe, 1959	<i>Protankyra bidentata</i>	NW Pacific	Morton and Scott 1989
<i>Borniopsis mortoni</i> sp. n.	<i>Patinapta ooplax</i>	NW Pacific	this study
<i>Montacuta donacina</i> (Wood, 1848)	<i>Leptosynapta inhaerens</i>	NE Atlantic	Bateson 1923
<i>Montacuta percompressa</i> Dall, 1899	<i>Leptosynapta tenuis</i>	NE Atlantic	Fox 1979
<i>Scintillona bellerophon</i> Ó Foighil & Gibson, 1984	<i>Leptosynapta clarki</i>	NE Pacific	Ó Foighil and Gibson 1984
<i>Scintillona zelandica</i> (Odhner 1924)	<i>Taeniogyrus dendyi</i>	New Zealand	Finlay 1927

A. obshimai in the Ryukyu Islands is a different species to that used by *B. mortoni* in eastern Shikoku Island. The taxonomy of this group of synaptid holothurians remains poorly understood and thus it is highly probable that *P. ooplax* is a species complex.

A new species of the genus *Borniopsis* is described herein. This genus is restricted to eastern Asia, and each species apparently utilizes a different invertebrate host (Table 2), suggesting that its diversification was caused by repeated host shifting. The shell size varies among species (Table 2), which is probably a specialization to each particular commensal lifestyle. *Borniopsis macrophthalmensis*, *B. maipoensis* and *B. mortoni* are the three smallest species in this genus (Table 2). *Borniopsis macrophthalmensis* is attached to the body surface of highly-mobile intertidal crabs only by fine byssal threads (Morton and Scott 1989). If *B. macrophthalmensis* had a large, heavy shell, it would easily fall off the host crab when it scurries quickly on the surface of the mudflat. Thus, such a small shell is probably an adaptation for life on a highly-mobile crab. Similarly, another galeommatoidan species, *Arthritica japonica* Lützen & Takahashi, 2003, which is also attached to the body surface of crabs, has very small shells (up to 2.05 mm) (Lützen and Takahashi 2003). This may represent a morphological convergence to a similar commensal lifestyle and supports our hypothesis of reduction in shell size associated with fast-moving hosts mentioned above. On the other hand, the second species *B. maipoensis* lives commensally with the tanaid *Discapseudes* sp. (Morton and Scott 1989), whereas the third species *B. mortoni* lives with the synaptid sea cucumber *P. ooplax*. The diameter of these host burrows is very small. Thus, the small-sized shells of *B. maipoensis* and *B. mortoni* are probably an adaptation to the mode of life in narrow host burrows.

The mudflats of eastern Asia evidently possess one of the richest burrowing invertebrate faunas in the world. However, burrow associates remain poorly understood in many of them. Thus, further investigation in this area could reveal increased diversity of this distinctive bivalve genus *Borniopsis*.

Table 2. *Borniopsis*, its host and maximum size (shell length).

Species	Host	Max. SL (mm)	References
<i>Borniopsis ariakensis</i> Habe, 1959	holothurian (<i>Protankyra bidentata</i>)	12.3	Habe 1959
<i>Borniopsis macrophthalmensis</i> (Morton & Scott 1989)	crab (<i>Macrophthalmus</i>)	3.2	Morton and Scott 1989
<i>Borniopsis maipoensis</i> (Morton & Scott 1989)	probably tanaid (<i>Discapseudes</i> sp.)	3.0	Morton and Scott 1989
<i>Borniopsis mortoni</i> sp. n.	holothurian (<i>Patinapta ooplax</i>)	4.1	this study
<i>Borniopsis nodosa</i> (Morton & Scott 1989)	sipunculan (<i>Sipunculus nudus</i>)	6.1	Morton and Scott 1989
<i>Borniopsis ochetostomae</i> (Morton & Scott 1989)	echiuran (<i>Listriolobus sorbillans</i>)	10.1	Morton and Scott 1989
<i>Borniopsis sagamiensis</i> (Habe 1961)	unknown	19.4	Habe 1961
<i>Borniopsis subsinuata</i> (Lischke, 1871)	mantis shrimp (<i>Squilla</i> , <i>Oratosquilla</i>)	12.0	Morton 1972
<i>Borniopsis tsurumaru</i> Habe, 1959	holothurian (<i>Protankyra bidentata</i>)	10.7	Habe 1959
<i>Borniopsis yamakawai</i> (Yokoyama, 1922)	echiuran (<i>Ochetostoma erythrogrammon</i>)	11.0	Goto and Kato 2012; this study

Acknowledgements

We wish to express our sincere gratitude to the following people: Y. Hamamura (Hiroshima Prefecture) for providing the information of *B. tsurumaru* and *B. ariakensis* and the images on *B. mortoni*; P. Valentich-Scott (Santa Barbara Natural History Museum) for allowing us to loan the specimens and use of his image on *B. maipoensis*; D. Ó Foighil (University of Michigan) for hosting RG as a research fellow for this study and permission to use the facilities in the laboratory; L. Taehwan (University of Michigan) for helping RG to set up the laboratory facilities to observe specimens; R. C. Willan (Museum and Art Gallery of the Northern Territory), E. V. Coan (Santa Barbara Natural History Museum) and A. Holmes (National Museum Wales) for comments that greatly improved the manuscript. This study was partially supported by grants to RG from the Japan Society for the Promotion of Science (JSPS) Research Fellowship for Young Scientists and the JSPS Postdoctoral Fellowships for Research Abroad.

References

- Bateson G (1923) Specimen of *Synapta inhaerens* with *Montacuta donacina* attached thereto. Proceedings of the Malacological Society of London 15: 266–267.
- Bieler R, Carter GJ, Coan EV (2010) Classification of bivalve families. Malacologia 52: 113–184.
- Boss KJ (1965) Symbiotic erycinacean bivalves. Malacologia 3: 183–195.
- Bouchet P, Lozouet P, Maestrati P, Heros V (2002) Assessing the magnitude of species richness in tropical marine environments: exceptionally high numbers of molluscs at a New Caledonia site. Biological Journal of the Linnean Society 75: 421–436. doi: 10.1046/j.1095-8312.2002.00052.x

- Finlay HJ (1927) A further commentary on New Zealand molluscan systematics. Transactions and Proceedings of the New Zealand Institute 57: 320–485.
- Fox TH (1979) Reproductive adaptations and life histories of the commensal leptonacean bivalves. Unpublished Ph.D. dissertation, University of North Carolina, Chapel Hill, 207 pp.
- Goto R, Kato M (2012) Geographic mosaic of mutually exclusive dominance of obligate commensals in symbiotic communities associated with a burrowing echinuran worm. Marine Biology 159: 319–330. doi: 10.1007/s00227-011-1810-8
- Goto R, Kawakita A, Ishikawa H, Hamamura Y, Kato M (2012) Molecular phylogeny of the bivalve superfamily Galeommatoidea (Heterodonta, Veneroidea) reveals dynamic evolution of symbiotic life style and interphylum host switching. BMC Evolutionary Biology 12: 172. doi: 10.1186/1471-2148-12-172
- Goto R, Ishikawa H, Hamamura Y (2016) Symbiotic association of the bivalve *Tellimya fujitaniana* (Galeommatoidea) with the heart urchin *Echinocardium cordatum* (Spatangoida) in the northwestern Pacific. Zoological Science 33: 434–440. doi: 10.2108/zs150215
- Habe T (1959) Five new minute bivalves from Japan (Erycinacea: Pelecypoda). Publications of the Seto Marine Biological Laboratory 7: 291–294.
- Huber M (2015) Compendium of Bivalves 2. A full-color guide to the remaining seven families. A systematic listing of 8500 bivalve species and 10500 synonyms. ConchBooks, Harxheim, 907 pp.
- Jespersen Å, Lützen J, Morton B (2002) Ultrastructure of dimorphic sperm and seminal receptacle in the hermaphrodites *Barrimysia siphonosomae* and *Pseudopythina ochetostomae* (Bivalvia, Galeommatoidea). Zoomorphology 121: 159–172. doi: 10.1007/s00435-001-0053-5
- Kai T, Henmi Y (2008) Description of zoeae and habitat of *Elamenopsis ariakensis* (Brachyura: Hymenosomatidae) living within the burrows of the sea cucumber *Protankyra bidentata*. Journal of Crustacean Biology 28: 342–351. doi: 10.1163/20021975-99990379
- Kato M (1998) Morphological and ecological adaptations in montacutid bivalves endo- and ecto-symbiotic with holothurians. Canadian Journal of Zoology 76: 1403–1410. doi: 10.1139/z98-077
- Kawahara T (1942) On *Devonia ohshimai* sp. n., a commensal bivalve attached to the synaptid *Leptosynapta ooplax*. Venus 11: 153–164.
- Kosuge T (2001) Ecological notes on the populations of the galeommatid bivalve *Anisodevonia ohshimai* in Ishigaki Island, Southern Japan. The Biological Magazine Okinawa 39: 5–10.
- Kuroda T (1937) The habit of "*Kellia*" *subsinuata* Lischke. Venus 7: 188–189.
- Li J, Ó Foighil D, Middelfart P (2012) The evolutionary ecology of biotic association in a mega-diverse bivalve superfamily: sponsorship required for permanent residency in sediment. PLoS ONE 8: e42121. doi: 10.1371/journal.pone.0042121
- Lützen J, Nielsen C (2005) Galeommatid bivalves from Phuket, Thailand. Zoological Journal of the Linnean Society 144: 261–308. doi: 10.1111/j.1096-3642.2005.00168.x
- Lützen J, Takahashi T (2003) *Arthritica japonica*, sp. n. (Bivalvia: Galeommatoidea: Leptonidae), a commensal with the pinnotherid crab *Xenophthalmus pinnotheroides* White, 1846. Yuriyagai 9: 11–19.
- Lützen H, Jespersen Å, Takahashi T, Kai T (2004) Morphology, structure of dimorphic sperm, and reproduction in the hermaphroditic commensal bivalve *Pseudopythina tsurumaru* (Galeommatoidea: Kellidae). Journal of Morphology 262: 407–420. doi: 10.1002/jmor.10158

- Malard AE (1903) Sur un lamellibranche nouveau, parasite des synaptés. Bulletin du Muséum National d'Histoire Naturelle (Paris) 9: 342–346.
- Middelfart P, Craig M (2004) Description of *Austrodevonia sharnae* n. gen. n. sp. (Galeommataceae: Bivalvia), an ectocommensal of *Taeniogyrus australianus* (Stimpson, 1855) (Synaptidae: Holothuroidea). Molluscan Research 24: 211–219. doi: 10.1071/MR04014
- Morton B (1972) Some aspects of the functional morphology and biology of *Pseudopythina sub-sinuata* (Bivalvia: Leptonacea) commensal on stomatopod crustaceans. Journal of Zoology 166: 79–96. doi: 10.1111/j.1469-7998.1972.tb04078.x
- Morton B (1988) Partnerships in the Sea: Hong Kong's Marine Symbioses. Hong Kong University Press, Hong Kong, 140 pp. <http://www.jstor.org/stable/j.ctt2jc6c7>
- Morton B, Scott PH (1989) The Hong Kong Galeommataceae (Mollusca: Bivalvia) and their hosts, with descriptions of new species. Asian Marine Biology 6: 129–160.
- Morton JE (1957) The habits of *Scintillona zelandica* (Odher) 1924 (Lamellibranchia: Galeommataceae). Journal of Molluscan Studies 32: 185–188.
- Ó Foighil D, Gibson A (1984) The morphology, reproduction and ecology of the commensal bivalve *Scintillona bellerophon* spec. n. (Galeommataceae). Veliger 27: 72–80.
- Ohshima H (1930) Preliminary note on *Entovalva semperi* sp. n., a commensal bivalve living attached to the body of a synaptid. Annotationes Zoologicae Japonenses 13: 25–27.
- Paulay G (2003) Marine Bivalvia (Mollusca) of Guam. Micronesica 35–36: 218–243.
- Ponder WF (1998) Superfamily Galeommatoidea. In: Beesley PL, Ross GJB, Wells AE (Eds) Mollusca: the Southern Synthesis. CSIRO Publishing, Melbourne, 316–318.
- Spärck R (1932) *Cycladoconcha amboinensis* n. gen. n. sp., a commensalistic lamellibranch. Papers from Dr Th. Mortensen's Pacific-Expedition 1914–1916 58: 227–237.
- Semper C (1868) Reisen im Archipel der Philippinen. 2. Wissenschaftliche Resultate. 1. Holothurien. Wilhelm Engelmann, Leipzig.