



An ammonite trapped in Burmese amber

Tingting Yu^{a,b,c,1}, Richard Kelly^{a,d,e,1}, Lin Mu^{a,1}, Andrew Ross^e, Jim Kennedy^{f,g}, Pierre Broly^h, Fangyuan Xiaⁱ, Haichun Zhang^{a,b}, Bo Wang^{a,b,j,2}, and David Dilcher^{k,2}

^aState Key Laboratory of Palaeobiology and Stratigraphy, Nanjing Institute of Geology and Palaeontology, Chinese Academy of Sciences, Nanjing 210008, China; ^bCAS Center for Excellence in Life and Palaeoenvironment, Chinese Academy of Sciences, Nanjing 210008, China; ^cSchool of Earth and Space Sciences, University of Science and Technology of China, Hefei 230026, China; ^dSchool of Earth Sciences, University of Bristol, Bristol B58 1TQ, United Kingdom; ^eDepartment of Natural Sciences, National Museum of Scotland, Edinburgh EH1 1JF, United Kingdom; ^fOxford University Museum of Natural History, Oxford OX1 3PW, United Kingdom; ^gDepartment of Earth Sciences, University of Oxford, Oxford OX1 3AN, United Kingdom; ^hPrivate address, 59840 Lompref, France; ⁱLingpoge Amber Museum, Shanghai 201108, China; ^jKey Laboratory of Zoological Systematics and Evolution, Institute of Zoology, Chinese Academy of Sciences, Beijing 100101, China; and ^kDepartment of Geology and Atmospheric Science, Indiana University, Bloomington, IN 47405

Contributed by David Dilcher, February 27, 2019 (sent for review December 13, 2018; reviewed by Phillip Barden and Enrique Peñalver Mollá)

Amber is fossilized tree resin, and inclusions usually comprise terrestrial and, rarely, aquatic organisms. Marine fossils are extremely rare in Cretaceous and Cenozoic ambers. Here, we report a record of an ammonite with marine gastropods, intertidal isopods, and diverse terrestrial arthropods as syninclusions in mid-Cretaceous Burmese amber. We used X-ray-microcomputed tomography (CT) to obtain high-resolution 3D images of the ammonite, including its sutures, which are diagnostically important for ammonites. The ammonite is a juvenile *Puzosia* (*Bhimaites*) and provides supporting evidence for a Late Albian–Early Cenomanian age of the amber. There is a diverse assemblage (at least 40 individuals) of arthropods in this amber sample from both terrestrial and marine habitats, including Isopoda, Acari (mites), Araneae (spiders), Diplopoda (millipedes), and representatives of the insect orders Blattodea (cockroaches), Coleoptera (beetles), Diptera (true flies), and Hymenoptera (wasps). The incomplete preservation and lack of soft body of the ammonite and marine gastropods suggest that they were dead and underwent abrasion on the seashore before entombment. It is most likely that the resin fell to the beach from coastal trees, picking up terrestrial arthropods and beach shells and, exceptionally, surviving the high-energy beach environment to be preserved as amber. Our findings not only represent a record of an ammonite in amber but also provide insights into the taphonomy of amber and the paleoecology of Cretaceous amber forests.

(at least 40 individuals) of arthropods in this amber sample that live today in both terrestrial and marine habitats. Of the terrestrial fauna, Acari (mites) are the most abundant, with 23 specimens; also present are Araneae (spiders), Diplopoda (millipedes), and representatives of the insect orders Blattodea (cockroaches), Coleoptera (beetles), Diptera (true flies), and Hymenoptera (wasps). The arthropod assemblage consists mostly of forest floor-dwelling taxa, and living representatives are generally associated with leaf litter or the top layers of soil. There are several isopods preserved which are consistent with littoral or supralittoral taxa. In addition to the ammonite itself, four definitively marine gastropod shells and one putatively marine isopod are present.

Ammonite. The ammonite is a juvenile (the adapertural septa are not crowded), has a maximum preserved diameter of 12 mm, and appears to retain the original aragonitic shell, on the basis of its appearance in reflected light (Fig. 2A). It is composed in part by the body chamber, but the apertural part is damaged, as revealed by the survival of a 60° sector of the umbilical wall extending beyond the fragment of the inner flanks of the shell (Fig. 2). Coiling is moderately involute, with ~64% of the previous whorl covered. The small, shallow umbilicus comprises 18% of the diameter, the low umbilical wall is very weakly convex, and the umbilical shoulder is broadly rounded. The whorl section is

amber | ammonite | fossil | paleoecology | taphonomy

Amber provides a unique mode of preservation for organisms, and when inclusions are present they are usually 3D fossils of terrestrial plants, microorganisms, arthropods, and even vertebrate remains (1–3). Amber deposits are therefore considered to be exceptional Lagerstätten, providing unique windows into past ecosystems (4–6). Given that amber is formed by the fossilization of terrestrial plant resins, the capture of marine inclusions may be considered extremely rare. However, some recent findings of marine and freshwater fossils, particularly, microfossils such as diatoms, radiolarians, ostracods, and copepods, have provided fresh insights into amber taphonomy (7–12).

Burmese amber (from northern Myanmar) contains the most diverse biota of all known Cretaceous ambers (13, 14). Over the last 100 years, and particularly in the past two decades, Burmese amber has received worldwide scientific interest; more than 500 families of invertebrates, vertebrates, protists, plants, and fungi have been reported (15). Here, we provide an account of an exceptional piece of amber that preserves a unique assemblage of marine macrofossils, alongside intertidal, fully terrestrial, and possibly freshwater aquatic arthropods.

Results

The ammonite-bearing piece of amber (BA18100) was obtained from an amber mine located near Noiye Bum Village, Tanaing Town (ref. 16 and Fig. 1). It is 33 mm long, 9.5 mm wide, and 29 mm high, and its weight is 6.08 g. There is a diverse assemblage

Significance

Aquatic organisms are rarely found in amber, but when they occur they provide invaluable evidence for the better understanding of amber taphonomy and past ecosystems. We report an ammonite and several marine gastropods alongside a mixed assemblage of intertidal and terrestrial forest floor organisms in mid-Cretaceous Burmese amber. Our discovery indicates that the Burmese amber forest was living near a dynamic and shifting coastal environment. The ammonite also provides supporting evidence for the age of the amber, which is still debated, and represents a rare example of dating using fossils present inside the amber.

Author contributions: B.W. and D.D. designed research; T.Y., R.K., L.M., A.R., J.K., and B.W. performed research; T.Y. and F.X. contributed new reagents/analytic tools; T.Y., R.K., L.M., A.R., J.K., P.B., H.Z., B.W., and D.D. analyzed data; and T.Y., R.K., L.M., A.R., J.K., B.W., and D.D. wrote the paper.

Reviewers: P.B., New Jersey Institute of Technology; and E.P.M., Instituto Geológico y Minero de España.

The authors declare no conflict of interest.

This open access article is distributed under [Creative Commons Attribution License 4.0 \(CC BY\)](https://creativecommons.org/licenses/by/4.0/).

¹T.Y., R.K., and L.M. contributed equally to this work.

²To whom correspondence may be addressed. Email: bowang@nigpas.ac.cn or dilcher@indiana.edu.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1821292116/-DCSupplemental.

Published online May 13, 2019.




Fig. 2. Ammonite *Puzosia* (*Bhimaites*) Matsumoto. (A) Lateral view under light microscopy. (B) Flattened sutures reconstructed by microtomography. (C) Microtomographic reconstruction, apparent view. (D) Microtomographic reconstruction, surface rendering; (E) Microtomographic reconstruction, virtual section. (Scale bars, 2 mm.)

There are a few other specimens that are probably isopods, including one that is badly damaged with most of its ventral side obscured by a gastropod. The coloration and coxal plates are similar to those of the specimen shown in Fig. 3A, but there is a partial eye preserved, and it is larger. It is closely associated with a gastropod, but this is probably taphonomic. There are two other very badly damaged specimens, which may be isopods and two others, which may be isopod larvae, but they are poorly preserved.

Although taxonomic assignment is difficult based on specimens in which key characters cannot be observed, the specimens present in the piece of amber seem to be consistent with littoral or supralittoral isopods, with one possible fully marine species.

Gastropods. Four marine gastropod shells are also preserved with the ammonite (Fig. 4), of which two are well-preserved and can be attributed to the genus *Mathilda* Semper (Mathildidae) by the small, conical shell with heterostrophic protoconchs, whorl sides rounded and basally subcarinate, base broadly arched, and ornament of strong spiral cords and fine axial threads (34). *Mathilda* was mainly distributed in the western Tethys sea during the Cretaceous, and our fossils are a Cretaceous record of this genus from the eastern Tethys sea.

Terrestrial Arthropod Assemblage. There are 22 oribatid mites in the piece, most of which are ptyctimous, meaning they can close their prodorsum over their legs as protection against predators (box mites). Most of the mites (15 individuals) are similar in appearance to Phthiracaridae (Fig. 5B), but the ventral shields would need to be examined to clarify this, which is difficult given their position in the amber (Fig. 5A). They have previously only been described from Baltic amber and younger deposits. Also, although obscured in the piece, there appear to be representatives

of Euphthiracaroida (Fig. 5C), based on the fusion of the plates, and some Brachypylina.

Ptyctimous oribatids are common in soil/forest floor communities and are usually considered to be secondary decomposers/fungivores,




Fig. 3. Isopods of uncertain taxonomic affinity, but generally consistent with littoral or supralittoral taxa. (A) Isopod 1. (B) Isopod 2. (C) Isopod 3. (D) Circular structure attached to the dorsal surface of isopod 2. (Scale bars, 1 mm in A and C. Scale bar, 0.5 mm in B and D.)




Fig. 4. Gastropods. (A) *Mathilda* sp. (B) *Mathilda* sp. (C) Undetermined specimen. (D) Undetermined specimen. (Scale bars, 1 mm.)

although some ptyctimous mites are known to be xylophagous and feed on dead wood as primary decomposers (35). Either way, they are generally found in the presence of decaying plant material (36, 37). Oribatids are generally free living in the upper soil layers, but Phthiracaroida are commonly found within fallen leaves or conifer needles (37), and ground-dwelling species can be found up to 4 m from the ground on live tree trunks (38) with a clear gradient of community species composition ascending the trunk (39).

There is one spider preserved (Fig. 5D) that is unfortunately partly decomposed, and the eyes and chelicerae are not well preserved, so it is difficult to identify, but it is similar in general appearance to some Cretaceous Oonopidae which have been found in amber from Canada and Myanmar (40, 41). Oonopidae (goblin spiders) are described by Penney (40) as wandering, active predators, fast moving and nocturnal, and are known from a varied range of habitats (42), including the forest floor or tree bark (43).

There are 12 adult insects preserved in the piece, eight of which are true flies (Diptera), two are beetles (Coleoptera), one is a parasitic wasp (Hymenoptera), and one is a cockroach (Blattodea). There are also several larval specimens. Diptera is mostly represented by small nematoceran midges or gnats (Ceratopogonidae, Cecidomyiidae, or Chironomidae), and there are two small brachyceran hump-backed flies with cyclorhaphan-type antennae and wing venation (Fig. 5F) consistent with scuttleflies (Phoridae). Some nematoceran midges (e.g., chironomids) have aquatic larvae, which are usually found in freshwater habitats, but others (ceratopogonids and cecidomyiids) could have terrestrial or plant gall stages. One of them may be a gall midge that could have been associated with trees, similar to the parasitoid wasp (Fig. 5G), which belongs to Chrysoidea.

There are two beetles preserved in the amber, but they are largely obscured by other material. The larger one (Fig. 5H) is obscured by a gastropod, but the characters that can be seen include hind and mid legs with the femur expanded; tibiae narrow at the base, expanding toward apex; thick tibial spines reaching at least the length of the first tarsomere; five-segmented tarsi, gradually reducing in length from the first to the fifth, with each expanding in width from base to apex; ring of clumped hairs around the apices of the tibiae and each tarsomere, except the fifth that has two claws; and fore leg curved in a raptorial style. The pronotum is transverse; the elytra are distorted but appear oval-shaped and may have a black-and-white-banded color pattern; the head (and antennae) are either not preserved or are obscured by a gastropod shell.

The cockroach (Fig. 5I) is about 20 mm long, with most of the head, thorax, and abdomen missing, but the general overall shape is preserved. Also, many important characters, such as the

antennae, maxillary and labial palps, and the right fore, mid, and hind legs are preserved. There are no remains of the wings. The maxillary palps are long; the head appears quite narrow, but not entirely preserved. The partial preservation suggests that not all of the cockroach became engulfed in the resin, and the exposed




Fig. 5. Amber inclusions. (A) Amber piece showing most large inclusions. (B) Acari: Phthiracaridae. (C) Acari: Euphthiracoidea. (D) Araneae: Oonopidae. (E) Diplopoda. (F) Diptera: Phoridae. (G) Hymenoptera: Chrysoidea. (H) Coleoptera. (I) Blattodea. (Scale bar, 5 mm in A. Scale bars, 1 mm in E and H. Scale bars, 0.5 mm in B–D, F, and G. Scale bar, 2 mm in I.)

parts decomposed before it was covered by another layer. Many cockroaches are found in forest floor, leaf litter habitats, and many are found associated with decaying wood, although others are arboreal or aquatic (44).

One millipede is preserved (Fig. 5E) that is around 15 mm long, very slender, and has relatively short legs. The amber around the millipede is cloudy, so it is difficult to observe characters in detail, certainly important characters such as the organ of Tömösváry or ozopores. Millipedes are important detritivores and are mainly forest floor dwellers and are considered to have been for their whole evolutionary history (45). The body form of the specimen probably matches the “borer” (platydesmoid) type suggested by Kime and Golovatch (46), which suggests that it inhabited the leaf litter or uppermost soil layers (stratobiont) or that it was an underbark xylobiont (47).

Discussion

U-Pb dating of zircons from the volcanoclastic matrix of the amber has given a maximum age of 98.8 ± 0.6 Ma (48), which places it in the Early Cenomanian based on the 100.5 ± 0.4 Ma age assigned to the base of the stage by Cohen et al. (49); however, this dates the amber-bearing horizon, not the amber itself. This age is incompatible with the record of the exclusively Upper Albian ammonite *Mortoniceras*, which was found in a sandstone above primary Burmese amber deposits (16). The specimen of this ammonite was neither described nor figured, and we could not examine it, as attempts to locate the specimen have not been successful; therefore we cannot confirm its identity. Thus, the incompatibility of the age and the *Mortoniceras* ammonite remains unresolved. The presence of borings of martesine bivalves in the outer rim of pieces of Burmese amber suggested that the amber could be older than the age of the bed it was collected from. Bivalves have also been found within the amber and therefore bored into it while the amber was still soft and are thus similar in age to the bed (50).

Amber pieces can be reworked and redeposited in younger deposits; therefore, dating amber is sometimes controversial. The amber-bearing strata can be dated from palynofloras, ammonites, and radiodating evidence, but the amber could be older. Marine inclusions can help date ambers, as marine diatoms and other marine microfossils supported an Albian–Cenomanian age of Charentese ambers of France (7, 51). The present discovery is another interesting example of dating using fossils present inside the amber.

How did the amber that would have flowed from a tree capture both terrestrial (insects, millipedes, spiders, and mites) and marine (ammonite, gastropods, and isopods) organisms? Analysis of the depositional environment supports the model of an estuarine, coastal landscape for the mid-Cretaceous amber forests. Poinar et al. (52) analyzed Burmese amber and found that the most likely origin of the resin which formed the amber was araucarian conifers (but see ref. 53), which can be closely associated with coastal habitats. Many pieces of Burmese amber were bored by martesine pholadid bivalves, indicating that the amber was deposited in a brackish nearshore environment (50). Martesine bivalves have also been found within the amber, indicating that the resin was still soft when the bivalves started boring into it, which suggests that resin-producing trees were growing near to the site of deposition. The ammonite and gastropods had suffered damage before entombment. For example, the ammonite had lost at least a 60° sector of its body chamber, indicating that this was not the shell of a live individual. There is no evidence of any soft-part preservation of any of the gastropods, which also suggests that these were dead shells. The amber also contains some shell sand.

Of the many thousands of specimens of Burmese amber studied, only one ammonite is known. It is an exceptional occurrence and may record an exceptional event. There are three

possible scenarios: (i) There was a sandy beach with resin-producing trees growing very close. The terrestrial insects were trapped in the resin while it was still on the tree, and as it traveled down the tree trunk it picked up the lower-lying terrestrial arthropods, such as the mites. When it reached the ground (the beach?) it landed on the sand and shells, trapping the supralittoral isopods as they traversed the beach. As these forests are considered to have been coastal, this scenario could have been commonplace, but the probability of such amber pieces surviving would be slight, owing to the dynamic nature of beaches, which would explain the rarity of such pieces of amber in the fossil record. (ii) There was a tsunami that flooded the amber-producing forest, bringing marine debris into the forest and thus into contact with numerous blobs of resin. This would certainly be an exceptional event, although it could possibly be expected that more diverse marine inclusions, including ones with soft-bodied preservation, would be found in the amber, if this scenario were true. (iii) Being a tropical environment, it could be assumed that tropical storms were fairly common and could therefore blow seashells and sand inland. This could also account for the martesine bivalve shells being found within the amber. However, if this was a fairly common event, it could be expected that occurrences of marine shells in amber would be more common as well.

Marine and terrestrial organisms may get trapped in a single resin piece located at the edge of a coastal forest, and more complicated scenarios such as liquid resin with sea water contact are not needed (6, 11, 54, 55), especially as Schmidt and Dilcher (8) found that resin barely solidifies when submerged in water. The incomplete preservation and lack of the soft body of the marine ammonite (Fig. 2A) and four gastropods (Fig. 4) indicate that they were dead and had experienced abrasion by the sea on the seashore before they were engulfed by resin. Moreover, the aperture of the ammonite is filled with coarse shell sand, which is also present in other parts of the amber piece (Fig. 5A), suggesting that the resin-producing trees were very close to the coast. Therefore, we consider that the first scenario is the most likely. Other marine inclusions in the future may suggest other scenarios, although it is possible for all of the scenarios above to have happened over the lifetime of the amber-producing forest. It seems clear, however, that the forest was living near a dynamic and ever changing coastal environment.

Conclusions

It is rare to find aquatic organisms in amber, and it is extremely rare to find marine organisms in amber, let alone macroscopic marine organisms mixed with intertidal, terrestrial, and potentially freshwater aquatic organisms. The exceptional occurrence of macroscopic marine macrofossils in the resin suggests that the amber forest was growing close to a coast, possibly next to a beach, and could have been subjected to exceptional events. The shells may record an exceptionally high, perhaps storm-generated tide, or even a tsunami or other high-energy event. Alternatively, and more likely, the resin fell to the beach from coastal trees, picking up terrestrial arthropods and beach shells and, exceptionally, surviving the high-energy beach environment to be preserved as amber.

Materials and Methods

The amber piece (BA18100) is deposited in the Lingpog Amber Museum in Shanghai. Photographs were taken using a Zeiss AXIO Zoom.V16 microscope system at the State Key Laboratory of Paleobiology and Stratigraphy, Nanjing Institute of Geology and Paleontology, Chinese Academy of Sciences (NIGPAS). In most cases, incident and transmitted light were used simultaneously. All images are digitally stacked photomicrographic composites of ~40 individual focal planes that were obtained using the software Helicon Focus 6 (<http://www.heliconsoft.com>) for better illustration of the 3D structures, as described by Schmidt et al. (56).

To three-dimensionally reconstruct the ammonite, we scanned the fossil at the micro-CT laboratory of NIGPAS, using a 3D X-ray microscope (3D-XRM), Zeiss Xradia 520 versa. Unlike conventional micro-CT, which relies on maximum geometric magnification and a flat panel detector to achieve high resolution, 3D-XRM uses CCD-based objectives to achieve higher spatial resolution. Based on the size of the fossil specimen, a CCD-based 0.4× objective was used, providing isotropic voxel sizes of 13.36 μm with the help of geometric magnification. During the scan, the acceleration voltage for the X-ray source was 70 kV (current 86 μA), and a thin filter (LE3) was used to avoid beam-hardening artifacts. To improve signal-to-noise ratio, 2001 projections over 360° were collected, and the exposure time for each projection was 3 s. Volume data processing was performed using software

VGStudio Max (version 3.0; Volume Graphics). The Nonplanar Clipping function program (VGStudio version 3.0) was used to reconstruct the suture along the curved surface of the ammonite.

ACKNOWLEDGMENTS. We are grateful to A. R. Schmidt and two reviewers for careful comments. Thanks to Z. Yin and S. Wu for the microcomputed tomography reconstruction and to H. Pan, E. Sidorchuk, N. Bruce, and H. Matuš for assistance with identifications. This research was supported by the Second Tibetan Plateau Scientific Expedition and Research (Grant 2019QZKK0706), Strategic Priority Research Program of the Chinese Academy of Sciences (Grants XDB26000000 and XDA19050101), and National Natural Science Foundation of China (Grants 41622201 and 41688103).

- Martínez-Delciós X, Briggs DEG, Peñalver E (2004) Taphonomy of insects in carbonates and amber. *Palaeogeogr Palaeoclimatol Palaeoecol* 203:19–64.
- Peñalver E, et al. (2012) Thrips pollination of Mesozoic gymnosperms. *Proc Natl Acad Sci USA* 109:8623–8628.
- Briggs DEG (2018) Sampling the insects of the amber forest. *Proc Natl Acad Sci USA* 115:6525–6527.
- Labandeira CC (2014) Amber. *Paleont Soc Pap* 20:163–215.
- Solórzano Kraemer MM, et al. (2018) Arthropods in modern resins reveal if amber accurately recorded forest arthropod communities. *Proc Natl Acad Sci USA* 115:6739–6744.
- Seyfullah LJ, et al. (2018) Production and preservation of resins—past and present. *Biol Rev Camb Philos Soc* 93:1684–1714.
- Girard V, et al. (2008) Evidence for marine microfossils from amber. *Proc Natl Acad Sci USA* 105:17426–17429.
- Schmidt AR, Dilcher DL (2007) Aquatic organisms as amber inclusions and examples from a modern swamp forest. *Proc Natl Acad Sci USA* 104:16581–16585.
- de Lourdes Serrano-Sánchez M, et al. (2015) The aquatic and semiaquatic biota in Miocene amber from the Campo La Granja mine (Chiapas, Mexico): Palaeoenvironmental implications. *J South Am Earth Sci* 62:243–256.
- Matzke-Karasz R, et al. (2019) Abundant assemblage of Ostracoda (Crustacea) in Mexican Miocene amber sheds light on the evolution of the brackish-water tribe Thalassocypridini. *Hist Biol* 31:65–101.
- Schmidt AR, et al. (2018) Marine microorganisms as amber inclusions: Insights from coastal forests of New Caledonia. *Foss Rec (Weinh)* 21:213–221.
- Xing L, et al. (2018) A gigantic marine ostracod (Crustacea: Myodocopa) trapped in mid-Cretaceous Burmese amber. *Sci Rep* 8:1365.
- Grimaldi DA, Ross AJ (2017) Extraordinary Lagerstätten in amber, with particular reference to the Cretaceous of Burma. *Terrestrial Conservation Lagerstätten: Windows into the Evolution of Life on Land*, eds Fraser NC, Sues HD (Dunedin Acad Press Ltd, Edinburgh), pp 287–342.
- Zheng D, et al. (2018) A Late Cretaceous amber biota from central Myanmar. *Nat Commun* 9:3170.
- Ross AJ (2019) Burmese (Myanmar) amber checklist and bibliography 2018. *Palaeoentomology* 2:22–84.
- Cruikshank RD, Ko K (2003) Geology of an amber locality in the Hukawng Valley, northern Myanmar. *J Asian Earth Sci* 21:41–455.
- Scotese CR (2014) *Atlas of Late Cretaceous Paleogeographic Maps, PALEOMAP Atlas for ArcGIS (PALEOMAP Project, Evanston, IL)*, Vol 2.
- Hitzel E (1902) Sur les fossils d'âge Albin recueillis par M. A. Guebard dans la région d'Escagnolles (Alpes-Maritimes). *Bull Soc Geol Fr* 4:874–880.
- Collignon M (1963) *Atlas des fossiles caractéristiques de Madagascar (Ammonites)* (Service Géologique, Tananarive, Madagascar).
- Kennedy WJ, Klinger HC (2012) Cretaceous faunas from Zululand and Natal, South Africa. The desmoceratoid genera *Moretella* Collignon, 1963, *Beudanticeras* Hitzel, 1902, and *Aioloceras* Whitehouse, 1926. *Afr Nat Hist* 8:55–75.
- Pervinquière L (1907) *Études de Paléontologie Tunisienne. 1. Céphalopodes des Terrains Secondaires* (Carte Géologique de la Tunisie, Paris).
- Latil JL (2011) Early Albin ammonites from central Tunisia and adjacent areas of Algeria. *Rev Paleobiol* 30:321–429.
- Matsumoto T (1954) *The Cretaceous System in the Japanese Islands* (The Japanese Soc for the Promotion of Sci Res, Tokyo).
- Crick GC (1907) Cretaceous fossils of Natal. *Third and Final Report of the Geological Survey of Natal and Zululand*, ed Anderson W (Nabu Press, London), pp 161–250.
- Kennedy WJ, Klinger HC (2014) Cretaceous faunas from Zululand and Natal, South Africa. *Valdedorsella*, *Pseudohaploceras*, *Puzosia*, *Bhimaites*, *Pachydesmoceras*, *Parapuzosia* (*Austiniceras*), and *P.* (*Parapuzosia*) of the ammonite subfamily *Puzosiinae* Spath, 1922. *Afr Nat Hist* 10:1–46.
- Kossmat F (1898) Untersuchungen über die Sudindische Kreideformation. *Beitr Paläontol Österreich-Ungarens Oriens* 11:89–152.
- Renz O (1972) Die Gattungen *Puzosia* Bayle, *Bhimaites* Matsumoto und *Desmoceras* Zittel (*Ammonoidea*) im Oberen Albin Venezuela. *Eclogae Geol Helv* 65:701–724.
- Poinar GO (August 14, 2018) A new genus of terrestrial isopods (Crustacea: Oniscidea: Armadillidae) in Myanmar amber. *Hist Biol*, 10.1080/08912963.2018.1509964.
- Broly P, Maillet S, Ross AJ (2015) The first terrestrial isopod (Crustacea: Isopoda: Oniscidea) from Cretaceous Burmese amber of Myanmar. *Cretac Res* 55:220–228.
- Bruce N (2005) New sphaeromatids (Crustacea: Isopoda: Sphaeromatidea) from coastal and freshwater habitats in New Zealand. *Zootaxa* 1002:1–20.
- Jaume D, Queinsec E (2007) A new species of freshwater isopod (Sphaeromatidea: Sphaeromatidae) from an inland karstic stream on Espiritu Santo Island, Vanuatu, southwestern Pacific. *Zootaxa* 1653:41–55.
- Botosaneanu L, Bruce N, Notenboom J (1986) Isopoda: Cirolanidae. *A Faunistic, Distributional and Ecological Synthesis of the World Fauna Inhabiting Subterranean Waters (Including the Marine Interstitial)*, ed Botosaneanu L (Brill Acad Publ, Leiden, The Netherlands), pp 412–422.
- Schmidt C (2008) Phylogeny of the terrestrial Isopoda (Oniscidea): A review. *Arthropod Syst Phylogeny* 66:191–226.
- Sohl NF (1960) Archeogastropoda, Mesogastropoda and stratigraphy of the Ripley, Owl Creek, and Prairie Bluff Formations. *US Geol Surv Prof Pap* 331A:1–151.
- Gulvik M (2007) Mites (Acari) as indicators of soil biodiversity and land use monitoring: A review. *Pol J Ecol* 55:415–440.
- Niedbala W (2012) Ptyctimous mites (Acari, Oribatida) of the Palaearctic region: Distribution. *Fauna Mundi* 5:1–348.
- Webb DP (1977) Regulation of forest litter decomposition by soil arthropods. *The Role of Arthropods in Forest Ecosystems*, ed Matton WJ (Springer, Berlin), pp 57–69.
- Lindo Z, Winchester NN (2007) Oribatid mite communities and foliar litter decomposition in canopy suspended soils and forest floor habitats of western redcedar forests, Vancouver Island, Canada. *Soil Biol Biochem* 39:2957–2966.
- Huhta V, Siira-Pietikäinen A, Penttinen R (2012) Importance of dead wood for soil mite (Acarina) communities in boreal old-growth forests. *Soil Org* 84:499–512.
- Penney D (2006) Fossil oonopid spiders in Cretaceous ambers from Canada and Myanmar. *Palaeontology* 49:229–235.
- Saube EE, et al. (2012) New *Orchestina* Simon, 1882 (Araneae: Oonopidae) from Cretaceous ambers of Spain and France: First spiders described using phase-contrast X-ray synchrotron microtomography. *Palaeontology* 55:127–143.
- Korenko S, Hamouzová K, Pekar S (2014) Trophic niche and predatory behavior of the goblin spider *Triaeris stenaspis* (Oonopidae): A springtail specialist? *J Arachnol* 42:74–78.
- Fannes W, De Bakker D, Loosveldt K, Jocqué R (2008) Estimating the diversity of arboreal oonopid spider assemblages (Araneae, Oonopidae) at Afrotropical sites. *J Arachnol* 36:322–330.
- Gullan PJ, Cranston PS (2010) *The Insects: An Outline of Entomology* (Wiley-Blackwell, West Sussex, UK).
- Shear WA, Kukulová-Peck J (1990) The ecology of Paleozoic terrestrial arthropods: The fossil evidence. *Can J Zool* 68:1807–1834.
- Kime RD, Golovatch SI (2000) Trends in the ecological strategies and evolution of millipedes (Diplopoda). *Biol J Linn Soc Lond* 69:333–349.
- Golovatch SI, Kime RD (2009) Millipede (Diplopoda) distributions: A review. *Soil Org* 81:565–597.
- Shi G, et al. (2012) Age constraint on Burmese amber based on U-Pb dating of zircons. *Cretac Res* 37:155–163.
- Cohen KM, Finney SC, Gibbard PL, Fan J (2013) The ICS International Chronostratigraphic Chart. *Episodes* 36:199–204.
- Smith RDA, Ross AJ (2018) Amberground pholadid bivalve borings and inclusions in Burmese amber: Implications for proximity of resin-producing forests to brackish waters, and the age of the amber. *Earth Environ Sci Trans R Soc Edinburgh* 107:239–247.
- Girard V, et al. (2009) Exceptional preservation of marine diatoms in Upper Albin amber. *Geology* 37:83–86.
- Poinar GO, Lambert JB, Wu Y (2007) Araucarian source of fossiliferous Burmese amber: Spectroscopic and anatomical evidence. *J Bot Res Inst Tex* 1:449–455.
- Dutta S, Mallick M, Kumar K, Mann U, Greenwood PF (2011) Terpenoid composition and botanical affinity of Cretaceous resins from India and Myanmar. *Int J Coal Geol* 85:49–55.
- Perrichot V (2004) Early Cretaceous amber from south-western France: Insight into the Mesozoic litter fauna. *Geol Acta* 2:9–22.
- Perrichot V, Girard V (2009) A unique piece of amber and the complexity of ancient forest ecosystems. *Palaios* 24:137–139.
- Schmidt AR, et al. (2010) Cretaceous African life captured in amber. *Proc Natl Acad Sci USA* 107:7329–7334.