

Review

Clinical review: Checklists – translating evidence into practice

Bradford D Winters¹, Ayse P Gurses², Harold Lehmann³, J Bryan Sexton^{2,4},
Carlyle Jai Rampersad⁵ and Peter J Pronovost^{2,4}

¹Departments of Anesthesiology and Critical Care Medicine, The Johns Hopkins University, 600 N. Wolfe Street, Meyer 299, Baltimore, MD 21287, USA

²Departments of Anesthesiology and Critical Care Medicine, The Johns Hopkins University, 1909 Thames Street, 1st floor, Baltimore, MD 21231, USA

³Departments of Pediatrics and Welch Health Sciences Informatics, The Johns Hopkins University, 2024 E. Monument Street, 1-201, Baltimore, MD 21205, USA

⁴Department of Health Policy and Management, The Johns Hopkins University, 1909 Thames Street, 2nd floor, Baltimore, MD 21231, USA

⁵US Airways, Philadelphia International Airport, Terminal B, Philadelphia, PA 19153, USA

Corresponding author: Peter J Pronovost, ppronovo@jhmi.edu

Published: 31 December 2009

This article is online at <http://ccforum.com/content/13/6/210>

© 2009 BioMed Central Ltd

Critical Care 2009, **13**:210 (doi:10.1186/cc7792)

Abstract

Checklists are common tools used in many industries. Unfortunately, their adoption in the field of medicine has been limited to equipment operations or part of specific algorithms. Yet they have tremendous potential to improve patient outcomes by democratizing knowledge and helping ensure that all patients receive evidence-based best practices and safe high-quality care. Checklist adoption has been slowed by a variety of factors, including provider resistance, delays in knowledge dissemination and integration, limited methodology to guide development and maintenance, and lack of effective technical strategies to make them available and easy to use. In this article, we explore some of the principles and possible strategies to further develop and encourage the implementation of checklists into medical practice. We describe different types of checklists using examples and explore the benefits they offer to improve care. We suggest methods to create checklists and offer suggestions for how we might apply them, using some examples from our own experience, and finally, offer some possible directions for future research.

Introduction

Checklists are common cognitive tools that can help complete a task as simple as shopping or as complex as flying a Boeing 737. The complexity and the time required for completing tasks and the context in which the work occurs vary widely. Yet all tasks are prone to human error given the natural limitations of our memory and attention span and our ability to cope with stress, fatigue, illness, interruptions, new situations, and production pressures. Intensive care units (ICUs) are complex and time-sensitive. On any given day in the ICU, the typical patient will require 178 interactions in their care [1]. A checklist standardizes the process to ensure that all elements or actions are addressed. The structure and predictability of checklists facilitate the careful and systematic

delivery of care, which reduces variability and improves performance. The science of developing checklists in health care is new. In an informal review of the literature, we did not find any standardized methodology to develop and design checklists in medicine [2]. Based on a review of checklist development in aviation and our own research and experience, we propose a process for creating checklists. If medical checklist elements are wise and based on best practices, they can help translate evidence into practice.

One great advantage of a checklist is the ability to democratize knowledge. Medicine, like many fields, is infused with jargon; physicians, nurses, and patients have distinct ways of describing the same thing. As such, translation errors are imminent when one member of the care team talks to another. Given that each caregiver has a unique perspective and professional culture and that clinical disciplines train separately, it is understandable why miscommunication is common and a major contributor of medical errors [3-6]. Checklists reduce these risks. They standardize and improve the reliable translation of information so the same knowledge is available to doctors, nurses, and patients.

Although checklists have tremendous potential to improve safety and quality and reduce the costs of health care, they are underused. Because we lack a clear definition for a medical checklist, it is difficult to know how widely they are used in health care. To be certain, other efforts to standardize work and create independent checks, such as protocols, critical pathways, and order sets, exist in health care. Yet they differ in significant ways from the quality/safety checklists we describe. Order sets are pre-printed instructions that attempt

CLABSI = central line-associated bloodstream infection; CPOE = computerized provider order entry; HCAP = health care-associated pneumonia; ICU = intensive care unit; OR = operating room; VAP = ventilator-associated pneumonia.

to include everything that a clinician may want to consider ordering for a particular group of patients (for example, those on anticoagulation therapy). Many things on this order set will not be checked. Conversely, quality/safety checklists include only the things that should be addressed. Clinical/critical pathways are time-continuum tools that support clinical decision-making and function primarily to improve work flow for a specific diagnosis. Pathways have nodes with drop-down checklists of tasks along their continuum. However, many pathways do not include explicit details regarding patient-level interventions. Rather, they generally present expected milestones along a patient care trajectory. Nevertheless, both order sets and pathways can be viewed as a type of checklist.

The underuse of quality/safety checklists is partly due to the paucity of scholarly research to identify where to use checklists, how to build and implement them and assess their effectiveness at improving patient outcomes, and whether or how checklist use is sustained over time [2,7]. In this review, we explore some of the principles and possible strategies to develop and encourage the implementation of checklists in medical practice.

Types of checklists

There are four principal types of checklists: static parallel, static sequential with verification, static sequential with verification and confirmation, and dynamic [8]. The differences among these checklists are the number of operators involved and the extent to which the correct action is verified. Static parallel checklists are completed by one operator (person) and executed as a series of read-and-do tasks. The anesthesia machine checklist (Figure 1) is an example of this format [9].

The static sequential with verification checklist involves a challenge and response. An operator reads a series of items and a second person verifies that each item has been completed or is within parameters. A checklist to ensure appropriate insertion of central lines is an example of this format [10,11]. As the physician performs the insertion, a nurse challenges the completion of each behavior/task and the physician responds to confirm whether the behavior/task has been accomplished.

Static sequential with verification and confirmation checklists are used more often in team-based settings where a set of items/tasks are done by varying team members. A designated person reads the items (challenge) and each responsible party verifies the completion of their specific task. Time-outs and briefings in the operating room (OR) use this format [12]. For example, the surgeon will state the patient's name, procedure, and surgical site and ask about availability of equipment (to which the nurse would confirm the information) and then ask about the patient's medical condition and availability of blood (to which the anesthesiologist would respond).

Finally, a dynamic checklist uses a flowchart to guide complex decision-making. Typically, there are multiple options to choose from and the care team must decide the optimal course (an algorithm). An example of a dynamic checklist is the American Society of Anesthesiologists difficult airway algorithm [13], which provides guidance on the intubation of a difficult airway. The team leader would use this algorithm checklist to develop a plan and then discuss it with other members of the team.

Checklists may be further categorized by their application to high-risk or normal operations. High-risk checklists are useful back-up plans to fix or mitigate harm when single or multiple failures of redundant systems occur [14]. High-risk checklists can be used during a crisis to prevent further mistakes and to ensure reliable communication and operations. For example, their use during disaster relief efforts would standardize tasks and processes, thereby organizing different groups. Pre-operative preparation of patients is an example of a normal operations checklist.

Structure of checklists

The structure of a checklist may or may not be important, depending on the context in which they are used. For some checklists, the sequential execution of each item is critical. For others, the focus is solely on executing or considering the items. When sequential action is required, an independent check is crucial to ensure that each task is completed prior to executing the next task. A static checklist would be risky here, particularly in high-risk situations given our natural cognitive and stress-induced limitations [15].

Failure to properly follow checklists (sequential or not) can have disastrous consequences [16]. For example, if the proper checklist procedure for identifying and verifying a patient and their blood type is not followed, the unit of blood transfused may be incompatible and lead to a potentially lethal hemolytic transfusion reaction.

Adoption and benefits of checklists

In medicine, physicians have largely resisted using checklists. Some feel that relying on a checklist insults their intelligence, whereas others doubt that a document with check boxes will prevent a medical mistake [1]. Physicians believe they know their job and do not need a prompter to guide or remind them [17]. However, modern medicine has become exceedingly complex, specialized, and interdisciplinary, offering hope for fantastic cures, but also inadvertently introducing potentially devastating risks.

Well-designed checklists standardize what, when, how, and by whom interventions are done and can reduce errors in routine and emergency situations. In addition, they provide a public framework to ensure adherence to clinical or procedural requirements. The shared knowledge of checklist content also allows caregivers to mutually support each other

Figure 1

NAME:	GRADE:
DATE:	TIME STARTED:
TIME COMPLETED:	
THEATER:	ANESTHETIC MACHINE: Anesthetic room/Theater

A. ANESTHETIC MACHINE
Note any labelling or service information attached to machine. Switch on electrical supply (if appropriate).

B. OXYGEN ANALYZER
Switch on.

1. Is analyzer calibrated?	Yes/No
2. Is analyzer functioning correctly?	Yes/No

Attach to common gas outlet.

C. MEDICAL GAS SUPPLIES
Identify gases supplied by pipeline and confirm correct connections with 'tug-test'.

1. Is machine connected to an O ₂ supply?	Yes/No
--	--------

Switch on spare O₂ cylinder.

2. Are cylinder contents adequate?	Yes/No
3. Is machine connected to N ₂ O supply (if intended for use)?	Yes/No
4. Are contents of spare N ₂ O cylinder adequate?	Yes/No
5. Is machine connected to compressed air supply (if intended for use)?	Yes/No
6. Are contents of spare air cylinder adequate?	Yes/No
7. Is CO ₂ cylinder attached to machine?	Yes/No
8. If yes, have you removed it?	Yes/No
9. Are blanking plugs fitted to all empty cylinder yokes?	Yes/No

D. FLOWMETERS

1. Do all flowmeter bobbins move freely throughout their range?	Yes/No
2. With O ₂ flowing at 5 L/min, does O ₂ analyzer approach 100%?	Yes/No

Turn off all flowmeters.

E. EMERGENCY OXYGEN BYPASS CONTROL

1. When the O ₂ bypass control is operated, does flow occur without significant drop in pipeline pressure?	Yes/No
2. Does O ₂ analyzer approach 100%?	Yes/No
3. Does flow cease when control is released?	Yes/No

F. VAPORIZERS

1. Are vaporizers for the required volatile agents present, correctly seated, and locked to the back-bar?	Yes/No
2. Are the vaporizers adequately filled?	Yes/No
3. Are the filling ports tightly closed?	Yes/No
4. Does the control knob for each vaporizer move through the full range?	Yes/No

Only perform the following tests where the back-bar is protected by a pressure relief valve:
With O₂ flow of 5 L/min, occlude common gas outlet.

5. Does flowmeter bobbin dip?	Yes/No
-------------------------------	--------

Turn on each vaporizer in turn and briefly occlude the common gas outlet.

6. Do any leaks occur from the filling ports of the vaporizers?	Yes/No
---	--------

Turn off all vaporizers.

G. BREATHING SYSTEM

1. Is breathing system correctly assembled, with all connections tight?	Yes/No
2. Do any leaks occur when the system is pressurized?	Yes/No
3. Does the adjustable pressure relief valve open and close fully?	Yes/No
4. In a circle system, do the unidirectional valves move correctly?	Yes/No

H. VENTILATOR

1. Is ventilator correctly assembled with all connections tight?	Yes/No
--	--------

Set the controls for use and switch on.

2. Is adequate pressure generated during the inspiratory phase?	Yes/No
3. Does the pressure relief valve operate correctly when the patient port is occluded?	Yes/No
4. Is the disconnection alarm present and operating correctly?	Yes/No
5. Is alternative means of ventilation available?	Yes/No

I. SCAVENGING

1. Is scavenging system correctly attached and functioning?	Yes/No
---	--------

J. ANCILLARY EQUIPMENT
Confirm presence, size range, and function of all ancillary equipment that may be needed.

1. Are all laryngoscopes in working order?	Yes/No
2. Is suction apparatus present and able to generate adequate negative pressure rapidly?	Yes/No
3. Does patient trolley tip head-down?	Yes/No

K. MONITORING EQUIPMENT
Check that appropriate monitoring equipment is present, switched on, and calibrated.
Set alarm limits as appropriate.

Checklist for anesthetic machines [9].

by cross-checking what is being done and in what order. These assurances are important when time is short, the pressure is on, and competing priorities distract our attention.

Checklists have been implemented in isolated clinical settings to improve processes of care [11,18-25] and in the ICU to facilitate bedside teaching and assessment of resident performance [26]. Checklists can be implemented as a stand-alone intervention [22], but in most cases they are part of an intervention bundle with several other components to improve quality and safety of care [11,27]. Studies to improve care using checklists have reduced uncertainty over the correct surgical site [20] and problems with laparoscopic equipment in the OR [21], clarified [22] or changed patient care plans [23] to mitigate or prevent medical errors, dramatically reduced central line-associated bloodstream infections (CLABSIs) [11], and helped diagnose communication deficiencies and depression in adults with intellectual disabilities [24,25]. Despite the corpus of evidence regarding the benefits of checklists, medicine remains slow in broadly adopting them into practice.

Seminal work in the psychology of memory found that we can retrieve seven (plus or minus two) pieces of information from our memory with relative accuracy [28]. When complex procedures, stress, and fatigue are introduced, our memory becomes increasingly unreliable [29]. And as the number of tasks/problems we simultaneously manage exceeds three, we show significant decline in the accuracy and speed of handling these tasks/problems [30]. A checklist can compensate for these fallibilities. The mandatory use of checklists can empower nurses and junior-level physicians to insist that those higher in the traditional hierarchy adhere to approved and safe procedures.

Although there is no published evidence indicating a negative impact by using checklists, they could pose risks. Any time that we change the system to improve safety, we may defend against some risks but invariably will introduce new risks. Checklists are not immune to this tendency. Poorly designed checklists or excessive use of checklists could overburden clinicians, complicate tasks, and reduce efficiency. If emerging evidence is not incorporated into checklists, they could hinder patients from getting state-of-the-art care. Another potentially negative effect could occur if clinicians strictly adhere to a checklist rather than exercise critical thinking when evidence is incomplete, when an individual patient's risk-benefit analysis favors not using the checklist, or when an unforeseen event that requires different interventions occurs. A recent example of this last point is the pilot who landed the Airbus A320 plane on the Hudson River without fatalities. Hence, it is imperative to be cautious when deciding when to use checklists and to be mindful of potential negative effects.

Creating checklists

The literature review of medical checklists by Hales and colleagues [2] found few strategies to develop checklists and


no standardized methodologies. Ideally, the process would incorporate diverse input (for example, empiric evidence, tacit experience, local input, regulatory input, and community input). Unfortunately, the process is often delegated to committees with homogeneous members, who may lack the experience and knowledge to explore appropriate items for checklists and the potential risks of these items. Checklists with elements that pose risks or those that exclude important elements may be neither effective nor efficient at improving patient care. While we need investment in the 'basic science' of checklist development, the human factors engineering literature describes how to design information tools, which can be applied in quality and safety. Briefly, the recommended steps to develop checklists include the following: review the existing literature, understand the needs and workplace of the users, include a multidisciplinary group in the design, and use an iterative approach for rigorous pilot testing and validation of your tool [31-33]. We recently published an approach to develop checklists which adds simulation to the above steps [34]. Below, we provide more detailed guidance to develop checklists.

First, we must pick a patient population, procedure, and/or outcome. For example, we sought to reduce infections from central lines in ICU patients [10]. This could include checklists for diagnosing, treating, or monitoring patients. Then, we need to assemble an interdisciplinary team to lead the development. This team should review and identify empiric evidence from published literature and tacit evidence from experience to decide on the content of the checklist (Figure 2).

In identifying evidence, the team could start by looking for evidence summaries (for example, systematic reviews or practice guidelines). For most items, however, empiric evidence will be either absent or incomplete. As such, teams should tap into the 'wisdom of crowds' by getting broad input regarding checklist items from diverse sources [35]. In addition to literature database sources such as PubMed, Embase, and Cochrane reviews, teams should consider knowledge banks, expert and social networking, and focus groups with frontline practitioners. Once a list of potential interventions is compiled, the team should consider those with the strongest impact and the lowest barriers to use in clinical practice. Because role and/or task ambiguity is associated with failure to use the checklist as intended [36], each intervention should be translated into an explicit, concise, and unambiguous behavior. This culling process is critical to make the checklist cognitively and logistically functional. In the case of health care-associated pneumonia (HCAP), there are over a hundred potential steps to prevent this complication, with multiple tasks for each step. Without discipline and data reduction, developing an HCAP prevention or similar checklist could become unwieldy.

In our experience, long unorganized checklists are often not useful or used. As previously mentioned, our ability to manage

Figure 2



Elements of creating a wise checklist.

information deteriorates as the number of items increases [28,30,37]. Memory checklists should especially adhere to this principle. If a checklist requires a large number of items, it is better to separate the process into substeps and create a checklist for each substep. In the OR, for example, there are critical steps that could be broken down into individual checklists, such as preventing wrong-sided surgery and surgical site infection. Each of these is part of the pre-induction/incision process but involves distinct tasks and behaviors. The creation process should be iterative and undergo as many revisions as necessary to achieve consensus on the checklist before attempting any pilot testing.

The checklist should provide unambiguous guidance on what, when, how, and who should do a particular intervention and should be logistically efficient and easily performed. The aviation industry offers guidance. Figure 3 shows a standard aviation procedure checklist for landing a Boeing 737-800 aircraft. This checklist uses the challenge/response format with simple, direct, and unambiguous language. Health care checklists should seek to emulate this model. The anesthesia machine static 'read-and-do' checklist (Figure 1) [9] is a close example of the aviation model with concise questions.

Once a beta-version of a checklist is completed, it should be pilot-tested by potential users and revised according to their findings. This testing could be done on the clinical units where it will be used and/or in a simulation setting. Usability and potential risks of a checklist can be assessed by (a) testing different real-life scenarios in a simulated environment, (b) heuristic evaluation by a human factors/usability testing expert, and (c) subjective evaluation by potential users through interviews, focus groups, and surveys [38]. Validity, reliability, and potential harms of a checklist should be assessed before it is broadly implemented. Face validity can be accomplished by asking potential users to review the tool for relevance and completeness and to suggest how it can be improved. Theoretical saturation (collecting data from

additional people until no new information emerges) [39] can be used to determine the number of people needed to pilot-test a checklist. Reliability of a checklist can be assessed using inter-rater and intra-rater reliability methods.

Finally, checklists must remain wise. To accomplish this, checklists must be dynamic and evolve, and empiric and tacit evidence must be continually evaluated to support each checklist item and to explore unintended consequences that are the norm rather than the exception. If this is accomplished, checklists can be efficient and effective.

When creating checklists, developers could use the following principles applied in human factors engineering [7]:

1. Design checklists based on caregivers' needs and the realities of their work by doing ethnographic studies of the clinical work and involvement of potential users.
2. List the most critical items at the beginning of the checklist whenever possible.
3. Avoid long checklists if possible. Subdivide long checklists into small meaningful sections or group checklists in time and space (for example, one checklist for this moment in time).
4. Pay close attention to usability, including the time it takes to complete the checklist, potential negative effects on caregivers' work and patient safety, and feedback from potential users.
5. Perform rigorous pilot testing and validation of the checklist before full-scale implementation.
6. Include potential users, content experts, and human factors/usability experts on the design team.
7. Re-evaluate and update checklists periodically based on new literature and organizational experiences.

Applying checklists

First, we will consider several widely used checklists in health care. The anesthesia machine checklist is universally accepted by anesthesiologists [40,41] and, over the course of

several decades of use, has undergone modifications to keep pace with new technology. This checklist helps to ensure that every practitioner's equipment has undergone the same standardized assessment (although machines of individual manufacturers may vary somewhat) before transporting the patient to the OR. Like many aviation checklists, this one focuses on a particular piece of equipment. One practitioner completes the checklist and the effectiveness depends on strict adherence to its completion [16].

We developed the catheter insertion checklist to address the problem of CLABSIs in the ICU [10]. It was informed by a detailed practice guideline developed by the Centers for Disease Control and Prevention (Atlanta, GA, USA) [42]. From this guideline, the team selected five practices that had the strongest evidence and lowest barriers to use. Each of the interventions was worded as a behavior, and we pilot-tested the checklist in our ICUs at the Johns Hopkins Hospital. This checklist uses the static sequential with verification format. The 'challenger' (nurse) is empowered to prevent the procedure from proceeding until the physician inserting the catheter confirms that the checklist item was completed. This empowerment of the nurse to halt the process makes the checklist particularly robust. Nevertheless, it requires a culture in which nurses are comfortable questioning physicians and in which physicians are willing to accept and modify their behavior. The successful use of interdisciplinary checklists such as this one often requires efforts to improve the culture of safety [43].

Another checklist we developed was a daily goals sheet (Figure 4) [22]. This checklist was built after residents and nurses on the ICU health care team stated that communication errors posed substantial risks to patients. It clarifies the patient's daily care plan, helps ensure delivery of evidence-based interventions, and plans for potential safety risks. It follows a static sequential with verification format. At the end of rounds on a patient, a member of the team reads off the items and team members (including the nurse) respond with the status of those items and discuss the patient's goals. Items that need to be changed, deleted, or added become the tasks for the day. The checklist is completed for every patient to ensure uniformity of care delivery and to ensure that all care team members understand the patient's status and care plan prior to moving to the next patient.

The daily goals checklist is fairly complex and has many items. To reduce the burden on clinicians, items are grouped into categories consistent with how ICU clinicians think. Each category contains a limited number of items. It was decided to keep this list intact since the checklist is kept by the bedside and revisited throughout the day.

In addition to addressing specific problems and goals, checklists may be applied to the whole spectrum of the care process. While acknowledging the complexity of this

Figure 3

Landing gear	<input type="checkbox"/> Check down
Autopilot	<input type="checkbox"/> Off
Landing speed	<input type="checkbox"/> 140 KIAS
After touchdown	<input type="checkbox"/> Apply reverse thrust
	<input type="checkbox"/> 60 KIAS: cancel reverse thrust
Spoilers	<input type="checkbox"/> Verify extended
Brakes	<input type="checkbox"/> As required

Landing procedure checklist for a Boeing 737-800 aircraft. Adapted from the Atlantic Sun Airways CAT B pilot procedures and checklists series [52]. KIAS, knots indicated airspeed.

undertaking, the process of medicine can be divided into three major phases: diagnosing, treating, and monitoring. Each phase can be broken down into three elements: decision (whether and what to do), execution (carrying the process out), and interpretation (what is the result and what does it mean). Medical errors may occur anywhere along this hypothetical spectrum of patient care. As such, checklists can help improve care anywhere along this spectrum. For example, we are developing checklists (from the patient's perspective) for diagnosing, treating, and monitoring lung cancer. The use of these checklists should help to reduce variability and errors in patient care.

Ventilator-associated pneumonia (VAP) is a useful model to examine how checklists can improve a spectrum of care. The diagnostic criteria and the recommended therapies for preventing and treating VAP are well established. A section on our daily goals form includes a checklist of evidence-based prevention interventions. The weaning item on the checklist prompts clinicians to monitor and wean the patient as soon as safely possible. This is followed by a separate checklist to evaluate the patient's readiness for extubation. This approach has shortened the average duration of mechanical ventilation and reduced ventilator days, failed extubations, and VAP rates [44].

The *diagnosis* phase checklist would cognitively prompt us to recognize the possibility of a VAP and then execute the entire process to interpret whether the patient meets the criteria for VAP. It would also assist in deciding whether to treat the patient. A *treatment* phase checklist would include therapies based on evidence-based medicine guidelines, local knowledge of probable pathogens, and anti-microbial sensitivity patterns. In addition, there may be a policy or structural-level checklist for preventing VAP (such as not routinely changing ventilator circuits). A *monitoring* checklist can help to

Figure 4

Room number _____ MD/NP COVERING Pt today: _____ Date ____/____/____ (Protocols available if **bolded**)

		AM shift (7 a.m.)	PM shift (7 p.m.) **Note changes from AM**
Safety	What needs to be done for patient to be discharged from the ICU?		
	<ul style="list-style-type: none"> ▪ Patient's greatest safety risk? ▪ How can we decrease risk? 		
	What events or deviations need to be reported? PSNs?	<input type="checkbox"/> PSN	
Patient care	IMPRESSION/PLAN	<input type="checkbox"/> Better <input type="checkbox"/> Unchanged <input type="checkbox"/> Worse	
	NEURO Pain mgt / Sedation	Pain goal ____/10 w/ <input type="checkbox"/> Daily lightening of sedation	<input type="checkbox"/> Wean sedation for extubation in AM
	RESP <ul style="list-style-type: none"> ▪ Pulmonary: ▪ Ventilator: (vent bundle: HOB elevated), RTW/Weaning) 	<input type="checkbox"/> OOB/ pulm toilet/ambulation <input type="checkbox"/> Maintain current support <input type="checkbox"/> FIO ₂ < ____ PEEP < ____ <input type="checkbox"/> Wean as tol <input type="checkbox"/> Swallow eval <input type="checkbox"/> PS/Trach trial ____h x ____ <input type="checkbox"/> Mechs before/after	<input type="checkbox"/> Wean vent <input type="checkbox"/> Mechanics by __a.m. <input type="checkbox"/> Plan to extubate
	CARD Cardiac Review EKGs	HR goal ____ <input type="checkbox"/> at goal <input type="checkbox"/> ↑ <input type="checkbox"/> ↓ <input type="checkbox"/> β block ____	
	GI GI / Nutrition / Bowel regimen (TPN line, NDT, PEG needed?)	<input type="checkbox"/> NPO <input type="checkbox"/> TF type ____ goal ____ <input type="checkbox"/> TPN INSULIN REQ ____Adj needed y/n	
	GU Volume status Net goal for midnight	<input type="checkbox"/> Net even <input type="checkbox"/> Net positive <input type="checkbox"/> Net neg: ____ w/ ____ <input type="checkbox"/> Pt determined	
	ID SIRS/infection/sepsis Evaluation SIRS criteria <input type="checkbox"/> Temp. >38°C or <36°C <ul style="list-style-type: none"> <input type="checkbox"/> HR >90 bpm <input type="checkbox"/> RR >20 breaths/min or PaCO₂ <32 torr <input type="checkbox"/> WBC >12K <4K or >10% bands	<input type="checkbox"/> No current SIRS/sepsis issues <input type="checkbox"/> Known/suspected infection: <input type="checkbox"/> PAN Cx <input type="checkbox"/> Bld x2 <input type="checkbox"/> Urine <input type="checkbox"/> Sputum <input type="checkbox"/> Other <input type="checkbox"/> ABx changes: Initiate / D/C <input type="checkbox"/> AG levels: <input type="checkbox"/> Sepsis bundle	
	Can catheters/tubes/lines be removed/rewired?	<input type="checkbox"/> Y <input type="checkbox"/> N	
	Is this patient receiving DVT/PUD prophylaxis? Is the patient receiving PT/OT/ROM	DVT: <input type="checkbox"/> Hep q8 / q12 / gtt (protocol?) PUD: <input type="checkbox"/> PPI <input type="checkbox"/> TEDS/SCDs <input type="checkbox"/> H ₂ B <input type="checkbox"/> LMWH PT/OT <input type="checkbox"/> ROM <input type="checkbox"/> Contraindication if any <input type="checkbox"/>	
	Anticipated LOS >2 days: TGC 3 days: fluconazole PO, KCl SS	<input type="checkbox"/> TGC <input type="checkbox"/> Fluc <input type="checkbox"/> KCl <input type="checkbox"/> N/A	<input type="checkbox"/> Transition from TGC→ SSI by AM
	Can any meds be discontinued, converted to PO, adjusted?	<input type="checkbox"/> N/A <input type="checkbox"/> D/C: <input type="checkbox"/> PO: <input type="checkbox"/> Renal: <input type="checkbox"/> Liver:	
	Tests / Procedures / OR today	<input type="checkbox"/> N/A <input type="checkbox"/> Consents needed/obtained	<input type="checkbox"/> Line change
	Scheduled labs (Reassess need q12 hrs) <i>AM lab needed</i> CXR?	<input type="checkbox"/> N/A <input type="checkbox"/> CMP <input type="checkbox"/> BMP <input type="checkbox"/> H8 <input type="checkbox"/> Coags <input type="checkbox"/> ABG <input type="checkbox"/> Lactate <input type="checkbox"/> Core 4 <input type="checkbox"/> CXR Wed: <input type="checkbox"/> Transferrin <input type="checkbox"/> Iron <input type="checkbox"/> Prealb <input type="checkbox"/> 24-hr urine	
Consultations	<input type="checkbox"/> Y <input type="checkbox"/> N		
Disposition	Is the primary service up-to-date?	<input type="checkbox"/> Y <input type="checkbox"/> N	

Daily goals checklist. ABG, arterial blood gas; ABx, antibiotics; AG, aminoglycoside; Bld, blood; BMP, basic metabolic panel; bpm, beats per minute; Card, cardiology; CMP, comprehensive metabolic profile; CXR, chest x-ray; D/C, discontinue; DVT, deep venous thrombosis; EKG, electrocardiogram; eval, evaluation; FIO₂, fraction of inspired oxygen; Fluc, fluconazole; GI, gastrointestinal; gtt, drops; GU, genitourinary; Hep, heparin; HOB, head of bed; HR, heart rate; ICU, intensive care unit; ID, infectious disease; KCl SS, potassium chloride sliding scale; LMWH, low-molecular-weight heparin; LOS, length of stay; mgt, management; NDT, nasoduodenal tube; Neuro, neurology; NPO, *nil per os* (nothing by mouth); OOB, out of bed; OR, operating room; OT, occupational therapy; PaCO₂, arterial partial pressure of carbon dioxide; PAN Cx, pan-culture; PEEP, positive end-expiratory pressure; PEG, percutaneous endoscopic gastronomy; PO, *per os* (by mouth); PPI, proton pump inhibitor; Prealb, prealbumin; PS, pressure support; PSN, patient safety network; Pt, patient; PT, physical therapy; PUD, peptic ulcer disease; pulm, pulmonary; q, every; Resp, respiratory; ROM, range of motion; RR, respiratory rate; RTW, ready to wean; SCD, sequential compression device; SIRS, systemic inflammatory response syndrome; SSI, sliding scale insulin; TEDS, thromboembolic deterrent stockings; TF, tube feeding; TGC, tight glucose control; tol, tolerated; TPN, total parenteral nutrition; WBC, white blood cells.

determine whether the patient requires continued ICU care, monitoring of oxygen saturation, or additional respiratory care.

This seems like a lot of checklists. Yet each step in the diagnosis, treatment, and monitoring process poses risks for error that we need to defend against. We do not know how many checklists are too many, when they are most useful, and when we have overloaded the checklist users. However, we do know that checklists should be short and simple, and the benefits demonstrated rather than assumed. In aviation, processes are broken down into manageable segments, each with an assigned checklist. This model has been successful in preventing errors. Although there may be some redundancy from this design, redundancy is a crucial element of safe and reliable systems [45].

One final example of a medical checklist that underscores this redundancy is decision support tools built into computerized provider order entry (CPOE) systems. While implementation of CPOE may have difficulties and unintended consequences [46,47], smooth and effective implementation of these systems can be enhanced by building a large number of order sets (checklists) for therapies and management. The more order sets created, the more efficient and effective the CPOE system [48,49]. Information technology will likely be the best mechanism to manage a large number of checklists. As such, we look forward to the day when our checklists are all automated and the required checklist can be selected electronically.

Future directions: create a more efficient health care knowledge market

While the science of checklists, much like the science of safety and quality, is immature, many believe that medical checklists can help prevent errors, mitigate harm, and reduce the costs associated with them. Yet using the process we describe to develop checklists for the multitude of diagnoses and procedures is a daunting and likely slow process. When we reflect on the underlying problem in translating evidence into practice, like many health care quality and safety shortcomings, it is clearly impaired by an inefficient knowledge market, much like the subprime credit crisis. Lack of diverse input and timely access to such information contributes to this inefficiency.

The delay between knowledge creation, dissemination, and translation into practice is significant – sometimes taking years [50,51]. Clearly, there are many elements to this process which must be addressed. Nevertheless, more efficient knowledge markets that quickly synthesize diverse input into usable tools and make these tools widely accessible could substantially improve some of the delays in knowledge translation.

A strategy to improve the efficiency and effectiveness of the health care knowledge market would be to create broad

learning communities to continually create and update checklists. Clinicians organized around a particular diagnosis or procedure can create checklists for a particular disease or process. These checklists would be informed by empiric and tacit evidence and revisited often to incorporate new evidence. If we believe in the ‘wisdom of crowds’, in which the broader community has tremendous knowledge and collectively will get it right, then this learning community can effectively codify this wisdom into checklists that can improve patient care.

Conclusions

Checklists are powerful tools to standardize work processes and create independent checks for key processes. Although they can have wide application in medicine, they are relatively underused. Checklists could create a more efficient and effective knowledge market by summarizing evidence into explicit behaviors, incorporating empiric and tacit evidence, and being continually updated by the health care community.

Further research is needed to advance the science for developing, implementing, and evaluating checklists. Although their format and content may vary, simple steps to identify, check, and verify what you have done or are about to do can determine whether you succeed or fail. They should be succinct, unambiguous, focused, and ultimately effective and efficient. When faced with a crisis, we can react quickly and decisively, knowing that the items we act out from the checklist are well thought out, tested, and will provide us with the results we want. We hope the science advances to help us build, refine, and use checklists wisely.

Competing interests

The authors declare that they have no competing interests.

Acknowledgments

The authors wish to thank Christine G Holzmueller for her assistance in editing this manuscript; no funding was provided. The authors had no source of funding for this research.

References

1. Gawande A: **The checklist: if something so simple can transform intensive care, what else can it do?** *New Yorker*. 10 December 2007; *Annals of Medicine* section:1-8.
2. Hales B, Terblanche M, Fowler R, Sibbald W: **Development of medical checklists for improved quality of patient care.** *Int J Qual Health Care* 2008, **20**:22-30.
3. Pronovost PJ, Thompson DA, Holzmueller CG, Lubomski LH, Dorman T, Dickman F, Fahey M, Steinwachs DM, Engineer L, Sexton JB, Wu AW, Morlock LL: **Toward learning from patient safety reporting systems.** *J Crit Care* 2006, **21**:305-315.
4. Lingard L, Espin S, Whyte S, Regehr G, Baker GR, Reznick R, Bohnen J, Orser B, Doran D, Grober E: **Communication failures in the operating room: an observational classification of recurrent types and effects.** *Qual Saf Health Care* 2004, **13**:330-334.
5. Sutcliffe KM, Lewton E, Rosenthal MM: **Communication failures: an insidious contributor to medical mishaps.** *Acad Med* 2004, **79**:186-194.
6. Gawande AA, Zinner MJ, Studdert DM, Brennan TA: **Analysis of errors reported by surgeons at three teaching hospitals.** *Surgery* 2003, **133**:614-621.
7. Degani A, Wiener EL: **Cockpit checklists: Concepts, design, and use.** *Hum Factors* 1993, **35**:345-359.

8. Peleg M, Boxwala AA, Ogunyemi O, Zeng Q, Tu S, Lacson R, Bernstam E, Ash N, Mork P, Ohno-Machado L, Shortliffe EH, Greenes RA: **GLIF3: the evolution of a guideline representation format.** *Proc AMIA Symp* 2000;645-649.
9. Kendall J, Barthram C: **Revised checklist for anaesthetic machines.** *Anaesthesia* 1998, **53**:887-890.
10. Berenholtz SM, Pronovost PJ, Lipsett PA, Hobson D, Earsing K, Farley JE, Milanovich S, Garrett-Mayer E, Winters BD, Rubin HR, Dorman T, Perl TM: **Eliminating catheter-related bloodstream infections in the intensive care unit.** *Crit Care Med* 2004, **32**:2014-2020.
11. Pronovost P, Needham D, Berenholtz S, Sinopoli D, Chu H, Cosgrove S, Sexton B, Hyzy R, Welsh R, Roth G, Bander J, Kepros J, Goeschel C: **An intervention to decrease catheter-related bloodstream infections in the ICU.** *N Engl J Med* 2006, **355**:2725-2732.
12. Makary MA, Holzmüller CG, Thompson DA, Rowen L, Heitmiller ES, Maley WR, Black JH, Stegner K, Freischlag JA, Ulatowski JA, Pronovost PJ: **Operating room briefings: working on the same page.** *Jt Comm J Qual Saf* 2006, **32**:351-355.
13. American Society of Anesthesiologists homepage [<http://www.asahq.org>].
14. Burian B: **Aeronautical emergency and abnormal checklists: expectations and realities.** In *Proceedings of the Human Factors and Ergonomics Society 50th Annual Meeting: 16-20 October 2006; San Francisco*. Santa Monica, CA: Human Factors and Ergonomics Society; 2006:101-105.
15. Swain AD, Guttman HE: *Handbook of Human Reliability Analysis with Emphasis on Nuclear Power Plant Applications (NUREG/CR-1278)*. Washington, DC: Nuclear Regulatory Commission; 1983.
16. Hayashi I, Wakisaka M, Ookata N, Fujiwara M, Odashiro M: **Actual conditions of the check system for the anesthesia machine before anesthesia. Do you really check?** *Masui* 2007, **56**:1182-1185.
17. Colon-Emeric CS, Lekan D, Utley-Smith Q, Ammarell N, Bailey D, Corazzini K, Piven ML, Anderson RA: **Barriers to and facilitators of clinical practice guideline use in nursing homes.** *J Am Geriatr Soc* 2007, **55**:1404-1409.
18. Hewson KM, Burrell AR: **A pilot study to test the use of a checklist in a tertiary intensive care unit as a method of ensuring quality processes of care.** *Anaesth Intensive Care* 2006, **34**:322-328.
19. Huang HC, Lin WC, Lin JD: **Development of a fall-risk checklist using the Delphi technique.** *J Clin Nurs* 2008, **17**:2275-2283.
20. Makary MA, Mukherjee A, Sexton JB, Syn D, Goodrich E, Hartmann E, Rowan L, Behrens DC, Marohn M, Pronovost PJ: **Operating room briefings and wrong site surgery.** *J Am Coll Surg* 2007, **204**:236-243.
21. Verdaasdonk EG, Stassen LP, Hoffmann WF, van der Elst M, Dankelman J: **Can a structured checklist prevent problems with laparoscopic equipment?** *Surg Endosc* 2008, **22**:2238-2243.
22. Pronovost P, Berenholtz S, Dorman T, Lipsett PA, Simmonds T, Haraden C: **Improving communication in the ICU using daily goals.** *J Crit Care* 2003, **18**:71-75.
23. Lingard L, Regehr G, Orser B, Reznick R, Baker GR, Doran D, Espin S, Bohnen J, Whyte S: **Evaluation of a preoperative checklist and team briefing among surgeons, nurses, and anesthesiologists to reduce failures in communication.** *Arch Surg* 2008, **143**:12-17; discussion 18.
24. Torr J, Iacono T, Graham MJ, Galea J: **Checklists for general practitioner diagnosis of depression in adults with intellectual disability.** *J Intellect Disabil Res* 2008, **52**:930-941.
25. Iacono T, West D, Bloomberg K, Johnson H: **Reliability and validity of the revised Triple C: Checklist of Communicative Competencies for adults with severe and multiple disabilities.** *J Intellect Disabil Res* 2009, **53**:44-53.
26. Clay AS, Que L, Petrusa ER, Sebastian M, Govert J: **Debriefing in the intensive care unit: a feedback tool to facilitate bedside teaching.** *Crit Care Med* 2007, **35**:738-754.
27. Levy MM, Pronovost PJ, Dellinger RP, Townsend S, Resar RK, Clemmer TP, Ramsay G: **Sepsis change bundles: Converting guidelines into meaningful change in behavior and clinical outcome.** *Crit Care Med* 2004, **32**:S595-S597.
28. Miller GA: **The magical number seven plus or minus two: some limits on our capacity for processing information.** *Psychol Rev* 1956, **63**:81-97.
29. Lorist MM, Boksem MA, Ridderinkhof KR: **Impaired cognitive control and reduced cingulate activity during mental fatigue.** *Brain Res Cogn Brain Res* 2005, **24**:199-205.
30. Halford GS, Baker R, McCreeden JE, Bain JD: **How many variables can humans process?** *Psychol Sci* 2005, **16**:70-76.
31. Hutchins E: *Cognition in the Wild*. Cambridge, MA: MIT Press; 1996.
32. Norman DA: **Design principles for cognitive artifacts.** *Res Eng Des* 1992, **4**:43-50.
33. Xiao Y: **Artifacts and collaborative work in healthcare: methodological, theoretical, and technological implications of the tangible.** *J Biomed Inform* 2005, **38**:26-33.
34. Rodriguez-Paz JM, Mark LJ, Herzer KR, Michelson JD, Grogan KL, Herman J, Hunt D, Wardlow L, Armour EP, Pronovost PJ: **A novel process for introducing a new intraoperative program: a multidisciplinary paradigm for mitigating hazards and improving patient safety.** *Anesth Analg* 2009, **108**:202-210.
35. Surowiecki J: *The Wisdom of Crowds: Why the Many Are Smarter than the Few and How Collective Wisdom Shapes Business, Economics, Societies and Nations*. 1st edition. New York, NY: Random House; 2004.
36. Gurses AP, Seidl KL, Vaidya V, Bochicchio G, Harris AD, Hebden J, Xiao Y: **Systems ambiguity and guideline compliance: a qualitative study of how intensive care units follow evidence-based guidelines to reduce healthcare-associated infections.** *Qual Saf Health Care* 2008, **17**:351-359.
37. Oberauer K, Kliegl R: **Simultaneous cognitive operations in working memory after dual-task practice.** *J Exp Psychol Hum Percept Perform* 2004, **30**:689-707.
38. Nielsen J: *Usability Engineering*. San Diego, CA: Academic Press; 1993.
39. Strauss A, Corbin J: *Basics of Qualitative Research: Techniques and Procedures for Developing Grounded Theory*. 2nd edition. Newbury Park, CA: Sage Publications; 1998.
40. Wicker P, Smith B: **Checking the anaesthetic machine.** *J Perioper Pract* 2006, **16**:585-590.
41. Brockwell R: **The anesthesia machine: what's new besides the name?** *ASA Newsletter* 2006, 70. Available at: http://www.asahq.org/Newsletters/2006/05-06/whatsNew05_06.html.
42. O'Grady NP, Alexander M, Dellinger EP, Gerberding JL, Heard SO, Maki DG, Masur H, McCormick RD, Mermel LA, Pearson ML, Raad II, Randolph A, Weinstein RA: **Guidelines for the prevention of intravascular catheter-related infections.** *Centers for Disease Control and Prevention. MMWR Recomm Rep* 2002, **51**:1-29.
43. Pronovost PJ, Berenholtz SM, Goeschel C, Thom I, Watson SR, Holzmüller CG, Lyons JS, Lubomski LH, Thompson DA, Needham D, Hyzy R, Welsh R, Roth G, Bander J, Morlock L, Sexton JB: **Improving patient safety in intensive care units in Michigan.** *J Crit Care* 2008, **23**:207-221.
44. El-Khatib MF, Bou-Khalil P: **Clinical review: liberation from mechanical ventilation.** *Crit Care* 2008, **12**:221.
45. Pronovost PJ, Berenholtz SM, Goeschel CA, Needham DM, Sexton JB, Thompson DA, Lubomski LH, Marsteller JA, Makary MA: **Creating high reliability in healthcare organizations.** *Health Serv Res* 2006, **41**:1599-1617.
46. Koppel R, Metlay JP, Cohen A, Abaluck B, Localio AR, Kimmel SE, Strom BL: **The impact of computerized physician medication order entry in hospitalized patients—a systematic review.** *JAMA* 2005, **293**:1197-1203.
47. Eslami S, de Keizer NF, Abu-Hanna A: **The impact of computerized physician medication order entry in hospitalized patients—a systematic review.** *Int J Med Inform* 2008, **77**:365-376.
48. Windle J, Van-Milligan G, Duffy S, McClay J, Campbell J: **Rapid deployment of physician order entry using web-based, disease-specific order sets.** In *AMIA Annual Symposium Proceedings*. Omaha, NE: University of Nebraska Medical Center; 2003:1079.
49. Wrona S, Chisolm DJ, Powers M, Miler V: **Improving processes of care in patient-controlled analgesia: the impact of computerized order sets and acute pain service patient management.** *Paediatr Anaesth* 2007, **17**:1083-1089.
50. Donaldson NE, Rutledge DN, Ashley J: **Outcomes of adoption: measuring evidence uptake by individuals and organizations.** *Worldviews Evid Based Nurs* 2004, **1** Suppl 1:S41-51.
51. Titler MG: **Methods in translation science.** *Worldviews Evid Based Nurs* 2004, **1**:38-48.
52. **Atlantic Sun Airways CAT B pilot procedures and checklists series** [www.atlanticsunairways.com/training/checklist_b727f.pdf].