



Factors associated with clinical trials that fail and opportunities for improving the likelihood of success: A review

David B. Fogel

Trials.ai, 4520 Executive Dr., Suite 200, San Diego, CA, 92121, United States



ARTICLE INFO

Keywords:

Clinical trials
Enrollment
Patient burden
Pharmaceutical trials
Retention
Recruitment

ABSTRACT

Clinical trials are time consuming, expensive, and often burdensome on patients. Clinical trials can fail for many reasons. This survey reviews many of these reasons and offers insights on opportunities for improving the likelihood of creating and executing successful clinical trials. Literature from the past 30 years was reviewed for relevant data. Common patterns in reported successful trials are identified, including factors regarding the study site, study coordinator/investigator, and the effects on participating patients. Specific instances where artificial intelligence can help improve clinical trials are identified.

1. Background

Clinical trials for pharmaceuticals and medical devices offer many opportunities for failure. Failures can arise from a lack of efficacy, issues with safety, or a lack of funding to complete a trial, as well as other factors such as failing to maintain good manufacturing protocols, failing to follow FDA guidance, or problems with patient recruitment, enrollment, and retention. Generating accurate and sufficient results to determine whether or not there is merit in continuing is important at each stage in the clinical trial process. The investments of resources, time, and funding grow with successive stages, from pre-clinical through phase 3. Thus, the cost of a failed phase 3 trial is not just the cost associated with the trial itself but the cost of all prior trials as well as the cost of lost time pursuing a potentially viable alternative.

It is important to maintain a philosophy of continual improvement with respect to clinical trials broadly and specifically with an aim towards optimizing every aspect of the research and development process. A comprehensive survey of all possible points of failure in clinical trials is beyond the scope of this publication. Still, there are many factors associated with failed trials that can be distilled with evidence, along with recommendations for improving the chances of success.

2. Failing to demonstrate efficacy or safety

The primary source of trial failure has been and remains an inability to demonstrate efficacy. Hwang et al. [58] assessed 640 phase 3 trials with novel therapeutics and found that 54% failed in clinical development, with 57% of those failing due to inadequate efficacy. There are many reasons that potentially efficacious drugs can still fail to

demonstrate efficacy, including a flawed study design, an inappropriate statistical endpoint, or simply having an underpowered clinical trial (i.e., sample size too small to reject the null hypothesis), which may result from patient dropouts and insufficient enrollment.

Clinical trials also fail with respect to safety. Hwang et al. [58] found that 17% of the failed phase 3 trials examined were due to safety. Safety is addressed in every clinical trial in every phase, but issues with safety may only become apparent with the larger populations associated with phase 3 studies, or at post-approval (phase 4) or post-market [24]. Identifying safety issues is not always straightforward. Patients have individual concerns about various adverse events that may not match what physicians are concerned about. This can influence which adverse events are reported, particularly if they are mild to moderate in severity.

For example, Henon et al. [49] studied 27 phase 1 trials in diverse settings between 2014 and 2015. Prior to the start of these particular trials, patients most feared adverse events of hematuria, vomiting, and hyperglycemia, and after the trials they feared some of the same events, but also personality change, fever, and dizziness. The physicians in these trials were concerned instead with eye disorders, confusion, and blurred vision. People may have a greater propensity to present for care when they experience an adverse event that is of concern to them, and not necessarily when experiencing an adverse event of less concern to them but greater concern to the physician. Reminding patients of the importance of reporting adverse events, particularly events of special interest, is recommended for improving the likelihood of detecting safety issues earlier rather than later (e.g. [22]).

It is important also to recognize the desire for a sponsor to move a drug or device forward in the clinical trial process. Rushing studies into

E-mail address: david@trials.ai.

<https://doi.org/10.1016/j.conctc.2018.08.001>

Received 29 May 2018; Received in revised form 19 July 2018; Accepted 6 August 2018

Available online 07 August 2018

2451-8654/© 2018 The Author. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

phase 3 after successful phase 2 trials may not provide time for reflection on how best to address safety in phase 3 [107]. Research also has identified that having higher-educated nurses is associated with lower risks of mortality and failure to rescue ([3,121,128]), which may be helpful as a factor to include in study site selection.

It is critical at each stage of clinical development to have safety be a primary concern even if it is not a primary objective. The cost of uncovering a safety issue increases at each stage, including post-approval [118].

3. Financial impact

Hwang et al. [58] noted that 22% of the failed phase 3 studies they examined failed due to lack of funding. The costs required to complete the entire development process from discovery to bringing a drug to market vary, and so do estimates of these costs; however, they have been reported in excess of \$2.5 billion [34]. This includes \$1.5 billion of hard dollar out-of-pocket costs with the remainder being lost opportunity of investment costs, but does not include additional post-approval clinical trials. Focusing on phase 3 trials, the Pharmaceutical Research and Manufacturers of America estimated the cost at \$42,000 per patient in 2013, with \$10 billion spent on 1680 phase 3 clinical trials comprising over 600,000 patients.

Certain studies present unique cost considerations. For example, in a study of hospital-acquired bacterial pneumonia, the cost of a 200-site, 1000-patient phase 3 study was \$89,600 per patient [112], with screen failures being a principal driver of the cost. Pharmaceutical research and development is a costly endeavor. More generally, particularly in the United States, the cost of complying with an increasing regulatory burden is also impactful, necessitating more staff, storage, and financial outlay [43].

With such a large financial burden, many trials (in phase 3, but also earlier) are underfunded, and may not have any reasonable opportunity to generate a positive outcome (even if protocols are amended, at additional cost). This leads to ethical issues regarding patient involvement [127]. Patients generally have an expectation that their participation in a trial will lead to an advancement of knowledge based on the trial's successful completion [71]. Underfunded trials are by definition more likely to miss the enrollment needed to demonstrate statistical significance at a predefined level of efficacy.

4. Eligibility criteria

Ideally, inclusion/exclusion criteria should result in a population that matches statistically the intended general patient population [48,124]; however, study designers must account for additional concerns, including whether or not particular segments of a target population may have too many comorbidities, leading to additional higher risk of withdrawal and adverse events. For example, Hill et al. [50] noted the heterogeneous nature of pulmonary arterial hypertension (PAH), for which clinical studies have had varied eligibility, but have tended to exclude patients with advanced conditions (New York Heart Association functional class III or IV) and older women, among other categories. The correspondence between the study population and the actual population of concern can become unclear [50].

Inclusion/exclusion criteria also must be chosen in light of the expected effect on recruitment. In the case of patients with PAH, Hill et al. [50] noted the availability of competing therapies, which can suppress enrollment in any one particular study, along with investigators being influenced to recruit patients who will be stable for the duration of the study (3–4 months). Investigators may look for patients who have been stable recently, thus restricting the available population in a way that does not match the general patient population.

Inclusion criteria may vary widely across studies in a specific area, providing little guidance to a prospective sponsor or investigator. For example, in heart failure, Luo et al. [78] reported that there are no

uniform diagnostic criteria for heart failure with preserved ejection fraction (HFpEF), with approximately 55% of 121 trials using 50% as the cut-off value for diagnosing HFpEF, leaving 45% choosing another threshold.

Overly specific inclusion criteria can lead to problems in finding suitable participants. This is true particularly for conditions associated with small populations but also it applies generally. Many oncology studies, for example, have exclusions based on prior chemotherapy, having an advanced stage of disease, or not being newly diagnosed. Particularly in oncology, targeted treatments based on specific genetic markers [53] will exacerbate this issue as diagnostics screen out more individuals (hopefully with the benefit of improved efficacy). Making inclusion criteria too narrow may lead to longer recruitment times and also eventually to amending the study protocol in an attempt to recruit additional participants. Getz et al. [44] reported that 16% of protocol amendments are due to changes in inclusion/exclusion criteria, which can lead to differences in the patient populations before and after the amendment [76].

It is clear that the choice of inclusion and exclusion criteria can affect the duration and cost of a clinical trial [4], as well as the likelihood of the trial meeting desired enrollment levels and retaining sufficient participants to have an opportunity to meet a statistical endpoint. Getz et al. [44] noted that across 3400 clinical trials, more than 40% had amended protocols prior to the first subject visit, delaying trials by 4 months. Some protocol amendments cannot be avoided; however, the potential for amendments can be reduced with better planning and anticipation of the consequences from design choices.

Exclusion criteria are often presented without an explicit rationale [104]. Sometimes criteria can be put in place based on an expectation of excluding participants who may not show sufficient improvement against an endpoint, not because their health is too poor but because it is too good. For example, Hill et al. [50] reported on the endpoint of a 6-min walk test for patients with PAH. Patients who could walk more than 400 m prior to being included in trial might not be able to show much improvement (482 m in 6 min is already a 3 mph pace, which would be a moderate pace for a healthy individual). Thus, there would be pressure to exclude patients at this functional level in favor of those who could only walk between 100 and 150 m prior to inclusion. Without background knowledge, someone reviewing exclusion criteria for such a trial might not have explicit motivation to intuit the rationale for this sort of exclusion criterion.

Performing a requisite literature review for related studies remains a labor-intensive task requiring personnel with specific knowledge who can interpret the framework, criteria, and results of prior clinical trials. Future protocol development will benefit from the use of artificial intelligence tools, such as natural language processing [2,17,32,53], which will be able to extract meaningful information across published documents and present systematically organized data to the study designer for consideration. Still, the study designer must think through the implications of different inclusion/exclusion criteria (as well as objectives and endpoints) and the effects they will have on recruitment, enrollment, retention, and ultimately time and cost to completion.

5. Patient recruitment

Patients are often willing to consent to participation in a clinical trial if they believe that they have an opportunity to receive better treatment or if the results can help others [29,45,89]. Still, failing to enroll a sufficient number of subjects in a trial is a long-standing problem [82,101]. A study of 114 trials in the UK [10] indicated that only 31% met enrollment goals. In addition, Campbell et al. [15] reported that one-third of publicly funded trials required a time extension because they failed to meet initial recruitment goals.

Feller [39] reported that 25% of cancer trials failed to enroll a sufficient number of patients, and 18% of trials closed with less than

half of the target (the number of enrollees divided by the number of subjects screened) number of participants after 3 or more years. Only a relatively small percentage (2%–5%) of adult cancer patients enroll in clinical trials [41]. Furthermore, enrollment fractions (the number of enrollees divided by the number of subjects screened) can be very low [50,116].

Stensland et al. [111] reported that, particularly for cancer trials, between 2005 and 2011, the cumulative incidence of trials failing to complete was about 20%, with almost 48,000 enrolled patients in these failed trials. An earlier report from the Institute of Medicine [60] indicated that 40% of National Cancer Institute-sponsored trials were not completed.

Some studies offer remuneration to patients, generally to cover the patients' time and expenses but also in the hope that recruitment will be improved. While logic suggests that trials that offer remuneration to patients should fare better with respect to recruitment than those that do not and, moreover, patients sometimes report this as being important to them [109], evidence supporting this has been generally inconclusive. Bryant and Powell [13] found no controlled studies aimed at testing the hypothesis that paying patients to participate in a trial generates superior recruitment or retention.

Several trials have reported no observed relationship between financial incentives and recruitment [31,52,98]. On the other hand, Edwards et al. [36] reported that monetary incentives increased participant response to postal and electronic questionnaires that were designed to improve retention. Martinson et al. [81] reported effective remuneration in a smoking trial, but the trial focused on adolescents, who may have a different reaction to receiving funds than adults. Surveys show, however, that a high remuneration is often associated in patients' minds as being associated with a perception of higher risk in the trial [25,109] and thus a reluctance to enroll. The effect and effectiveness of remuneration may depend on many factors and should remain an open area of research.

6. Additional costs associated with recruitment

Beyond remuneration, the additional costs associated with patient recruitment can be difficult to estimate and highly variable, even within the same investigative area [21]. For example, Okuyemi et al. [95] conducted a study to assess the efficacy of nicotine gum and counseling to help low-income African-American smokers to quit. They reported spending \$156 per enrollee obtained via direct marketing, but \$5040 per enrollee obtained via gas-pump advertising. This suggests that marketing strategies can play an important role in the financial viability of some trials, and by consequence with the ultimate outcome as well.¹ The degree to which professional marketing expertise has been applied to help promote clinical trials is difficult to ascertain presently, but deserves specific research attention.

Healthcare providers can have a significant impact on patient recruitment and retention. Recruitment and retention can suffer when patients perceive support staff to be unavailable or uninterested, or if they have to interface routinely with new staff [33]. Encouraging patient trust in the clinical trial process may be expected to lead to better participation [125].

Incentivizing staff (providing funds for enrolling patients) has been shown to improve patient recruitment [33]. Using nurses instead of

¹ Logic also suggests that paying patients to adhere to a study protocol would lead to better adherence; however, consider the case of study HPTN 065, conducted by the HIV Prevention Trials Network in New York City and Washington D.C. in 2015 [84]. Patients were paid \$280 to take HIV medications, and an additional \$125 to be tested and have consultations aimed at reducing the spread of infection [37]. The trial is viewed generally as unsuccessful in that those receiving remuneration had only 5% higher adherence rates.

surgeons (physicians) to perform recruitment has not evidenced any difference in outcomes; however, cost savings have been realized [35,40] which may be important in supporting recruitment and retention, or other aspects of the clinical trial, indirectly.

7. Respecting the patient's concerns

Patient recruitment and retention is affected negatively when patients are concerned about being assigned to a control group rather than receiving active study drug. Part of this effect may be due to patients having poor knowledge about placebos [55] or what specific treatment is given in the control group. For patients with poor prognoses, the concern may center around not having effective treatment at all.

Ulrich et al. [122] addressed the burdens of patients participating in cancer clinical trials. Patients reported burdens including potential side effects from treatment, additional tests that would have to be endured, financial concerns (including loss of job support and work disruption), and a general worry about the unknown future, including whether or not the study drug assigned would be beneficial. Sometimes patients are not presented with a clear rationale for why their participation is important and receive minimal feedback. These concerns were echoed in Rosbach and Andersen [102], which reported on burdens on patients with multimorbidity.

In addition, scientific literacy in the general population is limited, leading to difficulty understanding information associated with a clinical trial [9,67]. Hadden et al. [46] reviewed more than 200 approved ICFs and found the mean readability was at the 10th grade level, similar to results in Paasche-Orlow et al. [97]. In a survey by Lopienski [75] of various completed trials, patients who dropped out of a trial early were twice as likely to say that it was difficult to understand the informed consent form (ICF) than were patients who completed the trial (35% vs. 16%, respectively). Moreover, even for those who completed a trial, 1 in 6 patients found the ICF vexing.

It is of interest to determine if artificial intelligence tools employing sentiment analysis could be useful in crafting the language of the ICF and other materials to provide a more compassionate tone and greater patient confidence [38,91,114] in addition to maintaining an appropriate reading level. Davis et al. [30] reported that patients prefer simplified ICFs and testing showed no lower level of patient comprehension of the details of the clinical trial when using a simplified ICF. There is an impetus for simplifying the ICF, as well as other associated written materials.

In addition, Sood et al. [110] and Cartmell et al. [18] reported that providing information regarding the clinical trial process was helpful in improving patient satisfaction. Even receiving a simple biosketch card of a healthcare provider has been associated with improved patient satisfaction [90]. Surveys of patient satisfaction conducted by a personalized health network suggest patients often have a poor experience [132]. Communicating with the patient is important at all stages of the clinical trial and supports recruitment, enrollment, and retention [61].

8. Poor recruitment, dropouts, and underpowered trials

A repeated problematic pattern in the literature is that study centers report fewer eligible patients than anticipated [6,33]. Study centers with a track record of successful performance are historically more likely to meet enrollment targets [43]. There is considerable literature reporting results from studies in which numerous study sites failed to meet enrollment, or failed to enroll any subject at all [64,72,73,105]. Levett et al. [70] identified several factors most associated with above-average recruitment rate: implementation of a clearly defined "system" of recruitment, engagement of other staff, time from ethics approval to first recruit, and the provision of a dedicated trial coordinator. A site that has historically little focus on clinical trials or presents other non-scientific impediments may lead to low investigator enthusiasm [61].

Enthusiasm from the lead investigator at a study site was the most

important factor associated with positive recruitment across 60 study centers in trial assessing the management of local post-surgical pain [26,41,51]. Thoma et al. [117] noted the critical importance of a friendly and approachable study coordinator, as this can not only improve recruitment but also lower the study's dropout rate. Slow recruitment may come from an inadequate staff and a lack of prioritizing the clinical trial over day-to-day operations [117]. It can also come when the investigator has competing trials. In addition, retention can be improved by having dedicated recruiting/support people at each study center, as well as tailoring the protocol of the trial to conform to common practices at the study center [33].

When a trial suffers too many dropouts (either based on projected or actual enrollment), the trial may become underpowered. Underpowered clinical trials are problematic. The sponsor may adapt to low enrollment by expanding the number of sites (perhaps in additional countries, with corresponding costly protocol amendments and delays in further research), increasing funds allocated to the study in an effort to meet minimum enrollment. By consequence this sometimes necessitates eliminating certain planned tests in order to reallocate available funds. In turn, certain endpoints may have an insufficient sample size to detect an important result.

As an example [61], the STICH trial (surgical treatment for ischemic heart failure) that studied the effectiveness of coronary bypass surgery in patients with heart failure took place ultimately in 26 countries, in 127 study sites, and involved 2135 patients. Originally, the study was designed to cover 32 study sites in the USA and Canada, but low patient enrollment required expanding the study to 171 sites internationally. After the expansion, 44 sites that had been approved for the trial failed to enroll a patient. The cost per deactivated site was estimated at \$10,000. The extra costs involved in expanding the trial to numerous sites in various countries meant that funds intended originally for imaging studies had to be diverted. The imaging studies were removed from the protocol, creating additional expenses for protocol amendments.

Importantly, and as mentioned briefly earlier, underpowered trials have also been described as unethical [47], even though some patients may benefit from the trial, because patients who volunteer to be in the trial are unlikely to know or appreciate that their results will not be likely to contribute to a statistically significant outcome. Carlisle et al. [16] studied 2579 trials from the National Library of Medicine clinical trial registry and found that 48,027 patients had enrolled in trials that closed in 2011 that were “unable to answer the primary research question meaningfully,” notably for failed accrual or less-than-expected enrollment. Thus, poor recruitment, enrollment, and retention remains a primary area of concern for multiple reasons.

9. Employing quantitative measures

Formulating a list of factors to consider when designing and executing a clinical trial can provide a foundation for better outcomes. However, not all factors are equally important. A well-structured mathematical framework (e.g., a Valuated State Space [87]) for trading off degrees of achievement in various parameters can offer a quantitative measure for comparing alternative choices.

For example, increasing the speed of enrollment leads to faster completion, and may be associated with fewer dropouts, better statistical power, and increased confidence in results. In contrast, consistently slow recruitment may suggest problems in inclusion/exclusion criteria [117], which should be addressed by amending a protocol. Enrollment can be accelerated by spending money on recruitment, both in advertising and in having available friendly personnel. Thus, there is a direct trade-off between the speed of enrollment and the cost of executing the trial.

Budgets are not unlimited, and therefore various trade-offs need to be considered, including not only the speed of enrollment, but the likelihood of meeting the enrollment goal. A more-effective study

center with a long history of running clinical trials successfully and with a nearby population of prospective participants may be more expensive than another more remote site with less experience. But choosing the cheaper alternative may result in failing to meet recruitment. By consequence, this may necessitate spending more on additional study centers, which come with additional costs of evaluating, training, protocol amendments, and trial execution. Quantifying these trade-offs can assist with making better decisions.

10. Considering the patient's financial burden

Given the tremendous problem of clinical trials that fail to complete due to poor recruitment, enrollment, and retention, it's of primary importance in designing and executing clinical trials to consider the burden that each patient undergoes, with the belief that retention is correlated negatively with patient burden. All burdens to the patient should be given attention, but financial impacts deserve special consideration.

The financial impact to patients in clinical trials can be easily overlooked while focusing on the objectives, endpoints, and other aspects of a particular trial design. Patients may have out-of-pocket costs when participating in a clinical trial. These include the cost of transportation and lost work, but also medical costs for additional testing. Insurance may not cover medical care beyond that which is deemed routine. Even when it does, deductibles are often quite high and a given patient may not be able to afford to participate [92].

In addition, many trials require participants to travel to their specific study centers, even for tests or procedures that could be provided locally [92], or conducted at home. Patients may need to relocate close to a study center for some period of time [79]. The additional cost of participating under these circumstances biases participation to those in higher socioeconomic levels [5,103,115,119], particularly in oncology studies. Stump et al. [113] reported on a cancer study in which 99% of participants were insured and still greater than 30% reported concern about paying for treatment. Studies also show that the financial impact of some trials can adversely event patient adherence as well as retention [8,23,130].

11. Patient time investment

While some trial participants do need to relocate during a study, many are not willing to do so [19] and most participate in local trials. Patient recruitment and retention depends in part on the willingness of the participant to travel to and from the local study center [96]. Transportation is a long-standing particular challenge for elderly participants [77,83,96,99]. Regardless of patient age, long travel times, particularly in urban areas can dissuade participation.

Research that would provide a mathematical function describing the likelihood of patient recruitment or retention as a function of distance to a study center and other factors such as demographics appears missing currently. However, some related information can provide guidance in the absence of such specific research.

The Washington State Office of Financial Management [129] surveyed resident health care consumers. Based on more than 5000 responses, adults generally reported being willing to travel less than 30 min and less than 22 miles for urgent care, and about 10% less for routine care. Demographic analysis showed that males were willing to travel for longer duration (32.8 min), as were those in non-urban areas (34.8 min), and those who were uninsured (34.1 min), each being statistically significant at $\alpha = 0.05$. Interestingly, those 65 years of age or older were willing to travel only 26.4 min, which was also statistically significant at $\alpha = 0.05$.

These data suggest the importance of recruiting patients from proximate vicinities local to study centers when in urban settings. Moreover, when incorporating older patients it is important to assist in minimizing their total time investment as they may begin from a

perspective of being less patient than the average participant. Proper site selection can help minimize long travel times.

In addition, selecting a study site with a nearby larger population pool has been correlated positively, as expected, with the likelihood of meeting recruitment targets [123]. van den Bor et al. [123] also reported statistically significant results pertaining to recruitment by geographic region. In their analyzed data, study centers in China and India were more likely to meet recruitment targets, with centers in certain locations in Western Europe and North America being least likely. It remains an open question as to whether study adherence is equivalent across these sites, and if other factors influencing positive recruitment might be associated with any operational issues.

Artificial intelligence applications offer promise in helping reducing patient time investment regardless of constraints on study site location [7,20]. In particular, evolutionary algorithms [11,66], which use computer simulations of nature's processes of variation and selection to solve problems, can assign the most appropriate study center for each prospective patient in a trial based on patient and study center availability. There is also the opportunity not only to schedule staff to support a clinical trial appropriately [12] but also to match staff with patients so that patients tend to see familiar faces at each visit (and could also request having alternative staff to interact with if desired).

In certain cases, it is possible to schedule study center visits to minimize other conflicts that a patient may have. For example, the burden on a single parent of elementary school children who must come to a study center at 10am is different than the burden for the same procedure scheduled at 4pm, after school has let out for the day. Artificial intelligence software can examine the profile of each study participant and impute the least burdensome times for appointments within the constraints of a study center's activity and the constraints of the protocol (e.g., pharmacokinetic time points). The same software can search for opportunities to reschedule patients adaptively when openings develop, making the most efficient use of the clinical trial's time.

Effective scheduling also should incorporate the patient's time spent waiting after checking in before being seen. Waiting time has been offered as being associated negatively with patient satisfaction and how patients feel about the quality of their health care [28,93,94].

Vitals Index reported 30% of US patients have left an appointment because of an excessive wait and 20% have changed doctors due to habitually long waits [69]. This same study identified superior patient satisfaction when the average wait time was just over 13 min. The lowest level of satisfaction was generated at an average wait time of just over 34 min. Long waiting times are a source of stress and can leave patients feeling disrespected, which intuitively would be associated with lower retention.

The patient's perception of a long waiting time can be reduced by assigning an additional person to facilitate interaction with the patient [108]. This additional person can also help to relieve the burden on other doctor-office staff who would be dividing their attention between patients receiving routine care and those participating in a trial.

12. Discussion

Each of the facets of protocol design, execution, and successive trial planning offers opportunities for trading off different concerns, as well as simply making inappropriate judgments leading to poor outcomes.

Study site selection is an important aspect of the clinical trial process. Poor choices can lead directly to study failure, or to a costly exercise of including additional study sites, amended protocols, and the potential for patient populations receiving different treatment regimens. When possible, having contingency plans to open additional sites, perform extra recruitment, and cover protocol amendments is recommended. The practicality of holding out reserve funds to covers these and other contingencies, however, is case specific.

Many factors for study site selection are study specific. Hurtago-Chong et al. [57] reported on a multi-step study site evaluation process,

starting from a request for participation and extending through telephone follow-up, site selection questionnaires focusing on geriatric fracture management. It is straightforward to presume that many specific requirements for a study in this specific area would not carry over to a criteria for, say, a study on pediatric oncology. Still, there are many study-site-related factors that are common to successful trials.

Getz [42] cited research from pharmaceutical companies Lilly and Pfizer, suggesting a correlation between performing well on one trial and performing well on a subsequent trial, as well as the converse of performing poorly on one trial and performing poorly on a subsequent trial. Experience with clinical trials is also important, as experience facilitates effectiveness. A site that has conducted between 6 and 10 clinical trials has a greater probability of meeting enrollment within the required time than does a site with a history of fewer trials [42]. An additional indicator is time to enroll the first patient, which is correlated with better overall performance. Data in Ref. [123] corroborate that successful experience is a predictor of positive recruitment performance. In addition, as mentioned earlier, other positive factors include an enthusiastic investigator and experienced and involved staff.

A key item deserving more attention is the minimization of patient burden and maximizing patient appreciation. This encompasses: (1) providing materials that are easy to understand, (2) having empathetic and supportive staff, (3) leadership and enthusiasm from the principal investigator, (4) a schedule (time and events) that works in synergy with the patient's constraints rather than at odds with those constraints, (5) the opportunity to adaptively reschedule visits and assign appropriate personnel to support participants, (6) trial management software to send effective reminders about visits and protocol adherence via phone, text, or email, including supporting multiple languages in multi-lingual areas [14], and (7) understanding what the patient's day-to-day experience during the trial is likely to be. Even asking a simple question such as "are you planning on moving?" can help ensure that patients are more likely to remain involved in a study, particularly if the study is of longer duration [117].

Study support staff should be generally aware of how study participants are feeling during the trial, and seek to minimize patient stress. In a survey [75], patients who described site visits as stressful were more than twice as likely to drop out (38% of dropouts vs. 16% of those who completed). Hui et al. [56] reported that in one cancer study, 21% of patients who withdrew from the trial did so because of "symptom burden." Some burdens cannot be avoided, depending on the therapy and testing involved. Patients are more likely to withdraw from a trial when they perceive their condition as not improving, even though this may be anticipated. Support staff, as well as the investigator, should seek to set patient expectations appropriately and provide appropriate empathy for any burdens that a patient is undergoing during a trial.

Patients deserve to have access to reports from studies in which they participated. Yet, Ziv [132] reported that, after completion, most studies are not available via open access. Thus, patients have to pay to be able to read a published study, even one in which they have participated. Having already given much of their time in support of a clinical trial, they may feel disrespected to have to pay to find out what information was discovered during the trial. It would be easy for a sponsor to take the position that the patient has nothing left to offer to the trial after the trial concludes and thus any additional funds required to provide article access or to provide a copy of a publication would be better allocated elsewhere. This misses the point, however, that by ensuring patient participation from start all the way through publication, the patient may feel more respected and be less likely to dropout.

Study designers should employ methods to ensure that study populations are relevant to the real-world population that is intended to benefit from treatment. Eligibility criteria should be reviewed carefully in this regard. Older patients may be viewed, correctly in some cases, as presenting more potential for comorbidities, propensity for adverse events, and for ultimately withdrawing from a trial. However, some research in breast cancer [63] reported that older patients (> 65) were

Table 1

A list of factors associated with problems or challenges when preparing for or executing a clinical trial, along with the opportunities for artificial intelligence to help alleviate these issues. Abbreviation: NLP = natural language processing.

Factor	Opportunity	Role for Artificial Intelligence
Poor study design	More complete literature review	NLP of available literature, finding similar trials, trials addressing similar issues, or trials addressing different issues utilizing similar techniques, summarized for the study designer
	Appropriate endpoints	NLP of available literature, showing endpoints/measures used in other similar studies
	Inappropriate eligibility criteria	NLP assessment of similar published trials to determine suitability of eligibility criteria and any potentially important omissions.
	Appropriate statistical analysis	NLP of available literature, summarizing statistical methods and associating these methods with successful or failed outcomes.
	Determination of appropriate sample size	Nonlinear modeling, such as with neural networks, to predict patient drop-out rates and better estimate sample size to avoid becoming underpowered. Agent-based modeling to simulate trial before execution. Use of NLP to mine previously published trials to determine sample sizes used in successful trials
Ineffective site selection	Reducing likelihood of amendments	NLP and knowledge-based processing to present designer with pertinent information to consider.
	Inconsistencies in protocol	NLP (including table-based format) to check time and events schedule against text, as well as summary of changes for any amendments.
Poor recruitment	Effective measurement of trade-offs for each site	Nonlinear modeling, such as with neural networks, to assess trade-offs site history, staff experience, investigator enthusiasm, available population, expected patient burden, and financial impact. Potential use of fuzzy logic to provide linguistic measurement descriptions.
	Improved use of funds	Optimizing communication/advertising to maximize cost effectiveness. Targeting communication to meet patient profile, including sentiment analysis.
Patient burden/safety	Ensuring appropriate eligibility criteria	NLP on prior publications to identify suitable criteria, and also criteria associated with other trial failures.
	Facilitating locating eligible patients	Database coordination, prompting investigators and patients when appropriate trials are available for specific patients.
	Enrolling patients who are likely to complete the trial	NLP and machine learning to profile patients based on prior data on who is more likely to complete a trial, reducing drop-outs.
Poor trial execution	Minimize travel and wait times	Adaptive patient scheduling, also potentially turn-by-turn driving instructions, using evolutionary algorithms. Incorporate patient profiles to tailor site assignment/schedules to patient constraints where possible. Adapt site visit schedule if possible.
	Minimize out-of-pocket expenses	Systematic review of all patient costs to identify opportunities to minimize impacts.
	Minimize possibility of contraindicated medicines/procedures	Automatic review of prior and concomitant medications for contraindications, protocol violations.
Overall	Increase likelihood of feeling respected	Sentiment analysis and other NLP tools applied to all documents provided to patients. Prompts to interacting staff for personalizing interactions. Tailored messaging to participants to increase likelihood of retention.
	Automating reporting of events	Automated prompting of events for patients and staff, reporting requirements, notes missed events, prompts for required reporting, including protocol deviations and adverse events.
Overall	Preparing data and reporting for write-up	Automatic brand/generic conversion, skeletal form generation for narratives, table creation based on specified cut-offs.
	Lack of general awareness	Situation awareness provided to investigator/study coordinator monitoring study progress, patient progress, indicating interventions if needed.
Overall	Factor analysis to improve trade-offs based on budget and other constraints	Multicriteria decision making based on Pareto analysis or single aggregated evaluation function (Valuated State Space) to quantify and illuminate trade-offs.

less likely to be eligible for breast cancer trials but when eligible their participation rate was not statistically different from the participation rate of younger eligible subjects.

Collecting and reporting data on participation and withdrawal should be more common place in order to assist with a better understanding of how to design trials so that they can complete with representative subpopulations. For example, Hui et al. [56] reported on data from palliative oncology trials. Data revealed that patients of Hispanic ethnicity (odds ratio [OR] = 1.87), those holding an advanced degree (OR ≈ 1.5), non-Christians (OR not provided), and those with higher levels of dyspnea (OR = 1.06) and fatigue (OR = 1.08) were associated with statistically significantly higher dropout rates prior to end of study. Determining the repeatability of these factors across different types of trials remains for future work.

Future efforts should also be directed toward improving the efficiency and effectiveness of clinical trials broadly (e.g., using adaptive designs) and also based on specific genetic markers or other personalized factors. Bringing eligibility to the level of the individual holds the promise for establishing greater study drug efficacy but also has the drawback of limiting the available sample size [129] to more rapidly direct the use of study drug to targets of opportunity.

Success depends crucially on identifying genetic features reliably [86,131], which could benefit from establishing collaborative databases for academic research. It will also be important to determine quality of

life measures to better assess the cost effectiveness of these tailored trials. Current data do support the cost effectiveness (in terms of additional dollars spent per month of extended life) in the case of genetic markers for acute myeloid leukemia [52] but short-term extended life (cf. multiyear extension [100]) offers only a broad indication of the potential value of a treatment.

This review covers many aspects of clinical study design that can be affected positively by appropriate design considerations and trial execution. Study site selection and addressing patient concerns are two primary areas where the effectiveness of clinical trials can be affected positively. It is important to note that there are other factors that deserve attention, which arise even in “successful” trials, including (1) whether or not the appropriate outcome measure is chosen, particularly if it is a surrogate measure [27,48,68,120], (2) how missing data are handled, and whether values are imputed [59,65], (3) the use of subjective measures that are subject to observer bias [1,54], (4) defining what observation would constitute a clinically meaningful result rather than merely a statistically significant result [80,88], (5) the lack of long-term follow up [74], (6) minimizing protocol deviations [62,85], and (7) under reporting adverse events in peer-reviewed publications [106,126].

For convenience, Table 1 offers a summary of the factors associated with problems or challenges that occur when preparing for and executing clinical trials. Some of these issues may not lead directly to the

failure of a trial; however, a series of issues can lead to a critical failure. The table presents the opportunities for improving the likelihood of success and the role that artificial intelligence may play in that improvement. Many of these factors are correlated or interrelated, thus the table is not a substitute for the greater detailed explanation found in the text.

Acknowledgments

The author thanks E. Daly-DeJoy, M. Nicoletti, F. Rahman, J. Wallace, and T. Walpole for helpful comments that improved the quality of this review.

References

- [1] I. Abraha, A. Cherubini, F. Cozzolino, R. De Florio, M.L. Luchetta, J.M. Rimland, I. Folletti, M. Marchesi, A. Germani, M. Orso, P. Eusebi, A. Montedori, Deviation from intention to treat analysis in randomised trials and treatment effect estimates: meta-epidemiology study, *BMJ* 350 (2015) h2445.
- [2] H. Alam, R. Hartono, A. Kumar, F. Rahman, Y. Tarnikova, C. Wilcox, Web page summarization for handheld devices: a natural language approach, *Proc 7th Int Conf Doc Anal Recog*, IEEE Computer Society, Edinburgh, Scotland, 2013, pp. 1153–1157.
- [3] L.A. Audet, P. Bourgault, C.M. Rochefort, Associations between nurse education and experience and the risk of mortality and adverse events in acute care hospitals: systematic review of observational studies, *Int. J. Nurs. Stud.* 80 (2018) 128–146.
- [4] C.F. Babbs, Choosing inclusion criteria that minimize the time and cost of clinical trials, *World J. Meth.* 4 (2014) 109–122.
- [5] C.R. Baquet, G.L. Elison, S.I. Mishra, Analysis of Maryland cancer patient participation in national cancer institute-supported cancer treatment clinical trials, *J. Clin. Oncol.* 26 (2008) 3380–3386.
- [6] C.S. Bennette, S.D. Ramsey, C.L. McDermott, J.J. Carlson, A. Basu, D.L. Veenstra, Predicting low accrual in the National Cancer Institute's cooperative group clinical trials, *JNCI (J. Natl. Cancer Inst.)* 108 (2016), <https://doi.org/10.1093/jnci/djv324>.
- [7] B.P. Berg, B.T. Denton, S.A. Erdogan, T. Rohleder, T. Huschka, Optimal booking and scheduling in outpatient procedure centers, *Comp. Op Res.* 50 (2014) 24–37.
- [8] D.S. Bernard, S.L. Farr, Z. Fang, National estimates of out-of-pocket health care expenditure burdens among nonelderly adults with cancer: 2001 to 2008, *J. Clin. Oncol.* 29 (2011) 2821–2826.
- [9] S. Bostock, A. Steptoe, Association between low function health literacy and mortality in older adults: longitudinal cohort study, *BMJ* (2012) 344, <https://doi.org/10.1136/bmj.e1602>.
- [10] P. Bower, P. Wallace, E. Ward, J. Grafty, J. Miller, B. Delany, A.L. Kinmonth, Improving recruitment to health research in primary care, *Fam. Pract.* 26 (2009) 391–397, <https://doi.org/10.1093/fampra/cmp037>.
- [11] J. Branke, S. Nguyen, C.W. Pickardt, M. Zhang, Automated design of production scheduling heuristics. a review, *IEEE Trans. Evol. Comput.* 20 (2015) 110–124, <https://doi.org/10.1109/TEVC.2015.2429314>.
- [12] E.K. Burke, D.P. Causmaecker, G.V. Berghe, H. Van Landeghem, The state of the art of nurse rostering, *J. Sched.* 7 (2004) 441–499.
- [13] J. Bryant, J. Powell, Payment to healthcare professionals for patient recruitment to trials: a systematic review, *BMJ* (2005) 331, <https://doi.org/10.1136/bmj.331.7529.1377.0>.
- [14] M.M. Byrne, S.L. Tannenbaum, S. Gluck, J. Hurley, M. Antoni, Participation in cancer clinical trials: why are patients not participating? *Med. Decis. Making* 34 (2014) 116–126, <https://doi.org/10.1177/0272989X1347264>.
- [15] M.K. Campbell, C. Snowdon, D. Francis, D. Elbourne, A.M. McDonald, R. Knight, A. Grant, Recruitment to randomised trials: strategies for trial enrollment and participation study: the STEPS study, *Health Technol. Assess.* 11 (2007) 105 iii-ix.
- [16] B. Carlisle, J. Kimmelman, T. Ramsay, N. MacKinnon, Unsuccessful trial accrual and human subjects protections: an empirical analysis of recently closed trials, *Clin. Trials* 12 (2015) 77–83, <https://doi.org/10.1177/1740774514558307>.
- [17] D.S. Carrell, R.E. Schoen, D.A. Leffler, M. Morris, S. Rose, A. Baer, S.D. Crockett, R.A. Gourevitch, K.M. Dean, A. Mehrotra, Challenges in adapting existing clinical natural language processing systems to multiple, diverse health care settings, *JAMIA* 24 (2017) 986–991, <https://doi.org/10.1093/jamia/ocx039>.
- [18] K.B. Cartmell, H.S. Bonilha, T. Matson, D.C. Bryant, J. Zapka, T.A. Bentz, M.E. Ford, C. Hughest-Halberg, K.N. Simpson, A.J. Alberg, Patient participation in cancer clinical trials: a pilot test of lay navigation, *Contemp. Clin. Trials Comm.* 3 (2016) 86–93.
- [19] B.-H. Chang, A.M. Hendricks, M.T. Slawsky, J.S. Locastro, Patient recruitment to a randomized clinical behavioral therapy for chronic heart failure, *BMC Med. Res. Meth.* 4 (2004) 8.
- [20] C.-F. Chien, F.-P. Tseng, C.-H. Chen, An evolutionary approach to rehabilitation scheduling. A case study, *Euro. J. Op. Res.* 189 (2008) 1234–1253.
- [21] S.P. Chin Feman, L.T. Nguyen, M.T. Quilty, C.E. Kerr, B.H. Nam, L.A. Conboy, J.P. Singer, M. Park, A. Lembo, T.J. Kapchuk, R.B. Davis, Effectiveness of recruitment in clinical trials: an analysis of methods used in a trial for irritable bowel syndrome patients, *Contemp. Clin. Trials* 29 (2008) 241–251.
- [22] D. Cohen, Rosiglitazone: what went wrong? *BMJ* 341 (2010) c4848.
- [23] K.E. Covinsky, L. Goldman, E.F. Cook, R. Oye, N. Desbiens, D. Reding, W. Fulkerson, A.F. Connors, J. Lynn, R.S. Phillips, The impact of serious illness on patients' families. SUPPORT Investigators. Study to understand prognoses and preferences for outcomes and risks of treatment, *J. Am. Med. Assoc.* 272 (1994) 1839–1844.
- [24] M. Crowther, Phase 4 research: what happens when the rubber meets the road? *ASH Education Book* 2013 (2013) 15–18.
- [25] C. Cryder, A.J. London, K. Volpp, G. Lowenstein, Informative inducement: study payment as a signal of risk, *Soc. Sci. Med.* 70 (2010) 455–464, <https://doi.org/10.1016/j.socscimed.2009.10.047>.
- [26] C. Csimma, M.F. Swiontkowski, Large clinical trials in musculoskeletal trauma: are they possible? Lessons learned from the international study of the use of rhBMP-2 in open tibial fractures, *J. Bone Joint Surg. Am.* 87 (2005) 218–222.
- [27] R.B. D'Agostino, The slippery slope of surrogate outcomes, *Curr. Contr. Trials Cardiovasc. Med.* 1 (2000) 76–78.
- [28] K.H. Dansky, J. Miles, Patient satisfaction with ambulatory healthcare services: waiting time and filling time, *Hosp. Health Serv. Adm.* 42 (1997) 165–177.
- [29] C. Daugherty, M.J. Ratain, E. Grochowski, C. Stocking, E. Kodish, R. Mick, M. Siegler, Perceptions of cancer patients and their physicians involved in phase 1 trials, *J. Clin. Oncol.* 13 (1995) 1062–1072.
- [30] T.C. Davis, R.F. Holcombe, H.J. Berkel, S. Pramanik, S.G. Divers, Informed consent for clinical trials: a comparative study of standard versus simplified forms, *J. Natl. Cancer Inst.* 90 (1998) 668–674.
- [31] N.J. De Wit, A.O. Quartero, A.P. Zuihthoff, M.E. Numans, Participation and successful patient recruitment in primary care, *J. Fam. Pract.* 50 (2001) 97681.
- [32] D. Demner-Fushman, W.W. Chapman, C.J. McDonald, What can natural language processing do for clinical decision support? *J. Biomed. Inf.* 42 (2009) 760–772, <https://doi.org/10.1016/j.jbi.2009.08.007>.
- [33] S.A. Dickson, J. Logan, S. Hagen, D. Stark, C. Glazener, A.M. McDonald, G. McPherson, Reflecting on the methodological challenges of recruiting to a United Kingdom-wide, multi-centre, randomised controlled trial in gynaecology outpatient settings, *Trials* 14 (2013) 389, <https://doi.org/10.1186/1745-6215-14-389>.
- [34] J.A. DiMasi, H.G. Grabowski, R.W. Hansen, The cost of drug development, *N. Engl. J. Med.* 372 (2015) 1972, <https://doi.org/10.1056/NEJMc1504317>.
- [35] J.L. Donovan, T.J. Peters, S. Noble, P. Powell, D. Gillatt, S.E. Oliver, J.A. Lane, D.E. Neal, F.C. Hamdy, Who can best recruit to randomized trials? *J. Clin. Epidemiol.* 56 (2003) 605–609.
- [36] P.J. Edwards, I. Roberts, M.J. Clarke, C. Diguseppi, R. Wentz, I. Kwan, R. Cooper, L.M. Felix, S. Pratap, Methods to increase response to postal and electronic questionnaires, *Cochrane Database Syst. Rev.* 8 (2009) MR0000008.
- [37] W.M. El-Sadr, D. Donnell, G. Beauchamp, I. Hall, L.V. Torian, B. Zingman, G. Lum, M. Kharfen, R. Elion, J. Leider, F.M. Gordin, V. Elharrar, D. Burns, A. Zerbe, T. Gamble, B. Branson, Financial incentives for linkage to care and viral suppression among HIV-positive patients, *JAMA Internal Medicine* 177 (2017) 1083–1092, <https://doi.org/10.1001/jamainternmed.2017.2158>.
- [38] W. Fan, M.D. Gordon, The power of social media analysis, *Commun. ACM* 57 (2014) 74–81.
- [39] S. Feller, One in Four Cancer Trials Fails to Enroll Enough Participants, (2015) https://www.upi.com/Health_News/2015/12/30/One-in-four-cancer-trials-fails-to-enroll-enough-participants/2611451485504/.
- [40] B. Fletcher, A. Georghie, D. Moore, S. Wilson, S. Damery, Improving the recruitment activity of clinicians in randomised controlled trials: a systematic review, *BMJ Open* 2 (2012) e000496.
- [41] M.N. Fouad, J.Y. Lee, P.J. Catalano, T.M. Vogt, S.Y. Zafar, D.W. West, C. Simon, C.E. Klabunde, K.L. Kahn, J.C. Weeks, C.I. Kiefe, Enrollment of patients with lung and colorectal cancers onto clinical trials, *J. Oncol. Pract.* 9 (2) (2013) e40–47, <https://doi.org/10.1200/JOP.2012.000598>.
- [42] K. Getz, Predicting successful site performance, *Appl. Clin. Trials* (2011) 1–2 November 1 <http://www.appliedclinicaltrials.com/predicting-successful-site-performance>.
- [43] K.A. Getz, Characterizing the real cost of site regulatory compliance, *Appl. Clin. Trials* 24 (2015) June 1.
- [44] K.A. Getz, R. Zuckerman, A.B. Cropp, A.L. Hindle, R. Krauss, K.I. Kaitlin, Measuring the incidence, causes, and repercussions of protocol amendments, *Drug Inf. J.* 45 (2011) 265–275.
- [45] T. Godskesen, M.G. Hansson, P. Nygren, K. Nordin, U. Kihlbom, Hope for a cure and altruism are the main motives behind participation in phase 3 clinical trials, *Eur. J. Canc. Care* (2014), <https://doi.org/10.1111/ecc.12184>.
- [46] K. Hadden, J. Holland, L. James, Understanding the Subject – plain Language IRB Informed Consents for Research, PRIM&R Adv Ethical Res Conf (Poster), (2017) San Antonio, Abstract 26.
- [47] S.D. Halpern, J.H. Karlawish, J.A. Berlin, The continuing unethical conduct of underpowered clinical trials, *J. Am. Med. Assoc.* 288 (2002) 358–362.
- [48] C. Heneghan, B. Goldacre, K.R. Mahtani, Why clinical trial outcomes fail to translate into benefits for patients, *Trials* 18 (2017) 122.
- [49] C. Henon, D. Lissa, X. Paoletti, C. Thibault, C. Le Tourneau, E. Lanoy, A. Hollebecque, C. Massard, J.-C. Sorea, S. Postel-Vinay, Patient-reported tolerability of adverse events in phase 1 trials, *ESMO Open* 2 (2017) e000148.
- [50] N.S. Hill, I.R. Preston, K.E. Roberts, Patients with pulmonary arterial hypertension in clinical trials. Who are they? *Proc. Am. Thorac. Soc.* 5 (2008) 603–609.
- [51] M. Hjorth, E. Holmberg, S. Rodjer, A. Taube, J. Westin, Physician's attitudes toward clinical trials and their relationship to patient accrual in a Nordic multicenter study on myeloma, *Contr. Clin. Trials* 17 (1996) 37286.
- [52] L. Hörster, R.F. Schlenk, M. Stadler, M. Gabriel, F. Thol, J. Schildmann, J. Vollmann, U. Rochau, G. Sroczynski, J. Wasem, A. Ganser, M. Port,

- A. Neumann, Cost-effectiveness of methods in personalized medicine. Results of a decision-analytic model in patients with acute myeloid leukemia with normal karyotype, *Leuk. Res.* 62 (2017) 84–90.
- [53] G. Hripcsak, J.H.M. Austin, P.O. Alderson, C. Friedman, Use of natural language processing to translate clinical information from a database of 889,921 chest radiographic reports, *Radiology* 224 (2002) 157–163.
- [54] A. Hrobjartsson, A.S. Thomsen, F. Emanuelsson, B. Tendam, J. Hilden, I. Boutron, P. Ravaud, S. Brorson, Observer bias in randomised clinical trials with binary outcomes: systematic review of trials with both blinded and non-blinded outcome assessors, *BMJ* 344 (2012) e1119.
- [55] J. Hughes, M. Greville-Harris, C.A. Graham, G. Lewith, P. White, F.L. Bishop, What trial participants need to be told about placebo effects to give informed consent: a survey to establish existing knowledge among patients with back pain, *J. Med. Ethics* 0 (2017) 1–4, <https://doi.org/10.1136/medethics-2016-103964>.
- [56] D. Hui, I. Glitza, G. Chisholm, S. Yennu, E. Bruera, Attrition rates, reasons and predictive factors in supportive and palliative oncology clinical trials, *Cancer* 119 (2013) 1098–1105, <https://doi.org/10.1002/ncr.27854>.
- [57] A. Hurtado-Chong, A. Joeris, D. Hess, M. Blauth, Improving site selection in clinical studies: a standardised, objective, multistep method and first experience results, *BMJ Open* 7 (2017) 7, <https://doi.org/10.1136/bmjopen-2016-014796>.
- [58] T.J. Hwang, D. Carpenter, J.C. Lauffenburger, B. Wang, J.M. Franklin, A.S. Kesselheim, Failure of investigational drugs in late-stage clinical development and publication of trial results, *JAMA Intern. Med* 176 (2016) 1826–1833.
- [59] F. Ibrahim, B.D. Tom, D.L. Scott, A.T. Prevost, A systematic review of randomised controlled trials in rheumatoid arthritis: the reporting and handling of missing data in composite outcomes, *Trials* 17 (2016) 272.
- [60] Institute of Medicine, Committee on Cancer Clinical Trials, National Academic Press, Washington D.C, 2010.
- [61] Institute of Medicine, Public Engagement and Clinical Trials: New Models and Disruptive Technologies. Work Shop Summary, The National Academies Press, Washington, DC, 2012.
- [62] S.V. Jalgaonkar, S.S. Bhide, R.K. Tripathi, Y.C. Shetty, P.A. Marathe, J. Katkar, U.M. Thatte, An audit of protocol deviations submitted to an institutional ethics committee of a tertiary care hospital, *PLoS One* 11 (2016) e0146334.
- [63] S.H. Javid, J.M. Unger, J.R. Gralow, C.M. Moynour, A.J. Wozniak, J.W. Goodwin, P.N. Lara, P.A. Williams, E.F. Hutchins, C.C. Gotay, K.S. Albain, A prospective analysis of the influence of older age of physician and patient decision-making when considering enrollment in breast cancer clinical trials, *Oncology* 17 (2010) 1180–1190, <https://doi.org/10.1634/theoncologist.2011-0384>.
- [64] R.H. Jones, H. White, E.J. Velazquez, L.K. Shaw, R. Pietrobon, J.A. Panza, R.O. Bonow, G. Spoko, C.M. O'Connor, J.-L. Rouleau, STICH (surgical treatment for ischemic heart failure) trial enrollment, *J. Am. Coll. Cardiol.* 56 (6) (2010) 490–498.
- [65] H. Kang, The prevention and handling of the missing data, *Korean J Anesthesiol* 64 (2013) 402–406.
- [66] J.M. Keller, D. Liu, D.B. Fogel, *Fundamentals of Computational Intelligence*, IEEE/Wiley, NY, 2016.
- [67] J.L. Krieger, J.M. Neil, Y.A. Strekalova, M.A. Sarge, Linguistic strategies for improving informed consent in clinical trials among low health literacy patient, *J. Natl. Cancer Inst.* 109 (2017) djw233, <https://doi.org/10.1093/jnci/djw233>.
- [68] J.L. La Cour, J. Brok, P.C. Gotzsche, Inconsistent reporting of surrogate outcomes in randomised clinical trials: cohort study, *BMJ* 341 (2010) c3653.
- [69] G. Larson, 9th Annual Vitals Wait Time Report Released, *Vitals*, 2018 March 22.
- [70] K.M. Levett, C.L. Roberts, J.M. Simpson, J.M. Morris, Site-specific predictors of successful recruitment to a perinatal clinical trial, *Clin. Trials* 11 (2014) 584–589.
- [71] M. Lievre, J. Menard, E. Bruckert, J. Cogneau, F. Delahaye, P. Giral, E. Leitersdorf, G. Luc, L. Masana, P. Moulin, P. Passa, D. Pouchain, G. Siest, Premature discontinuation of clinical trials for reasons not related to efficacy, safety, or feasibility, *BMJ* 322 (2001) 603–606.
- [72] A.M. Lincoff, J.C. Tardif, B. Neal, S.J. Nicholls, L. Ryden, G.G. Schwartz, K. Malmberg, J.B. Buse, R.R. Henry, H. Wedel, A. Wichert, R. Cannata, D.E. Grobbee, Evaluation of the dual peroxisome proliferator-activated receptor α/γ agonist aleglitazar to reduce cardiovascular events in patients with acute coronary syndrome and type 2 diabetes mellitus: rationale and design of the AleCardio trial, *Am. Heart J.* 166 (2013) 429–434.e1.
- [73] A.M. Lincoff, J.C. Tardif, G.G. Schwartz, S.J. Nicholls, L. Ryden, B. Neal, K. Malmberg, H. Wedel, J.B. Buse, R.R. Henry, A. Wichert, R. Cannata, A. Svensson, D. Volz, D.E. Grobbee, Effect of aleglitazar on cardiovascular outcomes after acute coronary syndrome in patients with type 2 diabetes mellitus: the AleCardio randomized clinical trial, *J. Am. Med. Assoc.* 311 (2014) 1515–1525.
- [74] K. Linde, L. Kriston, G. Rücker, S. Jamil, I. Schumann, K. Meissner, K. Sigterman, A. Schneider, Efficacy and acceptability of pharmacological treatments for depressive disorders in primary care: systematic review and network meta-analysis, *Ann. Fam. Med.* 13 (2015) 69–79.
- [75] K. Lopienski, Retention in Clinical Trials – Keeping Patients on Protocols, (2015) June 1 <https://forteresearch.com/news/infographic/infographic-retention-in-clinical-trials-keeping-patients-on-protocols/>.
- [76] C. Löscher, M. Neuhäuser, The statistical analysis of a clinical trial when a protocol amendment changed the inclusion criteria, *BMC Med. Res. Meth.* 8 (2008) 16.
- [77] L.C. Lovato, K. Hill, S. Hertert, D.B. Hunninghake, J.L. Probstfield, Recruitment for controlled clinical trials: literature summary and annotated bibliography, *Contr. Clin. Trials* 18 (1997) 328–352.
- [78] H. Luo, Y. Xu, F. Yue, C. Zhang, C. Chen, Quality of inclusion criteria in the registered clinical trials of heart failure with preserved ejection fraction: is it time for a change? *Int. J. Cardiol.* 254 (2018) 210–214.
- [79] N.S. Majhail, J.D. Rizzo, T. Hahn, S.J. Lee, P.L. McCarthy, M. Ammi, E. Denzen, R. Drexler, S. Flesch, H. James, N. Omondi, T.L. Pedersen, E. Murphy, K. Pederson, Pilot study of patient and caregiver out-of-pocket costs of allogeneic hematopoietic cell transplantation, *Bone Marrow Transplant.* 28 (2013) 865–871, <https://doi.org/10.1038/bmt.2012.248>.
- [80] B. Make, How can we assess outcomes of clinical trials: the MCID approach, *COPD* 4 (2007) 191–194.
- [81] B.C. Martinson, D. Lazovich, H.A. Lando, C.L. Perry, P.G. McGovern, R.G. Boyle, Effectiveness of monetary incentives for recruiting adolescents to an intervention trial to reduce smoking, *Prev. Med.* 31 (2000) 706–713.
- [82] A.M. McDonald, R.C. Knight, M.K. Campbell, V.A. Entwistle, A.M. Grant, J.A. Cook, D.R. Elbourne, D. Francis, J. Garcia, I. Roberts, C. Snowdon, What influences recruitment to randomised controlled trials? A review of trials funded by two UK funding agencies, *Trials* 7 (2006) 9, <https://doi.org/10.1186/1745-6215-7-9>.
- [83] E.A. McNeely, S.D. Clements, Recruitment and retention of the older adult into research studies, *J. Neurosurg. Nurs.* 26 (1994) 57–61.
- [84] D.G. McNeil, Study that Paid Patients to Take HIV Drug Fails, *New York Times*, (2015) Feb. 24.
- [85] M. Mehra, K. Kurpanek, M. Petrizzo, S. Brenner, Y. McCracken, T. Katz, M. Gurian, The life cycle and management of protocol deviations, *Therapeut Innov Reg Sci* 49 (2014) 762–777.
- [86] A. Mehta, C.U. Awah, A.M. Sonabend, Topoisomerase II poisons for glioblastoma; existing challenges and opportunities to personalize therapy, *Front. Neurol.* 9 (2018) 459.
- [87] Z. Michalewicz, D.B. Fogel, *How to Solve it: Modern Heuristics*, second ed., Springer, Berlin, 2004, pp. 444–449.
- [88] F.J. Molnar, M. Man-Son-Hing, D. Fergusson, Systematic review of measures of clinical significant employed in randomized controlled trials of drugs for dementia, *J. Am. Geriatr. Soc.* 5793 (2009) 536–546.
- [89] S.Y. Moorcraft, C. Marriott, C. Peckitt, D. Cunningham, I. Chau, N. Starling, D. Watkins, S. Rao, Patients' willingness to participate in clinical trials and views on aspects of cancer research: results of a prospective patient survey, *Trials* 17 (2016) 17, <https://doi.org/10.1186/s13063-015-1105-3>.
- [90] B.J. Morris, J.E. Richards, K.R. Archer, M. Lasater, D. Rabalais, M.K. Sethi, A.A. Jahangir, Improving patient satisfaction in the orthopaedic trauma population, *J. Orthop. Trauma* 28 (2014) e80–84, <https://doi.org/10.1097/01.bot.0000435604.75873.ba>.
- [91] B.L. Neiger, R. Thackeray, S.A. Van Wagenen, Use of social media in health promotion, *Health Promot. Pract.* 13 (2012) 159–164.
- [92] R.D. Nipp, E. Powell, B. Chabner, B. Moy, Recognizing the financial burden of cancer patients in clinical trials, *Oncology* 20 (2015) 572–575.
- [93] M.H. Oermann, Effects of educational intervention in waiting room on patient satisfaction, *J. Ambul. Care Manag.* 26 (2003) 150–158.
- [94] M.H. Oermann, C.A. Needham, M.T. Doyal, L. Sinishaj, M.P. Lange, Filling the waiting time in the clinic with education about glaucoma, *Insight* 26 (2001) 77–80.
- [95] K.S. Okuyemi, L.S. Cox, N.L. Nollen, T.M. Snow, H. Kaur, W. Choi, N. Nazir, M.S. Mayo, J.S. Ahluwalia, Baseline characteristics and recruitment strategies in a randomized clinical trial of African-American light smokers, *Am. J. Health Promot.* 21 (2007) 183–191.
- [96] M.G. Ory, P.D. Lipman, P.L. Karlen, M.B. Gerety, V.J. Stevens, M.A. Singh, D.M. Buchner, K.B. Schechtman, FICSIT Group, Frailty and injuries: cooperative studies of intervention techniques, *Prev. Sci.* 3 (2002) 1–22.
- [97] M.K. Paasche-Orlow, H.A. Taylor, F.L. Brancati, Readability standards for informed-consent forms compared with actual readability, *NEJM* 348 (2003) 721–726.
- [98] A. Pearl, S. Wright, G. Gamble, R. Doughty, N. Sharpe, Randomised trials in general practice: a New Zealand experience in recruitment, *N. Z. Med. J.* 116 (2003) 6817.
- [99] H. Petrovitch, R. Byington, G. Bailey, P. Borhani, S. Carmody, L. Goodwin, J. Harrington, H.A. Johnson, P. Johnson, M. Jones, Systolic hypertension in the elderly program (SHEP). Part 2: screening and recruitment, *Hypertension* 17 (1991) II16–23.
- [100] K.A. Phillips, J.A. Sakowski, J. Trosman, M.P. Douglas, S.-Y. Liang, P. Neumann, The economic value of personalized medicine tests: what we know and what we need to know, *Genet. Med.* 16 (2014) 251–257.
- [101] R.J. Prescott, C.E. Counsell, W.J. Gillespie, A.M. Grant, I.T. Russell, S. Kiauka, I.R. Colthart, S. Ross, S.M. Shepherd, D. Russell, Factors that limit the quality, number and progress of randomised controlled trials, *Health Technol. Assess.* 3 (1999) 1143.
- [102] M. Rosbach, J.S. Andersen, Patient-experienced burden of treatment in patients with multimorbidity – a systematic review of qualitative data, *PLoS One* 12 (2017) e0179916, <https://doi.org/10.1371/journal.pone.0179916>.
- [103] W.B. Sateren, E.L. Trimble, J. Abrams, O. Brawley, N. Breen, L. Ford, M. McCade, R. Kaplan, M. Smith, R. Ungerleider, M.C. Christian, How sociodemographics, presence of oncology specialists, and hospital cancer programs affect accrual to cancer treatment trials, *J. Clin. Oncol.* 20 (2002) 2109–2117.
- [104] A.F. Schmidt, R.H.H. Groenwold, J.J.M. van Delden, Y. van der Does, O.H. Klungel, K.C.B. Roes, A.W. Hoes, R. van der Graaf, Justification of exclusion criteria was underreported in a review of cardiovascular trials, *J. Clin. Epidemiol.* 67 (2014) 635–644.
- [105] A.T. Schroen, G.R. Petroni, H.W.R. Gray, W. Cronin, D.J. Sargent, J. Benedetti, D.L. Wickerham, B. Djubegovic, C.L. Slingluff, Preliminary evaluation of factors associated with premature trial closure and feasibility of accrual benchmarks in phase III oncology trials, *Clin. Trials* 7 (2010) 312–321.
- [106] J.B. Schroll, E.I. Penninga, P.C. Götzsche, Assessment of adverse events in

- protocols, clinical study reports, and published papers of trials of Orlistat: a document analysis, *PLoS Med.* 13 (2016) e1002101.
- [107] A. Shanley, Preventing phase III failures, *Pharmaceut. Technol.* 2016 (2016) 24–27.
- [108] H.A. Sherwin, M. McKeown, M.F. Evans, O.K. Bhattacharyya, The waiting room "wait", *Can. Fam. Physician* 59 (2013) 479–481.
- [109] J. Slomka, S. McCurdy, E. Ratliff, P. Timpson, M. Williams, Perceptions of financial payment for research participation among African-American drug users in HIV studies, *J. Gen. Intern. Med.* 10 (2007) 1403–1409.
- [110] A. Sood, K. Prasad, L. Chhatwani, E. Shinozaki, S.S. Cha, L.L. Loehrer, D.L. Wahner-Roedler, Patients' attitudes and preferences about participation and recruitment strategies in clinical trials, *Mayo Clin. Proc.* 84 (2009) 243–247.
- [111] K.D. Stensland, R.B. McBride, A. Latif, J. Wisnivesky, R. Hendricks, N. Roper, P. Boffetta, S.J. Hall, W.K. Oh, M.D. Galsky, Adult cancer clinical trials that fail to complete: an epidemic? *JNCI (J. Natl. Cancer Inst.)* 106 (2014) dju299 <https://doi.org/10.1093/jnci/dju229>.
- [112] S. Stergiopoulos, S.B. Calvert, C.A. Brown, J.A. Awatin, P. Tenaerts, T.L. Holland, J.A. DiMasi, K.A. Getz, Cost drivers of a hospital-acquired bacterial pneumonia and ventilator-associated bacterial pneumonia phase 3 clinical trial, *Clin. Infect. Dis.* 66 (2018) 72–80, <https://doi.org/10.1093/cid/cix726>.
- [113] T.K. Stump, N. Eghan, B.L. Engleston, O. Hamilton, M. Pirolo, J.S. Schwartz, K. Armstrong, J.R. Beck, N.J. Meropol, Y.-N. Wong, Cost concerns of patients with cancer, *J. Oncol Pract* 9 (2013) 251–257.
- [114] M. Taboada, J. Brooke, M. Tofiloski, K. Voll, M. Stede, Lexicon-based methods for sentiment analysis, *Comput. Ling.* 37 (2011) 267–307.
- [115] L. Talarico, G. Chen, R. Pazdur, Enrollment of elderly patients in clinical trials for cancer drug registration: a 7-year experience by the US Food and Drug Administration, *J. Clin. Oncol.* 22 (2004) 4626–4631.
- [116] R.W. Taylor, J.L. Zimmerman, R.P. Dellenger, R.C. Straube, G.J. Criner, K. Davis, K.M. Kelly, T.C. Smith, R.J. Small, Inhaled nitric oxide in ARDS study group. Low-dose inhaled nitric oxide in patients with acute lung injury: a randomized controlled trial, *J. Am. Med. Assoc.* 291 (2004) 1603–1609.
- [117] A. Thoma, F. Farrokhyar, L. McKnight, M. Bhandari, How to optimize patient recruitment, *Can. J. Surg.* 53 (2010) 205–210.
- [118] C.H. Tong, L.-I. Tong, J.E. Tong, The Vioxx recall case and comments, *Compet. Rev.* 19 (2009) 114–118.
- [119] C.A. Townsley, R. Selby, L.L. Siu, Systematic review of barriers to the recruitment of older patients with cancer onto clinical trials, *J. Clin. Oncol.* 33 (2005) 3112–3124.
- [120] S. Twaddell, Surrogate outcome markers in research and clinical practice, *Aust. Prescr.* 32 (2009) 47–50.
- [121] D. Twigg, C. Duffield, A. Bremner, P. Rapley, J. Finn, Impact of skill mix variations of patient outcomes following implementation of nursing hours per patient day staffing: a retrospective study, *J. Adv. Nurs.* 68 (2012) 2710–2718.
- [122] C.M. Ulrich, K.A. Knafel, S.J. Ratcliff, T.S. Richmond, C. Grady, C. Miller-Davis, G.R. Wallen, Developing a model of the benefits and burdens of research participation in cancer clinical trials, *AJOB Prim Res* 3 (2012) 10–23.
- [123] R.M. van den Bor, D.E. Grobbee, B.J. Oosterman, P.W.J. Vaessen, K.C.B. Roes, Predicting enrollment performance of investigational centers in phase III multi-center clinical trials, *Contemp Clin Trials Comm* 7 (2017) 208–216.
- [124] J. Verster, A.J. van de Loo, T. Roehrs, T. Roth, Are clinical trial participants representative for patients with insomnia? *Sleep* 40 (2017) A148, <https://doi.org/10.1093/sleep/zsx050.397>.
- [125] P.R. Ward, P. Rokkas, C. Cenko, M. Pulvirenti, N. Dean, A.S. Carney, S. Meyer, "Waiting for" and "waiting in" public and private hospitals: a qualitative study of patient trust in South Australia, *BMC Health Serv. Res.* 17 (2017) 333.
- [126] B. Wieseler, N. Wolfram, N. McGauran, M.F. Kerekes, V. Vervolgyi, P. Kohlepp, M. Kamphuis, U. Groven, Completeness of reporting of patient-relevant clinical trial outcomes: comparison of unpublished clinical study reports with publicly available data, *PLoS Med.* 10 (2013) e1001526.
- [127] R.J. Williams, T. Tse, K. DePiazza, D. Zarin, Terminated trials in the ClinicalTrials.gov results database: evaluation of availability of primary outcome data and reasons for termination, *PLoS One* 10 (2015) e0127242, <https://doi.org/10.1371/journal.pone.0127242>.
- [128] S. Wilson, A. Bremner, Y. Hauck, J. Finn, The effect of nurse staffing on clinical outcomes of children in hospital: a systematic review, *Int. J. Evid. Base. Healthc.* 9 (2011) 97–121.
- [129] W. Yen, How Long and How Far Do Adults Travel for Primary Care? Research Brief No. 70 Washington State Health Services Research Project, 2013 (April).
- [130] S.Y. Zafar, J.M. Peppercorn, D. Schrag, D.H. Taylor, A.M. Goetzinger, X. Zhong, A.P. Abernethy, The financial toxicity of cancer treatment. A pilot study assessing out-of-pocket expenses and the insured cancer patient's experience, *Oncology* 18 (2013) 381–390.
- [131] D. Zardavas, M. Piccart-Gebhart, Clinical Trials of Precision Medicine through Molecular Profiling: Focus on Breast Cancer, *Am Soc Clin Oncol Educ Book*, 2015, pp. e183–190.
- [132] S. Ziv, How to Design a Better Clinical Trial with the Patient Experience in Mind, *Newsweek*, 2017 July 21 <http://www.newsweek.com/cancer-patients-clinical-trials-design-better-experience-639310>.