

Original Paper

Identifying National Availability of Abortion Care and Distance From Major US Cities: Systematic Online Search

Alice F Cartwright¹, MPH; Mihiri Karunaratne², BA; Jill Barr-Walker³, MPH, MS; Nicole E Johns¹, MPH; Ushma D Upadhyay¹, MPH, PhD

¹Advancing New Standards in Reproductive Health, Department of Obstetrics, Gynecology, and Reproductive Sciences, University of California, San Francisco, Oakland, CA, United States

²University of California, Berkeley, Berkeley, CA, United States

³Zuckerberg San Francisco General Library, University of California, San Francisco, San Francisco, CA, United States

Corresponding Author:

Ushma D Upadhyay, MPH, PhD

Advancing New Standards in Reproductive Health

Department of Obstetrics, Gynecology, and Reproductive Sciences

University of California, San Francisco

1330 Broadway, Suite 1100

Oakland, CA, 94612

United States

Phone: 1 510 986 8946

Email: ushma.upadhyay@ucsf.edu

Abstract

Background: Abortion is a common medical procedure, yet its availability has become more limited across the United States over the past decade. Women who do not know where to go for abortion care may use the internet to find abortion facility information, and there appears to be more online searches for abortion in states with more restrictive abortion laws. While previous studies have examined the distances women must travel to reach an abortion provider, to our knowledge no studies have used a systematic online search to document the geographic locations and services of abortion facilities.

Objective: The objective of our study was to describe abortion facilities and services available in the United States from the perspective of a potential patient searching online and to identify US cities where people must travel the farthest to obtain abortion care.

Methods: In early 2017, we conducted a systematic online search for abortion facilities in every state and the largest cities in each state. We recorded facility locations, types of abortion services available, and facility gestational limits. We then summarized the frequencies by region and state. If the online information was incomplete or unclear, we called the facility using a mystery shopper method, which simulates the perspective of patients calling for services. We also calculated distance to the closest abortion facility from all US cities with populations of 50,000 or more.

Results: We identified 780 facilities through our online search, with the fewest in the Midwest and South. Over 30% (236/780, 30.3%) of all facilities advertised the provision of medication abortion services only; this proportion was close to 40% in the Northeast (89/233, 38.2%) and West (104/262, 39.7%). The lowest gestational limit at which services were provided was 12 weeks in Wyoming; the highest was 28 weeks in New Mexico. People in 27 US cities must travel over 100 miles (160 km) to reach an abortion facility; the state with the largest number of such cities is Texas (n=10).

Conclusions: Online searches can provide detailed information about the location of abortion facilities and the types of services they provide. However, these facilities are not evenly distributed geographically, and many large US cities do not have an abortion facility. Long distances can push women to seek abortion in later gestations when care is even more limited.

(*J Med Internet Res* 2018;20(5):e186) doi:[10.2196/jmir.9717](https://doi.org/10.2196/jmir.9717)

KEYWORDS

abortion seekers; reproductive health; internet; access to information; information seeking; abortion patients; reproductive health services; information seeking behavior

Introduction

Women's ability to determine if and when they get pregnant and continue that pregnancy is key to their overall well-being. Women who are denied wanted abortions experience some negative outcomes compared with women who were able to obtain abortions, including increased economic insecurity [1] and continued exposure to violence from the man involved in the pregnancy [2]. While abortion rates have declined slightly in recent years, over 926,000 abortions were performed in the United States in 2014 [3]. This rate is equivalent to 1 in 4 women of reproductive age having an abortion within her lifetime [3], which underscores that abortion is common.

The explanations for the decline in abortion rates are varied, but part of this drop can likely be attributed to the decrease in facilities at which women can obtain abortion care across the United States over the past decade. Most abortions (95%) are performed in specialized abortion clinics (rather than private physicians' offices or hospitals), and the number of these clinics declined in half of US states from 2011 to 2014, with some regions experiencing up to a 22% decrease [3]. Because 90% of US counties do not have an abortion provider [3], many women seeking abortion must travel outside their home counties to obtain care. Other geographic disparities have been documented: women living in rural areas, the South and Midwest regions of the United States, and those seeking second-trimester or later abortions are more likely to travel farther for services, often 50 miles (80 km) or more one way [4-7]. These shifts in the availability of abortion-providing facilities indicate that women in underserved areas must travel increasingly far for abortion care.

Somedecline in the number of abortion facilities may be due to the more than 400 state laws regulating abortion that have been adopted between 2011 and 2017 [8], which, among other requirements, mandate that physicians have local hospital admitting privileges, facilities have formal transfer agreements with local hospitals, and facilities become ambulatory surgical centers. These laws have likely led to the closure of facilities that could not meet the financial or administrative requirements imposed by these laws. For example, after these types of laws were passed in Texas in 2013, the number of abortion facilities decreased by 54% over 15 months, requiring women whose nearest clinic had closed to travel 85 miles (137 km) one way to a facility [9]. Additional analyses of trends in abortion rates in Texas from 2012 to 2014 found a relationship between increases in distance to the nearest abortion facility and decreases in the county abortion rate [10]. Another analysis from Louisiana estimated that, if admitting privileges laws were to go into effect, 67% of women of reproductive age would live more than 150 miles (241 km) from the nearest abortion facility, thereby tripling the distance women have to travel to reach the nearest facility for care [11,12]. With distance come increased travel time, increased costs for transportation and childcare, lost wages, the need to take time off of work or school, the need to disclose the abortion to more people than desired, and overall delays in care [13-15]. Ultimately, delays in reaching and obtaining care can push women later into their pregnancies, even up to the point that they might not be able to obtain a

wanted abortion, depending on the gestational limits on abortion in their state [16].

To obtain abortion care in their communities, women who do not know where to go may use the internet to find abortion facility information [15]. Almost half (45%) of women seeking abortion services at clinics in Nebraska located the abortion clinic through an online search [17], and a recent study documented an interest in information on self-abortion among people searching online using the search engine Google [18]. Online searching for abortion information appears to be more prevalent in states with restrictive abortion laws and where abortion availability is limited, suggesting that women with reduced access to abortion are more likely to seek out information on abortion online [19,20].

We were interested in examining the question "What does the current landscape of abortion facilities look like to women searching online for abortion services?" There are no publicly available systematically documented and comprehensive lists of US abortion facilities, which makes it difficult to determine how far women must travel to obtain these services. Considering the trends in increased restrictions and decreasing numbers of abortion-providing facilities, it is important to generate accurate estimates of the distances women must travel to obtain abortion services in order to demonstrate potential impacts of closures. This study aimed to address this question by documenting the location of and abortion services available at abortion facilities identified through a systematic online search in all 50 US states (and the District of Columbia) and then calculating travel distances to these facilities from metropolitan areas with populations of 50,000 or more.

Methods

Data Collection

We conducted a systematic online search for abortion facilities using the Google (Google LLC), Bing (Microsoft Corporation), and Yahoo (Oath Inc) search engines between February 22, 2017 and May 22, 2017. Although Google alone accounts for a substantial portion of the market share in the United States (87.5%), together the 3 search engines comprised 99.1% of the total search engine market share as of February 2017 [21]. We conducted a search with the keywords "Abortion clinic in [state]" (no quotes) for all 50 states and the District of Columbia in each of the 3 search engines. In addition, we searched all cities (n=302) with populations over 100,000 based on 2015 US Census population estimates [22] using the keywords "Abortion clinic in [city]" (no quotes). For states that had fewer than 3 cities with populations over 100,000, we used the 3 most populous cities from the same US Census source. We conducted the keyword searches in Google's Chrome browser on Incognito mode and cleared the complete browsing history, including cookies and other site data and cached images and files, prior to each search. The researcher was logged into a Google account created specifically for this study during the searches. We chose keyword searches to reflect the natural language that women would use to search for local abortion services.

We assessed the first 20 results for each *city* for information on abortion-providing facilities, similar to previous analyses of search engine content [23,24], resulting in a review of a total of 18,120 city results across all 3 search engines. To capture the larger number of facilities expected in statewide searches, we reviewed the first 30 results for each state (for a total review of 4590 state results). For each result, we examined the website for relevant information. If the website belonged to an abortion facility, we included the result in our count of facilities and recorded whether they provided medication abortion, or aspiration or surgical abortion, as well as the facility gestational limit. Some facilities noted on their websites that they offered services beyond the gestational limit on a case-by-case basis; however, we recorded the limit that each facility offered to all patients seeking services. We included hospitals and clinics associated with universities and medical schools through the Ryan Residency Training Program in Abortion and Family Planning [25] in the analysis if they provided information about availability of abortion services on their website, even if they did not come up in our systematic Web searches. If a website did not provide information about a facility where abortion care could be obtained or explicitly stated that they did not provide abortion care, we excluded the facility. If the online information was incomplete or unclear, we called the facility using a mystery shopper method, which simulates the perspective of patients calling for services [26]. With these calls, we verified that it was not a crisis pregnancy center, confirmed that the facility was open and providing abortion services, and obtained additional information on its address, including state and zip code, types of abortion services provided, and gestational limits. Finally, because Planned Parenthood is the health care provider most widely known to provide abortion services in the United States, we reviewed all facilities listed by state on the Planned Parenthood Federation of America website as a validity check against the results from our systematic search. We confirmed that all Planned Parenthood facilities providing abortion had been captured by our searches. The study was approved by the institutional review board of the University of California, San Francisco.

Data Analysis

We described the number of facilities and the proportion of facilities that offered medication abortion only, aspiration or surgical abortion only, and both medication and aspiration or surgical abortion in each state and region. We grouped states by region and subregion based on US Census categories. The latest gestational limit at which facilities offered aspiration or surgical abortions was documented for each state. Using 2015 population estimates taken from 2010 US Census data [22], we determined the number of women of reproductive age (15-49 years) per abortion facility in each state.

To calculate the cities farthest from an abortion-providing facility, we defined cities based on the US Census's data on incorporated places of 50,000 or more [22], which amounted to 758 cities. After removing those cities that had at least one abortion provider, we calculated the distance from each city to all the abortion facilities within the state and in any neighboring or nearby states. For each city, we then took the minimum of

these distances to determine the closest provider. We calculated distances in Stata 14 (StataCorp LLC) using the `traveltime3` command, which uses a Google Maps application programming interface to calculate driving distances in miles and time. Mapping was performed in Redivis, a Stanford University-based online visualization platform. Rather than using Euclidean (straight-line) distance, Redivis uses road network information, including road type and corresponding average speed, sourced from OpenStreetMap [27] to implement a cost-distance algorithm to predict distance-access to abortion facilities.

Results

Distribution and Characteristics of Abortion Facilities in the United States

We identified 780 abortion facilities in the United States. The distribution of abortion facilities was not uniform across states. The largest numbers of facilities were in the Northeast and the West. California had the highest number of facilities ($n=152$), while Kentucky, Mississippi, Missouri, North Dakota, South Dakota, and West Virginia had 1 facility each. The geographic region with the highest ratio of women of reproductive age to facility was the Midwest, with 165,886 women per abortion facility (Table 1). The Northeast had the lowest ratio (55,662:1). While population density is not distributed evenly across all regions, the subregions with the highest ratios were the West South Central and East South Central subregions, with 298,733 and 288,463 women per facility, respectively. The state with the highest ratio of women to facility was Missouri, with 1,365,575 women per facility, and the lowest was in Maine, with 13,905 women per facility.

Most facilities reported providing both medication and aspiration or surgical abortion, although the proportion of facilities that provided different types of abortion care also varied by region and state. Over 30% (236/780, 30.3%) of all the facilities reported on their websites that they only provided medication abortion (Table 2), while 65.4% (510/780) provided both medication and aspiration or surgical abortion. Very few offered just aspiration or surgical abortion.

The South region had the highest proportion of facilities offering both medication abortion and aspiration or surgical abortion (169/193, 87.6%). While the Northeast and West had many more facilities overall, almost 40% of facilities in each of these regions offered medication abortion only (89/233, 38.2%; and 104/262, 39.7%, respectively). When looking at subregion, New England (34/74, 45.9%) and Pacific (87/207, 42.0%) had even greater proportions of facilities offering medication abortion only.

The highest gestational limit advertised by facilities also varied by state, subregion, and region (Table 2). The states with the lowest advertised gestation for abortions were Wyoming (12 weeks) and Indiana and South Dakota (both 13 weeks and 6 days), and the lowest subregions were West North Central in the Midwest and East South Central in the South, with limits of 22 weeks. Among all facilities, 50.9% (397/780) provided abortion services at 14 weeks or later and 26.5% (207/780) provided services at 20 weeks or later.

Table 1. Number of US abortion facilities by region and state, May 2017.

Region and state	Number of facilities	Population of women of reproductive age (ages 15-49 years) per facility
United States	780	95,033
Northeast	233	55,662
New England	74	45,655
Connecticut	20	40,632
Maine	20	13,905
Massachusetts	19	84,973
New Hampshire	6	48,395
Rhode Island	3	82,130
Vermont	6	22,743
Middle Atlantic	159	60,320
New Jersey	50	40,927
New York	92	51,293
Pennsylvania	17	166,210
Midwest	92	165,886
East North Central	68	155,508
Indiana	6	250,997
Illinois	25	120,135
Michigan	23	96,054
Ohio	11	235,016
Wisconsin	3	423,591
West North Central	24	195,289
Iowa	9	75,629
Kansas	4	161,029
Minnesota	5	245,486
Missouri	1	1,365,575
Nebraska	3	140,140
North Dakota	1	167,601
South Dakota	1	181,145
South	193	145,645
South Atlantic	147	98,787
Delaware	3	70,851
District of Columbia	3	66,863
Florida	65	67,757
Georgia	17	145,646
Maryland	25	56,665
North Carolina	15	155,709
South Carolina	3	370,733
Virginia	15	131,439
West Virginia	1	392,351
East South Central	15	288,463
Alabama	5	223,458

Region and state	Number of facilities	Population of women of reproductive age (ages 15-49 years) per facility
Kentucky	1	996,488
Mississippi	1	694,045
Tennessee	8	189,891
West South Central	31	298,733
Arkansas	3	222,577
Louisiana	3	363,228
Oklahoma	4	220,527
Texas	21	315,296
West	262	67,883
Mountain	55	97,547
Arizona	8	190,750
Colorado	21	60,902
Idaho	4	91,376
Montana	5	43,161
Nevada	8	83,522
New Mexico	5	91,243
Utah	2	363,970
Wyoming	2	63,250
Pacific	207	60,002
Alaska	6	27,969
California	152	61,740
Hawaii	3	103,715
Oregon	12	75,968
Washington	34	48,391

Table 2. Types of services offered by abortion care facilities (N=780) by US region and state, May 2017.

Region and state	Facilities offering only aspiration or surgical abortion, n (%)	Facilities offering only medication abortion, n (%)	Facilities offering both aspiration or surgical abortion and medication abortion, n (%)	Latest gestational limit (weeks since LMP ^a) as listed on website
United States	34 (4.4)	236 (30.3)	510 (65.4)	28
Northeast	18 (7.7)	89 (38.2)	126 (54.1)	27
New England	4 (5.4)	34 (45.9)	36 (48.6)	27
Connecticut	0 (0)	13 (65.0)	7 (35.0)	<24
Maine	0 (0)	17 (85.0)	3 (15.0)	<19
Massachusetts	4 (21.1)	0 (0)	15 (78.9)	27
New Hampshire	0 (0)	1 (16.7)	5 (83.3)	<16
Rhode Island	0 (0)	0 (0)	3 (100)	<22
Vermont	0 (0)	3 (50.0)	3 (50.0)	<19
Middle Atlantic	14 (8.8)	55 (34.6)	90 (56.6)	<25
New Jersey	6 (12.0)	22 (44.0)	22 (44.0)	<25
New York	8 (8.7)	29 (31.5)	55 (59.8)	24
Pennsylvania	0 (0)	4 (23.5)	13 (76.5)	23
Midwest	2 (2.2)	23 (25.0)	67 (72.8)	24
East North Central	1 (1.5)	16 (23.5)	51 (75.0)	24
Indiana	0 (0)	1 (16.7)	5 (83.3)	<14
Illinois	0 (0)	9 (36.0)	16 (64.0)	24
Michigan	0 (0)	5 (21.7)	18 (78.3)	24
Ohio	1 (9.1)	1 (9.1)	9 (81.8)	<22
Wisconsin	0 (0)	0 (0)	3 (100)	<23
West North Central	1 (4.2)	7 (29.2)	16 (66.7)	22
Iowa	0 (0)	6 (66.7)	3 (33.3)	<22
Kansas	0 (0)	1 (25.0)	3 (75.0)	<22
Minnesota	1 (20.0)	0 (0)	4 (80.0)	22
Missouri	0 (0)	0 (0)	1 (100)	<22
Nebraska	0 (0)	0 (0)	3 (100)	<22 ^b
North Dakota	0 (0)	0 (0)	1 (100)	16
South Dakota	0 (0)	0 (0)	1 (100)	<14
South	4 (2.1)	20 (10.4)	169 (87.6)	26
South Atlantic	2 (1.4)	15 (10.2)	130 (88.4)	26
Delaware	0 (0)	1 (33.3)	2 (66.7)	<16
District of Columbia	1 (33.3)	0 (0)	2 (66.7)	26
Florida	1 (1.5)	5 (7.7)	59 (90.8)	24
Georgia	0 (0)	4 (23.5)	13 (76.5)	24 ^c
Maryland	0 (0)	4 (16.0)	21 (84.0)	26
North Carolina	0 (0)	0 (0)	15 (100)	<21 ^b
South Carolina	0 (0)	0 (0)	3 (100)	14
Virginia	0 (0)	1 (6.7)	14 (93.3)	21
West Virginia	0 (0)	0 (0)	1 (100)	16

Region and state	Facilities offering only aspiration or surgical abortion, n (%)	Facilities offering only medication abortion, n (%)	Facilities offering both aspiration or surgical abortion and medication abortion, n (%)	Latest gestational limit (weeks since LMP ^a) as listed on website
East South Central	1 (6.7)	2 (13.3)	12 (80.0)	22 ^{b,c}
Alabama	1 (20.0)	0 (0)	4 (80.0)	21
Kentucky	0 (0)	0 (0)	1 (100)	22 ^{b,c}
Mississippi	0 (0)	0 (0)	1 (100)	16
Tennessee	0 (0)	2 (25.0)	6 (75.0)	<18
West South Central	1 (3.2)	3 (9.7)	27 (87.1)	<24 ^d
Arkansas	0 (0)	2 (66.7)	1 (33.3)	21
Louisiana	0 (0)	0 (0)	3 (100)	<19 ^{b,c}
Oklahoma	0 (0)	1 (25.0)	3 (75.0)	<22 ^b
Texas	1 (4.8)	0 (0)	20 (95.2)	<24 ^d
West	10 (3.8)	104 (39.7)	148 (56.5)	28
Mountain	4 (7.3)	17 (30.9)	34 (61.8)	28
Arizona	0 (0)	1 (12.5)	7 (87.5)	24
Colorado	2 (9.5)	8 (38.1)	11 (52.4)	26
Idaho	0 (0)	2 (50.0)	2 (50.0)	<16
Montana	0 (0)	2 (40.0)	3 (60.0)	21
Nevada	2 (25.0)	2 (25.0)	4 (50.0)	24
New Mexico	0 (0)	1 (20.0)	4 (80.0)	28
Utah	0 (0)	0 (0)	2 (100)	<22
Wyoming	0 (0)	1 (50.0)	1 (50.0)	12 ^b
Pacific	6 (2.9)	87 (42.0)	114 (55.1)	26
Alaska	0 (0)	1 (16.7)	5 (83.3)	<14
California	5 (3.3)	67 (44.1)	80 (52.6)	24
Hawaii	0 (0)	0 (0)	3 (100)	<24 ^b
Oregon	0 (0)	4 (33.3)	8 (66.7)	23
Washington	1 (2.9)	15 (44.1)	18 (52.9)	26

^aLMP: last menstrual period.

^bInformation on highest gestational limit obtained through phone call to facility.

^cInformation obtained in early 2017, and as of September 22, 2017, may no longer be accurate as a result of state laws or staffing changes.

^dOne facility in Texas listed the gestational limit on their website as <24 weeks; this was likely an error, as at the time of data collection, Texas had restrictions on abortion services after 22 weeks.


Abortion Deserts

We identified 27 “abortion deserts,” cities from which people would have to travel over 100 miles (160 km) to reach an abortion facility (Table 3). People living in Rapid City, SD had to travel the farthest, 318 miles (512 km), to reach an abortion

facility. Although the most cities in any one state (n=10) were located in Texas, there was a wide geographic diversity, with 15 unique states represented. These states were overwhelmingly in the South and Midwest. Figure 1 shows the geographic distribution of these distances, where large areas of the Midwest and Southwest had no abortion facility.

Table 3. Abortion deserts (cities >100 miles/160 km to closest facility) in the United States, May 2017.

City and state	2015 population, n	Distance to closest facility, miles (km)	Location of closest facility
1 Rapid City, SD	73,569	318 (512)	Billings, MT
2 Lubbock, TX	249,042	308 (496)	Fort Worth, TX
3 Midland, TX	132,950	293 (472)	Fort Worth, TX
4 Odessa, TX	118,968	282 (454)	El Paso, TX
5 Amarillo, TX	198,645	258 (415)	Warr Acres, OK
6 Casper, WY	60,285	223 (359)	Fort Collins, CO
7 San Angelo, TX	100,450	204 (328)	Austin, TX
8 Bismarck, ND	71,167	196 (315)	Fargo, ND
9 Laredo, TX	255,473	160 (257)	San Antonio, TX
10 Lake Havasu City, AZ	53,553	144 (232)	Henderson, NV
11 Abilene, TX	121,721	144 (232)	Fort Worth, TX
12 Corpus Christi, TX	324,074	139 (224)	San Antonio, TX
13 La Crosse, WI	52,306	137 (220)	Madison, WI
14 Springfield, MO	166,810	136 (219)	Fayetteville, AR
15 Lake Charles, LA	76,070	132 (212)	Baton Rouge, LA
16 Wichita Falls, TX	104,710	123 (198)	Fort Worth, TX
17 Columbia, MO	119,108	122 (196)	St Louis, MO
18 Evansville, IN	119,943	122 (196)	Bloomington, IN
19 St George, UT	80,202	121 (195)	Las Vegas, NV
20 Manhattan, KS	56,308	120 (193)	Overland Park, KS
21 Green Bay, WI	105,207	119 (192)	Milwaukee, WI
22 Pocatello, ID	54,441	114 (183)	Twin Falls, ID
23 Victoria, TX	67,574	107 (172)	San Antonio, TX
24 Fort Wayne, IN	260,326	106 (171)	Toledo, OH
25 Owensboro, KY	59,042	105 (169)	Louisville, KY
26 Dothan, AL	68,567	103 (166)	Montgomery, AL
27 Chattanooga, TN	176,588	101 (163)	Marietta, GA

Figure 1. Distance to nearest abortion facility in the contiguous United States, May 2017.

Discussion

Principal Results

Using an online search method, we identified almost 800 abortion facilities in the United States, which is consistent with other estimates of abortion clinics and nonspecialized clinics providing abortion [3]. These facilities were not distributed proportionately by state population. Through our analysis, we also found that 27 US cities, largely in the Midwest and the South, could be characterized as abortion deserts, as they did not have a publicly advertised abortion facility within 100 miles (160 km). These findings are consistent with those published by Bearak and colleagues [6], who found that the US counties where women would have to travel the farthest to reach the nearest abortion clinic were concentrated in the middle of the country, as well as several metropolitan areas in Texas. The lack of access to a common reproductive health service such as abortion is a public health concern in that more women in these cities could be forced to carry unwanted pregnancies to term if they are unable to travel long distances to obtain abortion care.

As states continue to pass, implement, and defend restrictions on abortion [8], it is possible that the number of abortion facilities will continue to decrease in those states with the most restrictions. The 6 states that have only 1 abortion facility have combined populations of almost 4 million women of reproductive age who will be forced to travel out of their home state to access abortion care if those facilities close.

For people seeking abortion services in the cities characterized as abortion deserts and in states with few facilities, reaching a facility for care could be incredibly challenging. Access to transportation is a barrier for people seeking all types of health care, in both urban and rural settings [28]. Lower-income women who are unable to access a car or money for gas may have to travel by bus, train, or other forms of transportation, which also becomes more difficult the farther they have to travel. Delays

in care due to distance or transportation can push women seeking abortion to later gestations [16,29,30] and are likely to disproportionately affect low-income women, who may struggle to cover the cost of transport [11,14]. Delays to abortion care may be particularly crucial to women in Wyoming, Alaska, Indiana, South Dakota, and South Carolina, where the abortion facilities had the lowest gestational limits. We found that 26.5% of identified facilities performed abortions at 20 weeks or later, which is lower than estimates from 2011-2012 [31], perhaps due to an increased number of state restrictions on abortion after 20 weeks since those estimates were published.

It seems likely that the larger number of facilities in the Northeast and West can be attributed to the fact that 40% to 50% of identified facilities in those regions are offering medication abortion only. The high proportion of facilities offering only medication abortion reflects the opportunities provided by medication abortion: the skills required for clinicians to provide it are minimal (compared with aspiration or surgical abortion) and the large majority of abortions in the United States (80.5%) occur at or before 10 weeks' gestation (the current accepted limit by which medication abortion can be provided) [32]. While the proportion of women choosing this method of abortion now accounts for 31% of nonhospital abortions (compared with 6% in 2001) [3], it is difficult to determine what the true demand would be if both medication and aspiration abortion were equally available. However, in states such as California, where fewer barriers to access exist for both types of abortion, medication abortion is now up to 46% of abortions in some populations, such as Medicaid recipients [33]. Additionally, states in the Northeast and West are less likely to have laws that limit the provision of medication abortion to physicians [34] and more likely to have policies that allow nurse practitioners, certified nurse midwives, and physicians assistants to offer medication abortion as part of their scope of practice.

These findings underscore the opportunities to pursue geographic expansion and other innovative models to achieve more equitable access to abortion care [35]. Some states have already managed this through an expansion of medication abortion-only services and increased use of telemedicine, which has been demonstrated to be safe and acceptable to women and to decrease travel for patients [36]. Indeed, in this analysis, Maine had the lowest ratio of women of reproductive age per facility, which was likely the result of an expansion of medication abortion through telemedicine programs offered from the existing Maine facilities [37]. While 19 states (almost exclusively in the South and Midwest) effectively prohibit telemedicine medication abortion [34], a recent Iowa Supreme Court decision could have implications for other states challenging similar restrictions that would allow expansion of medication abortion provision [38]. Given that it is less resource intensive, existing health care providers in the Midwest and the South, particularly in states where there is only 1 abortion provider or those states that contain cities classified as abortion deserts, could consider filling gaps in access by offering medication abortion alone as an entry point into abortion care, especially for primary care providers. Expanding the types of providers who can offer aspiration and medication abortion, such as nurse practitioners, would also increase the number of providers in smaller urban areas, thus expanding access to care [39]. However, it is important to note that in some states in which half or more facilities are only providing medication abortion, such as Idaho and Wyoming, the other facilities in the state offer abortion care up to 16 weeks and 12 weeks, respectively. Simply increasing the availability of medication abortion would not meet the needs of all women seeking abortion, some of whom may prefer aspiration abortion or need later abortion.

Supportive policy related to transportation for reproductive health services could also help reduce the burden on women in abortion deserts who have to travel extended distances to access services. California has recently introduced a Medicaid benefit to provide transportation for reproductive health care services (including abortion) to enrollees [40]. Further research is needed to determine what other policies can be enacted to reduce burdens of transportation and distance.

The internet will likely continue to be a key place for people to obtain the locations of abortion-providing facilities. However, both reduced geographic access and a desire to have more privacy and autonomy around the abortion process may lead women to seek out information on self-abortion [18] and obtain medication abortion pills through online sources, many of which have recently been shown to be selling effective medications with delivery to US-based mailing addresses [41]. There are no accurate estimates of how many women are obtaining abortion pills online, but the existence of online marketplaces and the documented feasibility of ordering from them implies online purchasing is occurring at volume.

Strengths and Limitations

This study is unique in that it systematically documented what people searching for abortion services online would find in search engine listings in early 2017 from a patient-centered perspective. A strength of this study is that it used up-to-date information on facilities of any volume to calculate distances, while other recent studies have been limited to a 2014 iteration of a proprietary database of only high-volume abortion-providing facilities (>400 abortions per year) that the Guttmacher Institute maintains [3,6]. An additional strength of the study is that it did not include abortion providers that offer abortion only to their existing patients or those that do not advertise their services, which would distort an accurate portrayal of the visibility of abortion availability. In addition, this analysis included the maximum gestations at which abortions were provided in each state, regardless of the state laws.

However, this study also has limitations. We used search terms that would enable us to locate abortion facilities in specific cities and states, but someone seeking abortion care might search “abortion clinic near me” and their results could vary from ours based on the location they are searching from. We attempted to eliminate geolocation bias by searching in Incognito mode and clearing both cache and cookies after each search. The information provided here is limited to what women seeking services would encounter—information that facilities chose to make available on their websites and provided through mystery shopper calls. Website information may be inaccurate or updated infrequently. It is possible that, if a woman called a facility to describe her unique situation, the staff could provide her with information about additional services that they do not wish to list on their website.

Distance is not the only barrier that people may face in trying to access abortion services—they may also face abortion stigma, waiting periods [13], and state gestational limits [16,42] as a result of state-level restrictions. In addition, the abortion facility that is closest to where a woman lives may not meet all her needs, particularly if it only provides medication abortion, has low limits on the gestation at which it provides abortion care, or cannot serve women with health conditions who may have higher risks.

Conclusions

Online searches provide information about abortion facilities and their services. The locations of these providers are not distributed equitably geographically across the United States. Having to travel long distances for abortion care can disproportionately affect low-income women and potentially push women to seek abortion at later gestations when care is even less available. Travel burdens may exist in addition to other restrictions on abortion in the state, including waiting periods and gestational limits, which can exacerbate inequities in the ability to access abortion care as part of the full range of reproductive health services.

Acknowledgments

We thank Ian Matthews, Redivis, for production of the map in this manuscript and Dr Sarah CM Roberts, University of California, San Francisco (UCSF), for her support in the initial development of this manuscript. This study was funded by a private, anonymous foundation. The funder had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. We also appreciate support from the UCSF Open Access Publishing Fund.

Authors' Contributions

AFC and UDU led the conceptual design of the study, analyzed and interpreted the data, and drafted the manuscript. MK and AFC led the collection of the data. JBW and NEJ contributed to the analysis and interpretation of the data and reviewed the manuscript for important intellectual content. All authors critically reviewed the paper drafts, gave final approval for the published draft, and are accountable for all aspects of the work.

Conflicts of Interest

None declared.

References

1. Foster DG, Biggs MA, Ralph L, Gerdtz C, Roberts S, Glymour MM. Socioeconomic outcomes of women who receive and women who are denied wanted abortions in the United States. *Am J Public Health* 2018 Mar;108(3):407-413. [doi: [10.2105/AJPH.2017.304247](https://doi.org/10.2105/AJPH.2017.304247)] [Medline: [29345993](https://pubmed.ncbi.nlm.nih.gov/29345993/)]
2. Roberts SC, Biggs MA, Chibber KS, Gould H, Rocca CH, Foster DG. Risk of violence from the man involved in the pregnancy after receiving or being denied an abortion. *BMC Med* 2014 Sep 29;12:144 [FREE Full text] [doi: [10.1186/s12916-014-0144-z](https://doi.org/10.1186/s12916-014-0144-z)] [Medline: [25262880](https://pubmed.ncbi.nlm.nih.gov/25262880/)]
3. Jones RK, Jerman J. Abortion incidence and service availability in the United States, 2014. *Perspect Sex Reprod Health* 2017 Mar;49(1):17-27 [FREE Full text] [doi: [10.1363/psrh.12015](https://doi.org/10.1363/psrh.12015)] [Medline: [28094905](https://pubmed.ncbi.nlm.nih.gov/28094905/)]
4. Johns NE, Foster DG, Upadhyay UD. Distance traveled for Medicaid-covered abortion care in California. *BMC Health Serv Res* 2017 Apr 19;17(1):287 [FREE Full text] [doi: [10.1186/s12913-017-2241-0](https://doi.org/10.1186/s12913-017-2241-0)] [Medline: [28420438](https://pubmed.ncbi.nlm.nih.gov/28420438/)]
5. Jones RK, Jerman J. How far did US women travel for abortion services in 2008? *J Womens Health (Larchmt)* 2013 Aug;22(8):706-713. [doi: [10.1089/jwh.2013.4283](https://doi.org/10.1089/jwh.2013.4283)] [Medline: [23863075](https://pubmed.ncbi.nlm.nih.gov/23863075/)]
6. Bearak JM, Burke KL, Jones RK. Disparities and change over time in distance women would need to travel to have an abortion in the USA: a spatial analysis. *Lancet Public Health* 2017 Nov;2(11):e493-e500. [doi: [10.1016/S2468-2667\(17\)30158-5](https://doi.org/10.1016/S2468-2667(17)30158-5)]
7. Jones RK, Jerman J. Characteristics and circumstances of U.S. women who obtain very early and second-trimester abortions. *PLoS One* 2017;12(1):e0169969 [FREE Full text] [doi: [10.1371/journal.pone.0169969](https://doi.org/10.1371/journal.pone.0169969)] [Medline: [28121999](https://pubmed.ncbi.nlm.nih.gov/28121999/)]
8. Nash E, Gold RB, Mohammed L, Ansari-Thomas Z, Cappello O. Policy trends in the States, 2017. New York, NY: Guttmacher Institute; 2018. URL: <https://www.guttmacher.org/article/2018/01/policy-trends-states-2017> [accessed 2018-04-30] [WebCite Cache ID 6xHYDHerL]
9. Gerdtz C, Fuentes L, Grossman D, White K, Keefe-Oates B, Baum SE, et al. Impact of clinic closures on women obtaining abortion services after implementation of a restrictive law in Texas. *Am J Public Health* 2016 May;106(5):857-864. [doi: [10.2105/AJPH.2016.303134](https://doi.org/10.2105/AJPH.2016.303134)] [Medline: [26985603](https://pubmed.ncbi.nlm.nih.gov/26985603/)]
10. Grossman D, White K, Hopkins K, Potter JE. Change in distance to nearest facility and abortion in Texas, 2012 to 2014. *JAMA* 2017 Dec 24;317(4):437-439. [doi: [10.1001/jama.2016.17026](https://doi.org/10.1001/jama.2016.17026)] [Medline: [28114666](https://pubmed.ncbi.nlm.nih.gov/28114666/)]
11. Roberts SCM, Fuentes L, Kriz R, Williams V, Upadhyay UD. Implications for women of Louisiana's law requiring abortion providers to have hospital admitting privileges. *Contraception* 2015 May;91(5):368-372. [doi: [10.1016/j.contraception.2015.02.001](https://doi.org/10.1016/j.contraception.2015.02.001)] [Medline: [25744615](https://pubmed.ncbi.nlm.nih.gov/25744615/)]
12. Roberts SC, Fuentes L, Kriz R, Williams V, Upadhyay UD. Corrigendum to "Implications for women of Louisiana's law requiring abortion providers to have hospital admitting privileges" [*Contraception* (2015) 91:368-72. doi: [10.1016/j.contraception.2015.02.001](https://doi.org/10.1016/j.contraception.2015.02.001)]. *Contraception* 2017 Feb;95(2):221-222. [doi: [10.1016/j.contraception.2016.08.015](https://doi.org/10.1016/j.contraception.2016.08.015)] [Medline: [27647624](https://pubmed.ncbi.nlm.nih.gov/27647624/)]
13. Roberts SCM, Turok DK, Belusa E, Combellick S, Upadhyay UD. Utah's 72-hour waiting period for abortion: experiences among a clinic-based sample of women. *Perspect Sex Reprod Health* 2016 Dec;48(4):179-187. [doi: [10.1363/48e8216](https://doi.org/10.1363/48e8216)] [Medline: [27010515](https://pubmed.ncbi.nlm.nih.gov/27010515/)]
14. Baum SE, White K, Hopkins K, Potter JE, Grossman D. Women's experience obtaining abortion care in Texas after implementation of restrictive abortion laws: a qualitative study. *PLoS One* 2016;11(10):e0165048 [FREE Full text] [doi: [10.1371/journal.pone.0165048](https://doi.org/10.1371/journal.pone.0165048)] [Medline: [27783708](https://pubmed.ncbi.nlm.nih.gov/27783708/)]
15. Jerman J, Frohwirth L, Kavanaugh ML, Blades N. Barriers to abortion care and their consequences for patients traveling for services: qualitative findings from two states. *Perspect Sex Reprod Health* 2017 Jun;49(2):95-102. [doi: [10.1363/psrh.12024](https://doi.org/10.1363/psrh.12024)] [Medline: [28394463](https://pubmed.ncbi.nlm.nih.gov/28394463/)]

16. Upadhyay UD, Weitz TA, Jones RK, Barar RE, Foster DG. Denial of abortion because of provider gestational age limits in the United States. *Am J Public Health* 2014 Sep;104(9):1687-1694. [doi: [10.2105/AJPH.2013.301378](https://doi.org/10.2105/AJPH.2013.301378)] [Medline: [23948000](https://pubmed.ncbi.nlm.nih.gov/23948000/)]
17. French V, Anthony R, Souder C, Geistkemper C, Drey E, Steinauer J. Influence of clinician referral on Nebraska women's decision-to-abortion time. *Contraception* 2016 Mar;93(3):236-243. [doi: [10.1016/j.contraception.2015.10.005](https://doi.org/10.1016/j.contraception.2015.10.005)] [Medline: [26593095](https://pubmed.ncbi.nlm.nih.gov/26593095/)]
18. Jerman J, Onda T, Jones RK. What are people looking for when they google "self-abortion"? *Contraception* 2018 Feb 22 [FREE Full text] [doi: [10.1016/j.contraception.2018.02.006](https://doi.org/10.1016/j.contraception.2018.02.006)] [Medline: [29477631](https://pubmed.ncbi.nlm.nih.gov/29477631/)]
19. Reis BY, Brownstein JS. Measuring the impact of health policies using internet search patterns: the case of abortion. *BMC Public Health* 2010;10:514 [FREE Full text] [doi: [10.1186/1471-2458-10-514](https://doi.org/10.1186/1471-2458-10-514)] [Medline: [20738850](https://pubmed.ncbi.nlm.nih.gov/20738850/)]
20. Stephens-Davidowitz S. The New York Times. 2016 Mar 05. The return of the D.I.Y. abortion URL: <https://www.nytimes.com/2016/03/06/opinion/sunday/the-return-of-the-diy-abortion.html> [accessed 2018-04-26] [WebCite Cache ID [6yyOKd5U9](https://www.webcitation.org/6yyOKd5U9)]
21. StatCounter. Search engine market share in United States of America. 2017. URL: <http://gs.statcounter.com/search-engine-market-share/all/united-states-of-america> [WebCite Cache ID [6vsle7oAp](https://www.webcitation.org/6vsle7oAp)]
22. U.S. Census Bureau Population Division. American Factfinder. 2016. Annual estimates of the resident population for incorporated places of 50,000 or more, ranked by July 1, 2015 population: April 1, 2010 to July 1, 2015 URL: <https://factfinder.census.gov/faces/tableservices/jsf/pages/productview.xhtml?src=bkmk> [accessed 2018-04-26] [WebCite Cache ID [6yyOXw6n4](https://www.webcitation.org/6yyOXw6n4)]
23. Robins S, Barr HJ, Idelson R, Lambert S, Zelkowitz P. Online health information regarding male infertility: an evaluation of readability, suitability, and quality. *Interact J Med Res* 2016 Oct 21;5(4):e25 [FREE Full text] [doi: [10.2196/ijmr.6440](https://doi.org/10.2196/ijmr.6440)] [Medline: [27769954](https://pubmed.ncbi.nlm.nih.gov/27769954/)]
24. Fast AM, Deibert CM, Hruby GW, Glassberg KI. Evaluating the quality of internet health resources in pediatric urology. *J Pediatr Urol* 2013 Apr;9(2):151-156. [doi: [10.1016/j.jpuro.2012.01.004](https://doi.org/10.1016/j.jpuro.2012.01.004)] [Medline: [22281281](https://pubmed.ncbi.nlm.nih.gov/22281281/)]
25. The Kenneth J. Ryan Residency Training Program in Abortion and Family Planning. Maps and location of Ryan program sites. 2017. URL: <https://www.ryanprogram.org/map-and-locations> [accessed 2017-12-21] [WebCite Cache ID [6vsm9IDOW](https://www.webcitation.org/6vsm9IDOW)]
26. Jacob S, Schifano N, Biard B. The mystery shopper: a tool to measure public service delivery? *Int Rev Adm Sci* 2016 Mar 14;84(1):164-184. [doi: [10.1177/0020852315618018](https://doi.org/10.1177/0020852315618018)]
27. OpenStreetMap. 2017. URL: <https://www.openstreetmap.org/> [accessed 2017-12-20] [WebCite Cache ID [6vq4PNy2b](https://www.webcitation.org/6vq4PNy2b)]
28. Syed ST, Gerber BS, Sharp LK. Traveling towards disease: transportation barriers to health care access. *J Community Health* 2013 Oct;38(5):976-993 [FREE Full text] [doi: [10.1007/s10900-013-9681-1](https://doi.org/10.1007/s10900-013-9681-1)] [Medline: [23543372](https://pubmed.ncbi.nlm.nih.gov/23543372/)]
29. Grossman D, Baum S, Fuentes L, White K, Hopkins K, Stevenson A, et al. Change in abortion services after implementation of a restrictive law in Texas. *Contraception* 2014 Nov;90(5):496-501 [FREE Full text] [doi: [10.1016/j.contraception.2014.07.006](https://doi.org/10.1016/j.contraception.2014.07.006)] [Medline: [25128413](https://pubmed.ncbi.nlm.nih.gov/25128413/)]
30. White K, deMartelly V, Grossman D, Turan JM. Experiences accessing abortion care in Alabama among women traveling for services. *Womens Health Issues* 2016;26(3):298-304. [doi: [10.1016/j.whi.2016.01.003](https://doi.org/10.1016/j.whi.2016.01.003)] [Medline: [26897655](https://pubmed.ncbi.nlm.nih.gov/26897655/)]
31. Jerman J, Jones RK. Secondary measures of access to abortion services in the United States, 2011 and 2012: gestational age limits, cost, and harassment. *Womens Health Issues* 2014;24(4):e419-e424 [FREE Full text] [doi: [10.1016/j.whi.2014.05.002](https://doi.org/10.1016/j.whi.2014.05.002)] [Medline: [24981401](https://pubmed.ncbi.nlm.nih.gov/24981401/)]
32. Jatlaoui TC, Ewing A, Mandel MG, Simmons KB, Suchdev DB, Jamieson DJ, et al. Abortion surveillance - United States, 2013. *MMWR Surveill Summ* 2016 Nov 25;65(12):1-44. [doi: [10.15585/mmwr.ss6512a1](https://doi.org/10.15585/mmwr.ss6512a1)] [Medline: [27880751](https://pubmed.ncbi.nlm.nih.gov/27880751/)]
33. Research and Analytics Studies Division, California Department of Health Care Services. Medi-Cal-funded induced abortions, 2014. Sacramento, CA: California Department of Health Care Services; 2016. URL: http://www.dhcs.ca.gov/dataandstats/statistics/Documents/Medi-Cal_Funded_Abortions_2014.pdf [WebCite Cache ID [6wipOASAK](https://www.webcitation.org/6wipOASAK)]
34. Guttmacher Institute. State policies in brief: medication abortion. New York, NY: Guttmacher Institute; 2017 Dec. URL: http://www.guttmacher.org/statecenter/spibs/spib_MA.pdf [accessed 2017-12-21] [WebCite Cache ID [6vsqpsBWF](https://www.webcitation.org/6vsqpsBWF)]
35. Upadhyay UD. Innovative models are needed for equitable abortion access in the USA. *Lancet Public Health* 2017 Nov;2(11):e484-e485. [doi: [10.1016/S2468-2667\(17\)30181-0](https://doi.org/10.1016/S2468-2667(17)30181-0)]
36. Grossman D, Grindlay K. Safety of medical abortion provided through telemedicine compared with in person. *Obstet Gynecol* 2017 Oct;130(4):778-782. [doi: [10.1097/AOG.0000000000002212](https://doi.org/10.1097/AOG.0000000000002212)] [Medline: [28885427](https://pubmed.ncbi.nlm.nih.gov/28885427/)]
37. Farwell J. Bangor Daily News. 2016 Feb 29. Maine Family Planning to offer telemedicine abortions URL: <https://bangordailynews.com/2016/02/29/health/maine-family-planning-to-offer-telemedicine-abortions/>
38. Yang YT, Kozhimannil KB. Medication abortion through telemedicine: implications of a ruling by the Iowa Supreme Court. *Obstet Gynecol* 2016 Feb;127(2):313-316 [FREE Full text] [doi: [10.1097/AOG.0000000000001251](https://doi.org/10.1097/AOG.0000000000001251)] [Medline: [26942359](https://pubmed.ncbi.nlm.nih.gov/26942359/)]
39. Weitz TA, Taylor D, Desai S, Upadhyay UD, Waldman J, Battistelli MF, et al. Safety of aspiration abortion performed by nurse practitioners, certified nurse midwives, and physician assistants under a California legal waiver. *Am J Public Health* 2013 Mar;103(3):454-461. [doi: [10.2105/AJPH.2012.301159](https://doi.org/10.2105/AJPH.2012.301159)] [Medline: [23327244](https://pubmed.ncbi.nlm.nih.gov/23327244/)]

40. Chen A. Q&A on non-medical transportation and reproductive health services. Los Angeles, CA: National Health Law Program; 2017 Aug 04. URL: <http://www.healthlaw.org/issues/reproductive-health/q-a-on-non-medical-transportation-and-reproductive-health-services#.WjwVs1WnHIU> [WebCite Cache ID 6vsptYDTA]
41. Murtagh C, Wells E, Raymond EG, Coeytaux F, Winikoff B. Exploring the feasibility of obtaining mifepristone and misoprostol from the internet. *Contraception* 2018 Apr;97(4):287-291 [FREE Full text] [doi: [10.1016/j.contraception.2017.09.016](https://doi.org/10.1016/j.contraception.2017.09.016)] [Medline: [29030227](https://pubmed.ncbi.nlm.nih.gov/29030227/)]
42. Roberts SCM, Gould H, Upadhyay UD. Implications of Georgia's 20-week abortion ban. *Am J Public Health* 2015 Aug;105(8):e77-e82. [doi: [10.2105/AJPH.2015.302728](https://doi.org/10.2105/AJPH.2015.302728)] [Medline: [26066924](https://pubmed.ncbi.nlm.nih.gov/26066924/)]

Abbreviations

LMP: last menstrual period

Edited by G Eysenbach; submitted 22.12.17; peer-reviewed by R Jones, S Robins, R Pankomera; comments to author 11.01.18; revised version received 08.03.18; accepted 02.04.18; published 14.05.18

Please cite as:

Cartwright AF, Karunaratne M, Barr-Walker J, Johns NE, Upadhyay UD

Identifying National Availability of Abortion Care and Distance From Major US Cities: Systematic Online Search

J Med Internet Res 2018;20(5):e186

URL: <http://www.jmir.org/2018/5/e186/>

doi: [10.2196/jmir.9717](https://doi.org/10.2196/jmir.9717)

PMID: [29759954](https://pubmed.ncbi.nlm.nih.gov/29759954/)

©Alice F Cartwright, Mihiri Karunaratne, Jill Barr-Walker, Nicole E Johns, Ushma D Upadhyay. Originally published in the Journal of Medical Internet Research (<http://www.jmir.org>), 14.05.2018. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in the Journal of Medical Internet Research, is properly cited. The complete bibliographic information, a link to the original publication on <http://www.jmir.org/>, as well as this copyright and license information must be included.