

RESEARCH ARTICLE

Gender-Specific Association between Tobacco Smoking and Central Obesity among 0.5 Million Chinese People: The China Kadoorie Biobank Study

Jun Lv¹, Wei Chen^{1,2}, Dianjianyi Sun^{1,2}, Shengxu Li^{1,2}, Iona Y. Millwood³, Margaret Smith³, Yu Guo⁴, Zheng Bian⁴, Canqing Yu¹, Huiyan Zhou⁴, Yunlong Tan⁴, Junshi Chen⁵, Zhengming Chen^{3*}, Liming Li^{1,4*}, China Kadoorie Biobank collaborative group

1 Department of Epidemiology and Biostatistics, School of Public Health, Peking University Health Science Center, Beijing, China, **2** Department of Epidemiology, School of Public Health and Tropical Medicine, Tulane University, New Orleans, Louisiana, United States of America, **3** Clinical Trial Service Unit & Epidemiological Studies Unit (CTSU), Nuffield Department of Population Health, University of Oxford, United Kingdom, **4** Chinese Academy of Medical Sciences, Beijing, China, **5** China National Center for Food Safety Risk Assessment, Beijing, China

* lmlee@vip.163.com (LML); zhengming.chen@ctsu.ox.ac.uk (ZMC)



OPEN ACCESS

Citation: Lv J, Chen W, Sun D, Li S, Millwood IY, Smith M, et al. (2015) Gender-Specific Association between Tobacco Smoking and Central Obesity among 0.5 Million Chinese People: The China Kadoorie Biobank Study. PLoS ONE 10(4): e0124586. doi:10.1371/journal.pone.0124586

Academic Editor: C. Mary Schooling, Hunter College, UNITED STATES

Received: July 22, 2014

Accepted: March 5, 2015

Published: April 21, 2015

Copyright: © 2015 Lv et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: The authors do not own the data for this study. Requests for the data should be submitted to the China Kadoorie Biobank (CKB) Data Access Committee (<http://www.ckbiobank.org/site/Research/Data+Access+Policy>). As stated in the policy, as data custodian, the CKB study group must maintain the integrity of the database for future use and regulate data access to comply with prior conditions agreed with the Chinese government. Data security is an integral part of CKB study protocols. Data can be released outside the

Abstract

Objectives

Lifestyle factors are well-known important modifiable risk factors for obesity; the association between tobacco smoking and central obesity, however, is largely unknown in the Chinese population. This study examined the relationship between smoking and central obesity in 0.5 million Chinese adults, a population with a low prevalence of general obesity, but a high prevalence of central obesity.

Subjects

A total of 487,527 adults (200,564 males and 286,963 females), aged 30-79 years, were enrolled in the baseline survey of the China Kadoorie Biobank (CKB) Study conducted during 2004-2008. Waist circumference (WC) and WC/height ratio (WHtR) were used as measures of central obesity.

Results

The prevalence of regular smokers was significantly higher among males (60.6%) than among females (2.2%). The prevalence of central obesity increased with age and BMI levels, with a significant gender difference (females>males). Of note, almost all obese adults (99.4%) were centrally obese regardless of gender. In multivariable regression analyses, adjusting for age, education, physical activity, alcohol use and survey site, regular smoking was inversely associated with BMI in males (standardized regression coefficients, $\beta = -0.093$, $p < 0.001$) and females ($\beta = -0.025$, $p < 0.001$). Of interest, in the BMI stratification analyses in

CKB research group only with appropriate security safeguards.

Funding: This work was supported by the CKB baseline survey and the first re-survey were supported by the Kadoorie Charitable Foundation in Hong Kong. The long-term follow-up of the project during 2009-14 is supported by the Wellcome Trust in the UK [088158/Z/09/Z] and the National Key Technologies Research and Development Program in the 12th Five-year Plan, Chinese Ministry of Science and Technology [2011BAI09B01, 2012-14]. Support for this analysis is provided by the National Natural Science Foundation of China [No. 81390541]. The sponsors had no role in the design, analysis, interpretation or drafting of this manuscript.

Competing Interests: The authors have declared that no competing interests exist.

18 groups, all β s of regular smoking for WHtR were positive in both genders; the β s showed a significantly greater increasing trend with increasing BMI in males than in females. In the analyses with model adjustment for BMI, the positive associations between regular smoking and WHtR were stronger in males ($\beta=0.021$, $p<0.001$) than in females ($\beta=0.008$, $p<0.001$) ($p<0.001$ for gender difference). WC showed considerably consistent results.

Conclusions

The data indicate that tobacco smoking is an important risk factor for central obesity, but the association is gender-specific and depends on the adjustment for general obesity.

Introduction

Obesity has reached an epidemic level worldwide in both children and adult populations despite efforts of research and prevention for decades [1–5]. Obesity has gained importance because of its association with subsequent cardiovascular disease and all-cause mortality [6–8]. Obesity has also been shown to be associated with a number of cardiovascular disease risk factors, such as hypertension, dyslipidemia, type 2 diabetes, and insulin resistance, all known as metabolic syndrome components [9–12]. Asian populations have a relatively lower BMI compared with Western populations, but are predisposed to central or abdominal obesity [7,11–16]. This central fat accumulation is reported to be more strongly associated with cardiovascular disease risk factors and metabolic syndrome than peripheral fat distribution [11,12].

The rapid economic growth has been accompanied by urbanization, westernization of lifestyle behaviors and rising rates of obesity in China during the past couple of decades [4,5,17,18]. Based on secular trends in the prevalence of obesity among Chinese adults during 1993–2009, it seems that the obesity epidemic has not plateaued in China [5]. Central obesity and related cardiometabolic abnormalities have been considered an expression of a "Civilization Syndrome" [19]. The modern, urbanized society-related features, including unhealthy dietary patterns, lower physical activity, alcohol consumption and tobacco smoking, are well-known important modifiable lifestyle risk factors for obesity [1,17–21]. Despite a huge body of literature on the inverse association between smoking and general obesity [22–24], data are limited regarding the association between tobacco smoking and central obesity in Asians; no previous studies have focused on the smoking-central obesity association among adults in the Chinese population, with a low prevalence of general obesity, but a high prevalence of central obesity. The objective of the present study is to examine the relationship between tobacco smoking and central obesity measures in 0.5 million male and female adults enrolled in the China Kadoorie Biobank (CKB) Study.

Methods

Study Design and Participants

The CKB Study is a prospective study of chronic disease in China. Details of the CKB study design and characteristics of the study participants have been described elsewhere [25,26]. Briefly, 512,891 participants without major disabilities living in administrative units (rural villages or urban residential committees), aged 30–79 years (mean age: 51.5 years), were recruited in the baseline survey between 2004 and 2008 from five urban and five rural areas in China. Selection

of the survey sites was based on local patterns of disease and exposure to certain risk factors, population stability, quality of death and disease registries, local commitment and capacity.

The current analysis included participants who had a weight between 30 kg and 160 kg, a height of 145–200 cm (males) and 140–200 cm (females), a BMI between 18.5 and 45.0, and a waist circumference between 50 cm and 150 cm. People who were very underweight or extremely obese or were at the extremes of the height distribution may have underlying metabolic or growth disorders. A total of 25,364 participants were excluded based on the above selection criteria, and 487,527 adults (200,564 males and 286,963 females) formed the sample for the current analysis.

Ethical approval for the CKB study was obtained from the Ethical Review Committee of the Chinese Center for Disease Control and Prevention (Beijing, China) and the Oxford Tropical Research Ethics Committee, University of Oxford (UK). In addition, approvals were obtained from the institutional research boards at the local Center for Disease Control and Prevention in each of the ten survey sites. Finally, written informed consent was obtained from all participants.

Questionnaire survey

In the baseline survey, trained interviewers administered a standardized questionnaire using a laptop-based direct data-entry system, with built-in functions to avoid logical errors and missing items. The questionnaire included detailed questions on socio-demographic status, medical history and health status, smoking, alcohol consumption, physical activity, and other lifestyle behaviors.

Education was classified into five levels: Illiterate, elementary, middle school, high school, college and above. Total physical activity was converted into metabolic equivalent hours per day (MET-hours/day) spent on work, transportation, housework, non-sedentary recreation as described in our previous study [27]. For assessment of alcohol consumption, participants were asked how often they had drunk alcohol during the previous 12 months, and those who had not drunk weekly were asked if there was a period of at least a year prior to that when they had drunk some alcohol at least once a week. Participants were classified into four main drinking categories: 1) nondrinkers; 2) ex-drinkers; 3) occasional drinkers; and 4) weekly drinkers.

For assessment of tobacco consumption, participants were asked how often they had smoked tobacco at the time of the survey, and those who did not smoke or only smoked occasionally were asked if there was a period prior to that when they had smoked some tobacco on most days or daily. In the present analysis, participants were classified into four main smoking categories: 1) nonsmokers were defined as those who did not smoke currently and had not smoked more than 100 cigarettes during his/her lifetime; 2) ex-smokers as those who currently do not smoke or only smoked occasionally but had smoked on most days or daily in the past; 3) occasional smokers as those who do not smoke currently but had smoked occasionally in the past or had smoked at least 100 cigarettes during his/her lifetime, or those who currently smoke occasionally and had not smoked on most days or daily in the past; 4) regular smokers as those who currently smoke daily or on most days. Information on duration (years) of smoking and number of packs smoked per day was obtained for regular smokers, and pack-years was calculated as a measure of cumulative burden of smoking.

Anthropometric measurements

Standing height and body weight were measured in light indoor clothing without shoes to the nearest 0.1 cm and 0.1 kg, respectively. Waist circumference (WC) was measured midway between the iliac crest and the lower rib margin at the end of normal expiration using a plastic

flexible tape to the nearest 0.1 cm. All measurements were made by trained staff using a standard protocol and instruments. WC/height ratio (WHtR) was calculated as WC divided by height and was used as a measure of central obesity [12]. Body mass index (BMI) was calculated as weight in kilograms divided by the square of height in meters and was used as a measure of general obesity. Normal-weight (BMI = 18.5–23.9), over weight (BMI = 24.0–27.9) and obesity (BMI = 28.0–45.0) groups were defined based on recommendations of the Working Group on Obesity in China [7]. Central obesity was defined as WHtR > 0.5 [11,12].

Statistical analyses

Statistical analyses were performed using Stata Statistical Software (version 13.1, 2013, Stata-Corp LP., College Station, TX). All analyses were performed separately by gender groups. Descriptive data were presented as mean and standard deviation (SD) for continuous variables, and percentages for categorical variables. General linear models were used to test differences in continuous study variables between gender groups, adjusting for age, and survey sites for the descriptive data. Chi-square test was used to test the differences in categorical study variables between gender groups.

Multivariable linear regression analyses were performed (dependent variables = BMI, WC or WHtR; predictor = smoking) by gender, adjusting for age, survey site (included in the regression models as nine dummy variables using Qingdao as a reference), education (categorized as no formal school, elementary school, middle school, high school, or college and above), total physical activity (MET-hours/day) and alcohol use (nondrinker, ex-drinker, occasional drinker and weekly drinker). Smoking was included in the model as both a categorical variable (nonsmoking, ex-smoking, occasional smoking and regular smoking) and a continuous variable (pack-years). In order to assess the influence of adjustment for BMI on the association of tobacco smoking with WC and WHtR, males and females were stratified into 18 subgroups according to levels of BMI with equal numbers of subjects in each subgroup. Multivariable linear regression analyses were performed within each subgroup, adjusting for the above mentioned covariates. Standardized regression coefficients of regular smoking for WC and WHtR in the 18 BMI subgroups were plotted in males vs females. Linear regression models were used to assess significance of trends in the effect size of regular smoking on WC and WHtR across these 18 BMI subgroups; interaction regression models were used to test the difference in slopes of the regression lines between gender groups.

In sensitivity analyses, the association of tobacco smoking with WC and WHtR was examined using multivariable linear regression models, with and without BMI adjustment, by groups of normal-weight (BMI = 18.5–23.9), over-weight (BMI = 24.0–27.9), and obesity (BMI = 28.0–45.0), adjusting for covariates as mentioned above.

Results

[Table 1](#) summarizes the characteristics of the study cohort by gender. Males were older and had higher education than females ($p < 0.001$); males had significantly higher levels of physical activity measured as MET-hours/day and prevalence of weekly alcohol drinkers than females. The prevalence of regular smokers was 27.5 times higher among males than that among females (60.6% vs 2.2%). Females had longer duration of smoking but lower packs/day and pack-years than males. Height, weight, and WC showed significant gender differences in mean values (males > females); however, BMI, WHtR, and the prevalence of central obesity were significantly lower among males than among females. Further, the prevalence of central obesity increased in the order of subgroups of normal-weight (BMI = 18.5–23.9), over-weight (BMI = 24.0–27.9), and obesity (BMI = 28.0–45.0) in both males and females, with significant

Table 1. Characteristics (mean±SD or %) of study variables by gender groups.

	Males (n = 200,564)	Females (n = 286,963)	P for gender difference
Age (year)	52.1 ± 10.8	50.7 ± 10.3	<0.001
Education (%)			<0.001
Illiterate and elementary	41.3	55.9	
Middle school and above	58.7	44.1	
Physical activity (MET-hours/day)^a	22.2 ± 15.3	20.5 ± 12.8	<0.001
Alcohol use (%)			<0.001
Nondrinker	19.7	63.0	
Ex-regular drinker	8.7	0.9	
Occasional drinker	38.0	34.1	
Regular drinker	33.7	2.0	
Tobacco use			<0.001
Nonsmoker	14.5	95.2	
Ex-regular smoker	13.4	0.8	
Occasional smoker	11.4	1.8	
Regular smoker	60.6	2.2	
Regular smoker			
Duration of smoking (years)	28.4 ± 10.9	29.6 ± 15.5	<0.001
Packs/day	0.9 ± 0.5	0.5 ± 0.4	<0.001
Pack-years/1000	9.5 ± 6.9	5.5 ± 5.4	<0.001
Height (cm)^a	165.3 ± 6.4	154.4 ± 5.7	<0.001
Weight (kg)^a	65.0 ± 10.4	57.5 ± 8.9	<0.001
BMI (kg/m²)^a	23.7 ± 3.1	24.1 ± 3.2	<0.001
Waist circumference (cm)^a	82.7 ± 9.4	79.8 ± 9.1	<0.001
WHtR ratio^a	0.500 ± 0.054	0.517 ± 0.060	<0.001
Central obesity (%)^b	48.8	58.2	<0.001
Central obesity (%)^b			
BMI 18.5–23.9	19.1	30.3	<0.001
BMI 24.0–27.9	84.0	85.8	<0.001
BMI 28.0–45.0	99.4	99.4	0.878

^a: P-values for gender difference were adjusted for age and survey site;

^b: Central obesity was defined as WHtR≥0.5.

MET-hours/day: metabolic equivalent hours per day

WHtR: waist circumference/height ratio

doi:10.1371/journal.pone.0124586.t001

gender differences in the normal-weight and over-weight groups (females>males). Of note, almost all Chinese adults (99.4%) with obesity in this cohort were centrally obese regardless of gender.

Fig 1. presents the prevalence of central obesity by gender and age in the total sample (A) and among individuals with normal-weight (B). In the total sample, the prevalence of central obesity increased with age, and females showed a significantly higher prevalence than males at ages above 45 years. Among normal-weight individuals, the prevalence of central obesity increased with age and was significantly higher among females than among males in all age groups, with a steeper increase in females than in males.

Table 2 shows mean levels of BMI, WC and WHtR by gender and smoking groups. Compared to nonsmokers, regular smokers had significantly lower values of BMI in males and females; the differences in WC and WHtR, however, showed an opposite direction in meals




Fig 1. Prevalence of central obesity in the total sample (BMI = 18.5–45.0) (A) and normal-weight adults (BMI = 18.5–23.9) (B).

doi:10.1371/journal.pone.0124586.g001

(regular smokers < nonsmokers) and females (regular smokers > nonsmokers). It has to be noted that the difference in unadjusted BMI between nonsmokers and regular smokers was greater in males (24.13–23.4 = 0.73) than in females (24.10–23.9 = 0.20). Apparently, the lower BMI resulted, at least partly, in lower WC and WHtR in male regular smokers.

Table 3 presents linear regression coefficients of smoking for BMI, WC and WHtR by gender groups, adjusting for age, education, physical activity, alcohol use and survey site. Compared with non-smoking, ex-smoking was positively associated with BMI, WC and WHtR in males and females ($p < 0.001$), but regular smoking was inversely associated with BMI, WC and WHtR in both gender groups ($p < 0.001$). Occasional smoking was positively associated with all adiposity measures in males and females except for BMI in males. Pack-years was inversely associated with BMI, WC and WHtR in both gender groups ($p < 0.001$) except for WC in males (positively). Gender differences in standardized regression coefficients of ex-smoking and regular smoking for WC and WHtR (males > females) were all significant; gender differences in standardized regression coefficients of pack-years were significant for BMI (males > females) and WC (an opposite direction). The regression coefficients of pack-years vs regular smoking for WC had different signs (0.006 vs -0.044) among males. The explanation is that the associations of light and heavy regular smoking and the duration of smoking with BMI may be different, and the large sample is highly sensitive to detect the change in the effect. Another

Table 2. Unadjusted means ± SD of adiposity measures by gender and smoking groups.

	Males			Females		
	BMI(kg/m ²)	WC(cm)	WHtR	BMI(kg/m ²)	WC(cm)	WHtR
Nonsmoker	24.13±3.0	83.8±9.3	0.507 ± 0.053	24.10±3.2	79.7±9.1	0.516 ± 0.060
Ex-smoker	24.4±3.1	85.0±9.7	0.515 ± 0.056	24.9±3.5	83.6±9.8	0.552 ± 0.063
Occasional smoker	24.1±3.0	83.4±9.0	0.505 ± 0.052	24.3±3.4	80.3±9.3	0.524 ± 0.061
Regular smoker	23.4±3.0	81.9±9.3	0.495 ± 0.053	23.9±3.4	80.4±9.6	0.528 ± 0.061
P-value ^a	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001

WC: waist circumference

WHtR: waist circumference/height ratio

^a: difference between nonsmokers and regular smokers, adjusted for age and survey sites.

Gender differences in means were all significant ($P < 0.001$), adjusted for age and survey sites.

doi:10.1371/journal.pone.0124586.t002

Table 3. Standardized (upper) and unstandardized (lower) linear regression coefficients (95% confidence interval) for BMI, WC and WHtR by gender groups.

Independent variable	BMI		WC ^a		WHtR ^a	
	Males	Females	Males	Females	Males	Females
Model 1						
Non-smoking (reference)	—	—	—	—	—	—
Ex-smoking	0.037**	0.009**†	0.057**	0.015**†	0.049**	0.014**†
	0.328**	0.323**	1.567**	1.488**	0.008**	0.009**
	(0.280, 0.377)	(0.195, 0.451)	(1.418, 1.715)	(1.135, 1.840)	(0.007, 0.009)	(0.007, 0.012)
Occasional smoking	-0.001	0.006**	0.006*	0.009**	0.007*	0.009**
	-0.006	0.144**	0.181*	0.612**	0.001*	0.004**
	(-0.056, 0.045)	(0.056, 0.233)	(0.027, 0.334)	(0.370, 0.854)	(0.0002, 0.002)	(0.003, 0.006)
Regular smoking	-0.093**	-0.025**†	-0.044**	-0.010**†	-0.061**	-0.011**†
	-0.582**	-0.548**	-0.853**	-0.626**	-0.007**	-0.005**
	(-0.620, -0.544)	(-0.629, -0.467)	(-0.970, -0.736)	(-0.849, -0.403)	(-0.007, -0.006)	(-0.006, -0.003)
Model 2^b						
Pack-years/1000	-0.031**	-0.022**†	0.006*	-0.008**†	-0.007*	-0.010**
	-0.013**	-0.062**	0.007*	-0.066**	-0.00005*	-0.001**
	(-0.015, -0.011)	(-0.073, -0.052)	(0.001, 0.014)	(-0.095, -0.038)	(-0.00009, -0.00001)	(-0.001, -0.0003)

WC: waist circumference

WHtR: waist circumference/height ratio

* p<0.05,

** p<0.01 for coefficients different from 0

† P<0.05 for gender difference

^a: BMI was not included in the models for adjustment.

^b: Only nonsmokers and regular smokers were included.

Age, education, physical activity, alcohol use and survey site were included in models for adjustment.

doi:10.1371/journal.pone.0124586.t003

explanation is that body height might have an influence in this regard when compared with the regression coefficient for WHtR (-0.007*, p = 0.011) in males.

In order to assess the influence of adjustment for BMI on the association analyses of smoking with WC and WHtR, males and females were stratified into 18 subgroups with equal numbers of subjects in each subgroup according to levels of BMI. Fig 2 depicts standardized regression coefficients of regular smoking for WHtR (A) and WC (B) by 18 BMI groups among males and females, adjusted for age, education, physical activity, alcohol use and survey site. The number of adults was 11,142–11,143 in the male subgroups and 15,942–15,943 in the female subgroups. The standardized regression coefficients of regular smoking were all positive in males and females and showed a significant increasing trend with increasing BMI for WHtR and WC in males, but not in females; the slopes of the regression lines were significantly different between males and females. It should be noted that the association parameters in the BMI stratification analysis, a powerful and reliable adjustment method, were completely different (opposite) from those in the regression analyses for WC and WHtR in Table 3 without BMI adjustment in the model.

In order to confirm the results of BMI stratification analyses, a model adjustment for BMI was used. Table 4 shows regression coefficients of smoking for WC and WHtR. When BMI was included in the regression models, both regular smoking and pack-years showed positive associations with WC and WHtR in males and females. The differences in standardized regression




Fig 2. Standardized regression coefficients of regular smokers for WHtR (A) and WC (B) by 18 BMI groups in males and females: Standardized regression coefficients greater than 0.026 for male smokers and 0.015 for female smokers were significant ($p < 0.05$).

doi:10.1371/journal.pone.0124586.g002

coefficients of regular smoking and pack-years between males and females in Table 4 were all significant for both WC and WHtR (males > females, $p < 0.01$). The results of the model adjustment for BMI with respect to regression coefficients of regular smoking were substantially consistent with those of the BMI stratification analyses shown in Fig 2.

Table 4. Standardized (upper) and unstandardized (lower) linear regression coefficients (95% confidence interval) for WC and WHtR by gender groups, with adjustment for BMI.

Independent variable	WC		WHtR	
	Males	Females	Males	Females
Model 1				
Nonsmoking	—	—	—	—
Ex-smoking	0.026** (0.639, 0.795)	0.008**† (0.555, 0.952)	0.017** (0.002, 0.003)	0.007**† (0.003, 0.006)
Occasional smoking	0.007** (0.114, 0.276)	0.004** (0.147, 0.420)	0.007** (0.001, 0.002)	0.005** (0.001, 0.003)
Regular smoking	0.034** (0.593, 0.716)	0.010**† (0.494, 0.745)	0.020** (0.002, 0.003)	0.008**† (0.003, 0.004)
Model 2^a				
Pack-years/1000	0.032** (0.038, 0.045)	0.010**† (0.059, 0.091)	0.021** (0.0001, 0.0002)	0.008**† (0.0003, 0.001)

WC: waist circumference

WHtR: waist circumference/height ratio

* $p < 0.05$,

** $p < 0.01$ for coefficients different from 0

† $P < 0.05$ for gender difference

^a: Only nonsmokers and regular smokers were included.

Age, education, physical activity, alcohol use, survey site and BMI were included in models for adjustment.

doi:10.1371/journal.pone.0124586.t004




Fig 3. Prevalence of central obesity by 18 BMI subgroups in males and females.

doi:10.1371/journal.pone.0124586.g003

Fig 3. illustrates the prevalence of central obesity by 18 BMI subgroups and gender. The prevalence of central obesity increased with increasing BMI at a faster rate at BMI levels below 24.0 in both males and females; the prevalence almost reached the ceiling in subgroups with a mean BMI above 28.0. Further, Pearson correlation coefficients between WHtR and BMI were 0.641, $p < 0.001$ (males) and 0.550, $p < 0.001$ (females) in the normal-weight group (BMI = 18.5–23.9); 0.487, $p < 0.001$ (males) and 0.416, $p < 0.001$ (females) in the over-weight group (BMI = 24.0–27.9); and 0.575, $p < 0.001$ (males) and 0.542, $p < 0.001$ (females) in the obesity group (BMI = 28.0–45.0). The positive BMI-WHtR correlations and negative smoking-BMI association indicated the importance of adjustment for BMI in the smoking-central obesity association analyses, especially in the normal-weight group.

In sensitivity analyses, the association between smoking and WHtR was examined in linear regression models by groups of normal-weight (BMI = 18.5–23.9), over-weight (BMI = 24.0–27.9) and obesity (BMI = 28.0–45.0), without and with adjusting for BMI along with other covariates. In males, the standardized linear regression coefficients of regular smoking for WC were -0.009* without adjusting for BMI and 0.040** with adjusting for BMI in the normal-weight group, 0.067** and 0.077** in the over-weight group and 0.066** and 0.072** in the obesity group (* $p < 0.05$; ** $p < 0.01$). In females, the standardized linear regression coefficients of regular smoking for WC were 0.001 without adjusting for BMI and 0.016** with adjusting for BMI in the normal-weight group, 0.016** and 0.018** in the over-weight group and 0.028** and 0.020** in the obesity group. In males, the standardized linear regression coefficients of regular smoking for WHtR were -0.032** without adjusting for BMI and 0.019** with adjusting for BMI in the normal-weight group, 0.043** and 0.054** in the over-weight group and 0.049** and 0.055** in the obesity group. In females, the standardized linear regression coefficients of regular smoking for WHtR were 0.001 without adjusting for BMI and 0.015** with adjusting for BMI in the normal-weight group, 0.013** and 0.015** in the over-weight group and 0.021** and 0.014** in the obesity group. The results of these sensitivity analyses indicated that the negative associations of regular smoking with WC and WHtR without adjusting for BMI were mainly because of the association patterns in the normal-weight group in males and females. The number of study subjects of the normal-weight group accounted for 54.2% of the entire study cohort.

Discussion

Asian populations have a relatively lower BMI, but a greater prevalence of central obesity than Caucasian populations [5,7,11–16,28]. The CKB study, one of the largest cohort studies of

chronic disease in the world, provides a great opportunity to investigate tobacco smoking and central obesity in Chinese population. In the current study, the prevalence of central obesity increased with increasing values of BMI, and female adults had a significantly higher prevalence of central obesity than male adults, especially in the normal-weight group. Of note, almost all Chinese adults (99.4%) with obesity (BMI = 28.0–45.0) in this cohort were centrally obese regardless of gender. Compared with males, females had a much lower prevalence of regular smokers (2.2% vs 60.6%), Regular smoking was associated with decreased BMI and increased WC and WHtR in both males and females when BMI was adjusted for in the model, and the association of smoking with WC, WHtR was much weaker in females than in males. These results indicate that female central obesity may have other important underlying mechanisms and risk factors in addition to tobacco smoking.

Epidemiologic studies have shown that body weight tends to be lower among smokers than among nonsmokers in many populations [22–24]. Smoking abstinence results in an increase in body weight for both males and females, and smoking is often considered as a way to control appetite and body weight [29]. The mechanisms that underlie this smoking-related weight phenomenon are complex and involve multiple neurochemical pathways. Most of the effects of smoking on body weight are mediated by nicotine inhaled from cigarette smoke. Nicotine acutely increases the levels of various neurotransmitters, such as the systemic release of catecholamines, dopamine and serotonin in the brain, suppresses appetite and consequently reduce food intake [23,30]. This process likely explains why smokers tend to decrease body weight, and why smoking cessation is frequently followed by weight gain [29]. As a result, the belief that cigarette smoking is helpful in controlling body weight has long been part of popular culture. Despite the universally accepted concept in this regard, there has been growing evidence that smoking is associated with abdominal fat accumulation; cross-sectional studies indicate that the prevalence of central obesity is higher in smokers than in nonsmokers [24,31–35]. Waist-to-hip ratio is positively associated with the number of pack-years of smoking [34], and there is a dose-response relation between waist-to-hip ratio and the number of cigarettes smoked [35]. The findings from the present study on the significantly inverse smoking-BMI association and positive smoking cessation-BMI association in both males and females are in concordance with observations from previous studies [22–24]. However, the association of regular smoking with WC and WHtR observed in the current study was dependent on the adjustment for BMI, especially in male normal-weight adults. After adjusting for BMI, regular smoking and the cumulative burden of smoking were significantly and positively associated with WC and WHtR; stratification analysis and model adjustment methods showed consistent results in this regard. The findings from the present study support the previous findings on the smoking-central obesity association [31–35]. Of interest, an important question has been raised in this study regarding whether BMI has to be adjusted for in the smoking-central obesity association analysis, particularly among normal-weight male adults.

By comparing the regression coefficients before and after adjustment for BMI in the model, completely opposite directions in the associations of regular smoking with WC and WHtR were noted in the current study. When BMI was not included in the model, current smoking showed a protective effect on WC and WHtR; whereas, current smoking was positively associated with these central obesity measures with BMI included in the model. The changes in the strength and direction of the association parameters suggested a considerable influence of BMI on these associations, particularly in the male normal-weight group as shown in the sensitivity analyses. When BMI was not adjusted for in regression analysis models, the negative association of regular smoking with WC ($\beta = -0.009$) and WHtR ($\beta = -0.032$) largely reflect the effect of smoking on BMI in the male normal-weight group. The smoking-BMI-central obesity relationship is similar to the mediation model proposed by Hernández-Díaz (Fig 3, panel 3.3) [36].

In the current analysis, the smoking→BMI indirect effect (-) and BMI↔WHtR correlation (+) are so strong that the absolute value of the negative total indirect effect (the product of the two indirect path effects) is supposed to be much greater than the positive value of smoking→WHtR direct effect (+). Thus, the total smoking→WHtR effect (the sum of indirect effect and direct effect) is expected to be negative without adjustment for BMI in the model as shown in [Table 3](#). Apparently, whether or not BMI is adjusted for in the analysis model would lead to different conclusions in this regard. For this reason, the adjustment for BMI has been widely used in the risk factor-central obesity association analysis in the literature [37,38]. In addition, a traditional and powerful adjustment method, stratification analysis, was used to confirm this concept. The regression coefficients of regular smoking for WC and WHtR in the 18 BMI subgroups were all positive, which provided reliable evidence that regular smoking is positively associated with central obesity at all BMI levels. Results of the BMI stratification analyses support the argument that BMI needs to be included in the model for adjustment in the analyses related to smoking-central obesity association, especially for normal-weight adults. The phenomenon that the direction of any relation between two variables is reversed after a third variable is introduced is statistically known as “reversal paradox” [36,39]. The reversal paradox makes it very challenging to correctly interpret the findings seen in observational studies. Although BMI cannot be considered a confounder by definition in this scenario, the reversal paradox experienced in the current study points to the need for the adjustment for BMI in the smoking-central obesity association analysis.

The prevalence of overweight and obesity is higher among women than men, especially in developing countries [40]. Women often report consuming healthier foods, sex hormones (estrogens and testosterone), however, differentially affect adipocyte physiology and body fat distribution [41]. The increases in overweight and obesity in menopausal women are important public health concerns [42]. The prevalence of obesity increases significantly in American women after they reach age 40 [43]. In this study cohort, women showed a significantly higher prevalence of central obesity than men after age 45 in both total sample and the normal-weight group. On the other hand, the positive association between current smoking and central obesity measures was found to be significantly stronger in males than in females in this study. The higher, negative association between smoking and BMI in males ($\beta = -0.093$) than in females ($\beta = -0.025$), stronger correlation between WHtR and BMI ($r = 0.641$) in the male normal-weight group and higher prevalence of smokers in males may account for, in part, the gender-specific association between smoking and central obesity. The current observations suggest that female adults might have other more important mechanisms like sex hormones underlying central obesity in addition to tobacco smoking.

This population-based epidemiologic study has certain limitations. First, subjects voluntarily participated in the study, which might result in selection bias to some extent with respect to the population representativeness despite the large sample size. Second, the pathophysiological mechanisms underlying the association between smoking and central obesity cannot be investigated in the current study because the baseline survey data are observational in nature. Finally, the CKB study is not a nutritional survey, and detailed information on dietary patterns is not available. Therefore, dietary factors, which may be important, were not included in the current analyses.

In conclusion, the current study demonstrates that there are considerable differences in the prevalence of tobacco smokers and central obesity between male and female Chinese adults; smoking is significantly and positively associated with central obesity measured as WC and WHtR, with males having substantially stronger association parameters than females. Importantly, the association patterns of regular smoking with central obesity were largely dependent on BMI adjustment. The present study emphasizes the importance of adjustment for BMI in

the smoking-central obesity association analyses, especially for normal-weight adults. Sophisticated statistical models like structural equation modeling or mediation analysis are needed to investigate this issue in depth. These gender-specific results indicate that other important underlying mechanisms like sex hormones may be involved in the development of central obesity in females in addition to tobacco smoking. The study findings underscore the importance of undertaking preventive strategies for central obesity through smoking behavior control.

Acknowledgments

We thank Judith Mackay in Hong Kong; Yu Wang, Gonghuan Yang, Zhengfu Qiang, Lin Feng, Maigeng Zhou, Wenhua Zhao, and Yan Zhang in China CDC; Lingzhi Kong, Xiucheng Yu, and Kun Li in the Chinese Ministry of Health; and Yiping Chen, Sarah Clark, Martin Radley, Mike Hill, Hongchao Pan, and Jill Boreham in the CTSU, Oxford, for assisting with the design, planning, organization, and conduct of the study. The most important acknowledgement is to the participants in the study and the members of the survey teams in each of the 10 regional centers, as well as to the project development and management teams based at Beijing, Oxford and the 10 regional centers.

Members of China Kadoorie Biobank collaborative group

(a) International Steering Committee. Liming Li, Zhengming Chen, Junshi Chen, Rory Collins, Richard Peto.

(b) Study Coordinating Centers

International Coordinating Center, Oxford: Zhengming Chen, Garry Lancaster, Xiaoming Yang, Alex Williams, Margaret Smith, Ling Yang, Yumei Chang, Iona Millwood, Yiping Chen, Sarah Lewington, Sam Sansome, Robin Walters, Om Kurmi

National Coordinating Center, Beijing: Yu Guo, Zheng Bian, Can Hou, Yunlong Tan, Xin Cai, Huiyan Zhou, Xuguang Chen

Regional Coordinating Centers, 10 areas in China:

Qingdao Qingdao Center for Disease Control: Zengchang Pang, Shanpeng Li, Shaojie Wang. Licang Center for Disease Control: Silu lv.

Heilongjiang Provincial Center for Disease Control: Zhonghou Zhao, Shumei Liu, Zhigang Pang. Nangang Center for Disease Control: Liqiu Yang, Hui He, Bo Yu.

Hainan Provincial Center for Disease Control: Shanqing Wang, Hongmei Wang. Meilan Center for Disease Control: Chunxing Chen, Xiangyang Zheng.

Jiangsu Provincial Center for Disease Control: Xiaoshu Hu, Minghao Zhou, Ming Wu, Ran Tao. Suzhou Center for Disease Control: Yeyuan Wang, Yihe Hu, Liangcai Ma. Wuzhong Center for Disease Control: Renxian Zhou.

Guangxi Provincial Center for Disease Control: Zhenzhu Tang, Naying Chen, Ying Huang. Liuzhou Center for Disease Control: Mingqiang Li, Zhigao Gan, Jinhuai Meng, Jingxin Qin.

Sichuan Provincial Center for Disease Control: Xianping Wu, Ningmei Zhang. Pengzhou Center for Disease Control: Guojin Luo, Xiangsan Que, Xiaofang Chen.

Gansu Provincial Center for Disease Control: Pengfei Ge, Xiaolan Ren, Caixia Dong. Maiji Center for Disease Control: Hui Zhang, Enke Mao, Zhongxiao Li.

Henan Provincial Center for Disease Control: Gang Zhou, Shixian Feng. Huixian Center for Disease Control: Yulian Gao, Tianyou He, Li Jiang, Huarong Sun.

Zhejiang Provincial Center for Disease Control: Min Yu, Danting Su, Feng Lu. Tongxiang Center for Disease Control: Yijian Qian, Kunxiang Shi, Yabin Han, Lingli Chen.

Hunan Provincial Center for Disease Control: Guangchun Li, Huilin Liu, LI Yin. Liuyang Center for Disease Control: Youping Xiong, Zhongwen Tan, Weifang Jia.

Author Contributions

Conceived and designed the experiments: JC ZC LL. Performed the experiments: JC ZC LL. Analyzed the data: JL WC DS. Contributed reagents/materials/analysis tools: YG ZB CY HZ YT. Wrote the paper: JL WC DS SL IYM MS.

References

1. Malik VS, Willett WC, Hu FB. Global obesity: trends, risk factors and policy implications. *Nat Rev Endocrinol*. 2013; 9:13–27. doi: [10.1038/nrendo.2012.199](https://doi.org/10.1038/nrendo.2012.199) PMID: [23165161](https://pubmed.ncbi.nlm.nih.gov/23165161/)
2. Flegal KM, Carroll MD, Kit BK, Ogden CL. Prevalence of obesity and trends in the distribution of body mass index among US adults, 1999–2010. *JAMA*. 2012; 307:491–497. doi: [10.1001/jama.2012.39](https://doi.org/10.1001/jama.2012.39) PMID: [22253363](https://pubmed.ncbi.nlm.nih.gov/22253363/)
3. Ogden CL, Carroll MD, Kit BK, Flegal KM. Prevalence of obesity and trends in body mass index among US children and adolescents, 1999–2010. *JAMA*. 2012; 307:483–490. doi: [10.1001/jama.2012.40](https://doi.org/10.1001/jama.2012.40) PMID: [22253364](https://pubmed.ncbi.nlm.nih.gov/22253364/)
4. Wildman RP, Gu D, Muntner P, Wu X, Reynolds K, Duan X, et al. Trends in overweight and obesity in Chinese adults: between 1991 and 1999–2000. *Obesity (Silver Spring)*. 2008; 16:1448–1453. doi: [10.1038/oby.2008.208](https://doi.org/10.1038/oby.2008.208) PMID: [18388899](https://pubmed.ncbi.nlm.nih.gov/18388899/)
5. Xi B, Liang Y, He T, Reilly KH, Hu Y, Wang Q, et al. Secular trends in the prevalence of general and abdominal obesity among Chinese adults, 1993–2009. *Obes Rev*. 2012; 13:287–296. doi: [10.1111/j.1467-789X.2011.00944.x](https://doi.org/10.1111/j.1467-789X.2011.00944.x) PMID: [22034908](https://pubmed.ncbi.nlm.nih.gov/22034908/)
6. Flegal KM, Kit BK, Orpana H, Graubard BI. Association of all-cause mortality with overweight and obesity using standard body mass index categories: a systematic review and meta-analysis. *JAMA*. 2013; 309:71–82. doi: [10.1001/jama.2012.113905](https://doi.org/10.1001/jama.2012.113905) PMID: [23280227](https://pubmed.ncbi.nlm.nih.gov/23280227/)
7. Zhou B-F. Predictive values of body mass index and waist circumference for risk factors of certain related diseases in Chinese adults—study on optimal cut-off points of body mass index and waist circumference in Chinese adults. *Biomed Environ Sci*. 2002; 15:83–96. PMID: [12046553](https://pubmed.ncbi.nlm.nih.gov/12046553/)
8. Whitlock G, Lewington S, Sherliker P, Clarke R, Emberson J, Halsey J, et al. Body-mass index and cause-specific mortality in 900 000 adults: collaborative analyses of 57 prospective studies. *Lancet*. 2009; 373:1083–1096. doi: [10.1016/S0140-6736\(09\)60318-4](https://doi.org/10.1016/S0140-6736(09)60318-4) PMID: [19299006](https://pubmed.ncbi.nlm.nih.gov/19299006/)
9. Cameron AJ, Boyko EJ, Sicree RA, Zimmet PZ, Söderberg S, Alberti KG, et al. Central obesity as a precursor to the metabolic syndrome in the AusDiab study and Mauritius. *Obesity (Silver Spring)*. 2008; 16:2707–2716. doi: [10.1038/oby.2008.412](https://doi.org/10.1038/oby.2008.412) PMID: [18820650](https://pubmed.ncbi.nlm.nih.gov/18820650/)
10. Srinivasan SR, Wang R, Chen W, Wei CY, Xu J, Berenson GS. Utility of waist-to-height ratio in detecting central obesity and related adverse cardiovascular risk profile among normal weight younger adults (from the Bogalusa Heart Study). *Am J Cardiol*. 2009; 104:721–724. doi: [10.1016/j.amjcard.2009.04.037](https://doi.org/10.1016/j.amjcard.2009.04.037) PMID: [19699351](https://pubmed.ncbi.nlm.nih.gov/19699351/)
11. Hsieh SD, Muto T. Metabolic syndrome in Japanese men and women with special reference to the anthropometric criteria for the assessment of obesity: Proposal to use the waist-to-height ratio. *Prev Med (Baltim)*. 2006; 42:135–139. PMID: [16376978](https://pubmed.ncbi.nlm.nih.gov/16376978/)
12. Hsieh SD, Muto T. The superiority of waist-to-height ratio as an anthropometric index to evaluate clustering of coronary risk factors among non-obese men and women. *Prev Med (Baltim)*. 2005; 40:216–220.
13. Deurenberg P, Deurenberg-Yap M, Guricci S. Asians are different from Caucasians and from each other in their body mass index/body fat per cent relationship. *Obes Rev*. 2002; 3:141–146. PMID: [12164465](https://pubmed.ncbi.nlm.nih.gov/12164465/)
14. Low S, Chin MC, Ma S, Heng D, Deurenberg-Yap M. Rationale for redefining obesity in Asians. *Ann Acad Med Singapore*. 2009; 38:66–69. PMID: [19221673](https://pubmed.ncbi.nlm.nih.gov/19221673/)
15. WHO Expert Consultation. Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *Lancet*. 2004; 363:157–163. PMID: [14726171](https://pubmed.ncbi.nlm.nih.gov/14726171/)
16. Ramachandran A, Chamukuttan S, Shetty SA, Arun N, Susairaj P. Obesity in Asia—is it different from rest of the world. *Diabetes Metab Res Rev*. 2012; 28 Suppl 2:47–51. doi: [10.1002/dmrr.2353](https://doi.org/10.1002/dmrr.2353) PMID: [23280866](https://pubmed.ncbi.nlm.nih.gov/23280866/)
17. Cheng TO. The current state of cardiology in China. *Int J Cardiol*. 2004; 96:425–439. PMID: [15301897](https://pubmed.ncbi.nlm.nih.gov/15301897/)
18. Reynolds K, Gu D, Whelton PK, Wu X, Duan X, Mo J, et al. Prevalence and risk factors of overweight and obesity in China. *Obesity (Silver Spring)*. 2007; 15:10–18.
19. Björntorp P. Visceral obesity: a “civilization syndrome”. *Obes Res*. 1993; 1:206–222. PMID: [16350574](https://pubmed.ncbi.nlm.nih.gov/16350574/)

20. Wadden TA, Webb VL, Moran CH, Bailer BA. Lifestyle modification for obesity: new developments in diet, physical activity, and behavior therapy. *Circulation*. 2012; 125:1157–1170. doi: [10.1161/CIRCULATIONAHA.111.039453](https://doi.org/10.1161/CIRCULATIONAHA.111.039453) PMID: [22392863](https://pubmed.ncbi.nlm.nih.gov/22392863/)
21. Hebden L, Chey T, Allman-Farinelli M. Lifestyle intervention for preventing weight gain in young adults: a systematic review and meta-analysis of RCTs. *Obes Rev*. 2012; 13:692–710. doi: [10.1111/j.1467-789X.2012.00990.x](https://doi.org/10.1111/j.1467-789X.2012.00990.x) PMID: [22413804](https://pubmed.ncbi.nlm.nih.gov/22413804/)
22. Molarius A, Seidell JC, Kuulasmaa K, Dobson AJ, Sans S. Smoking and relative body weight: an international perspective from the WHO MONICA Project. *J Epidemiol Community Health*. 1997; 51:252–260. PMID: [9229053](https://pubmed.ncbi.nlm.nih.gov/9229053/)
23. Audrain-McGovern J, Benowitz NL. Cigarette smoking, nicotine, and body weight. *Clin Pharmacol Ther*. 2011; 90:164–168. doi: [10.1038/clpt.2011.105](https://doi.org/10.1038/clpt.2011.105) PMID: [21633341](https://pubmed.ncbi.nlm.nih.gov/21633341/)
24. Chioloro A, Faeh D, Paccaud F, Cornuz J. Consequences of smoking for body weight, body fat distribution, and insulin resistance. *Am J Clin Nutr*. 2008; 87:801–809. PMID: [18400700](https://pubmed.ncbi.nlm.nih.gov/18400700/)
25. Li L, Lv J, Guo Y, Collins R, Chen JS, Peto R, et al. The China Kadoorie Biobank: related methodology and baseline characteristics of the participants. *Zhonghua Liu Xing Bing Xue Za Zhi*. 2012; 33:249–255. PMID: [22613372](https://pubmed.ncbi.nlm.nih.gov/22613372/)
26. Chen Z, Chen J, Collins R, Guo Y, Peto R, Wu F, et al. China Kadoorie Biobank of 0.5 million people: survey methods, baseline characteristics and long-term follow-up. *Int J Epidemiol*. 2011; 40:1652–1666. doi: [10.1093/ije/dyr120](https://doi.org/10.1093/ije/dyr120) PMID: [22158673](https://pubmed.ncbi.nlm.nih.gov/22158673/)
27. Du H, Bennett D, Li L, Whitlock G, Guo Y, Collins R, et al. Physical activity and sedentary leisure time and their associations with BMI, waist circumference, and percentage body fat in 0.5 million adults: the China Kadoorie Biobank study. *Am J Clin Nutr*. 2013; 97:487–496. doi: [10.3945/ajcn.112.046854](https://doi.org/10.3945/ajcn.112.046854) PMID: [23364014](https://pubmed.ncbi.nlm.nih.gov/23364014/)
28. Misra A, Shrivastava U. Obesity and dyslipidemia in South Asians. *Nutrients*. 2013; 5:2708–2733. doi: [10.3390/nu5072708](https://doi.org/10.3390/nu5072708) PMID: [23863826](https://pubmed.ncbi.nlm.nih.gov/23863826/)
29. Williamson DF, Madans J, Anda RF, Kleinman JC, Giovino GA, Byers T. Smoking cessation and severity of weight gain in a national cohort. *N Engl J Med*. 1991; 324:739–745. PMID: [1997840](https://pubmed.ncbi.nlm.nih.gov/1997840/)
30. Benowitz NL. Nicotine addiction. *N Engl J Med*. 2010; 362:2295–2303. doi: [10.1056/NEJMra0809890](https://doi.org/10.1056/NEJMra0809890) PMID: [20554984](https://pubmed.ncbi.nlm.nih.gov/20554984/)
31. Wakabayashi I. Relationship between smoking and metabolic syndrome in men with diabetes mellitus. *Metab Syndr Relat Disord*. 2014; 12:70–78. doi: [10.1089/met.2013.0110](https://doi.org/10.1089/met.2013.0110) PMID: [24266721](https://pubmed.ncbi.nlm.nih.gov/24266721/)
32. Kim JH, Shim KW, Yoon YS, Lee SY, Kim SS, Oh SW. Cigarette smoking increases abdominal and visceral obesity but not overall fatness: an observational study. *PLoS One*. 2012; 7:e45815. doi: [10.1371/journal.pone.0045815](https://doi.org/10.1371/journal.pone.0045815) PMID: [23029258](https://pubmed.ncbi.nlm.nih.gov/23029258/)
33. Shimokata H, Muller DC, Andres R. Studies in the distribution of body fat. III. Effects of cigarette smoking. *JAMA*. 1989; 261:1169–1173. PMID: [2915440](https://pubmed.ncbi.nlm.nih.gov/2915440/)
34. Canoy D, Wareham N, Luben R, Welch A, Bingham S, Day N, et al. Cigarette smoking and fat distribution in 21,828 British men and women: a population-based study. *Obes Res*. 2005; 13:1466–1475. PMID: [16129730](https://pubmed.ncbi.nlm.nih.gov/16129730/)
35. Jee SH, Lee SY, Nam CM, Kim SY, Kim MT. Effect of smoking on the paradox of high waist-to-hip ratio and low body mass index. *Obes Res*. 2002; 10:891–895. PMID: [12226137](https://pubmed.ncbi.nlm.nih.gov/12226137/)
36. Hernández-Díaz S, Schisterman EF, Hernán MA. The birth weight "paradox" uncovered? *Am J Epidemiol*. 2006; 164:1115–1120. PMID: [16931543](https://pubmed.ncbi.nlm.nih.gov/16931543/)
37. Heid IM, Jackson AU, Randall JC, Winkler TW, Qi L, Steinthorsdottir V, et al. Meta-analysis identifies 13 new loci associated with waist-hip ratio and reveals sexual dimorphism in the genetic basis of fat distribution. *Nat Genet*. 2010; 42:949–960. doi: [10.1038/ng.685](https://doi.org/10.1038/ng.685) PMID: [20935629](https://pubmed.ncbi.nlm.nih.gov/20935629/)
38. Liu CT, Monda KL, Taylor KC, Lange L, Demerath EW, Palmas W, et al. Genome-wide association of body fat distribution in African ancestry populations suggests new loci. *PLoS Genet*. 2013; 9(8): e1003681. doi: [10.1371/journal.pgen.1003681](https://doi.org/10.1371/journal.pgen.1003681) PMID: [23966867](https://pubmed.ncbi.nlm.nih.gov/23966867/)
39. Tu Y-K, West R, Ellison GTH, Gilthorpe MS. Why evidence for the fetal origins of adult disease might be a statistical artifact: the "reversal paradox" for the relation between birth weight and blood pressure in later life. *Am J Epidemiol*. 2005; 161:27–32. PMID: [15615910](https://pubmed.ncbi.nlm.nih.gov/15615910/)
40. Kanter R, Caballero B. Global gender disparities in obesity: a review. *Adv Nutr* 2012; 3:491–498. doi: [10.3945/an.112.002063](https://doi.org/10.3945/an.112.002063) PMID: [22797984](https://pubmed.ncbi.nlm.nih.gov/22797984/)
41. Lizcano F, Guzmán G. Estrogen Deficiency and the Origin of Obesity during Menopause. *Biomed Res Int*. 2014; 2014:757461. doi: [10.1155/2014/757461](https://doi.org/10.1155/2014/757461) PMID: [24734243](https://pubmed.ncbi.nlm.nih.gov/24734243/)

42. Wietlisbach V, Marques-Vidal P, Kuulasmaa K, Karvanen J, Paccaud F. The relation of body mass index and abdominal adiposity with dyslipidemia in 27 general populations of the WHO MONICA Project. *Nutrition, Metabolism and Cardiovascular Diseases*. 2013; 2:432–442.
43. Flegal KM, Carroll MD, Ogden CL, Curtin LR. Prevalence and trends in obesity among US adults, 1999–2008. *JAMA*. 2010; 303:235–241. doi: [10.1001/jama.2009.2014](https://doi.org/10.1001/jama.2009.2014) PMID: [20071471](https://pubmed.ncbi.nlm.nih.gov/20071471/)