

# RNA $N^6$ -methyladenosine methylation in post-transcriptional gene expression regulation

Yanan Yue,<sup>1,2</sup> Jianzhao Liu,<sup>1,2</sup> and Chuan He<sup>1,2</sup>

<sup>1</sup>Department of Chemistry, Institute for Biophysical Dynamics, The University of Chicago, Chicago, Illinois 60637, USA; <sup>2</sup>Howard Hughes Medical Institute, The University of Chicago, Chicago, Illinois 60637, USA

**$N^6$ -methyladenosine ( $m^6A$ ) is the most prevalent and internal modification that occurs in the messenger RNAs (mRNA) of most eukaryotes, although its functional relevance remained a mystery for decades. This modification is installed by the  $m^6A$  methylation “writers” and can be reversed by demethylases that serve as “erasers.” In this review, we mainly summarize recent progress in the study of the  $m^6A$  mRNA methylation machineries across eukaryotes and discuss their newly uncovered biological functions. The broad roles of  $m^6A$  in regulating cell fates and embryonic development highlight the existence of another layer of epigenetic regulation at the RNA level, where mRNA is subjected to chemical modifications that affect protein expression.**

Both DNA and histone proteins undergo dynamic and reversible chemical modifications to control gene expression (Strahl and Allis 2000; Bird 2001; Suzuki and Bird 2008; Bhutani et al. 2011; Jones 2012; Kohli and Zhang 2013). Although post-transcriptional modifications are known to occur to RNAs, the impact of these modifications on gene expression regulation has only recently begun to be explored (He 2010). To date, more than a hundred structurally distinct chemical modifications have been found in eukaryotic RNAs (Cantara et al. 2011; Machnicka et al. 2013); however, the enzymes responsible for each modification and the biological consequences of these modified RNAs are largely unknown. RNA modifications were once considered to be static, but a flurry of recent discoveries has demonstrated that some chemical modifications can be dynamic and participate in the regulation of diverse physiological processes (Motorin and Helm 2011; Yi and Pan 2011; Chan et al. 2012; Fu et al. 2014; Meyer and Jaffrey 2014; Kirchner and Ignatova 2015). The presence of  $N^6$ -methyladenosine ( $m^6A$ ) in pol-

yadenylated mRNA was first discovered in the 1970s (Desrosiers et al. 1974; Perry and Kelley 1974; Lavi and Shatkin 1975; Wei et al. 1975; Schibler et al. 1977; Wei and Moss 1977) by researchers who were characterizing the 5' cap structure of messenger RNA (mRNA) in mammalian cells. Since then,  $m^6A$  has been identified as the most prevalent internal modification in mRNA and long noncoding RNA (lncRNA) in higher eukaryotes. It is widely conserved among eukaryotic species that range from yeast, plants, and flies to mammals as well as among viral mRNAs that replicate inside host nuclei (Krug et al. 1976; Beemon and Keith 1977; Horowitz et al. 1984; Bokar 2005). In addition to its occurrence in mRNA,  $m^6A$  also exists in various classes of RNA in eukaryotes, bacteria, and archaea, including ribosomal RNAs, small nuclear RNAs, and transfer RNAs (Bjork et al. 1987; Maden 1990; Shimba et al. 1995; Gu et al. 1996; Agris et al. 2007; Piekna-Przybylska et al. 2008). Despite its widespread distribution in the mammalian transcriptome (on average, approximately three  $m^6A$  sites per mRNA), functional insight has been lacking, possibly due to the low abundance of  $m^6A$  mRNA and technical difficulties in global detection.

Interest in the biological relevance of  $m^6A$  in mRNA resurfaced after the discovery of two mammalian RNA demethylases, FTO (fat mass and obesity-associated protein) (Jia et al. 2011) and its homolog, ALKBH5 (Zheng et al. 2013), which selectively reverse  $m^6A$  to adenosine in nuclear RNA. FTO is associated with human obesity (Dina et al. 2007; Frayling et al. 2007; Loos and Yeo 2014) and mental development (Hess et al. 2013), while ALKBH5 is shown to affect mouse spermatogenesis in a demethylation-dependent manner (Zheng et al. 2013), suggesting broad roles of  $m^6A$  in various physiological processes. Shortly after these findings, YTHDF2 (YTH domain-containing family protein 2) was identified as the first  $m^6A$  reader protein that preferentially recognizes

[**Keywords:**  $N^6$ -methyladenosine,  $m^6A$  methyltransferase, RNA demethylase, METTL3–METTL14, mRNA methylation, post-transcriptional regulation]

**Corresponding author:** [chuanhe@uchicago.edu](mailto:chuanhe@uchicago.edu)

Article is online at <http://www.genesdev.org/cgi/doi/10.1101/gad.262766.115>.

© 2015 Yue et al. This article is distributed exclusively by Cold Spring Harbor Laboratory Press for the first six months after the full-issue publication date (see <http://genesdev.cshlp.org/site/misc/terms.xhtml>). After six months, it is available under a Creative Commons License (Attribution-NonCommercial 4.0 International), as described at <http://creativecommons.org/licenses/by-nc/4.0/>.


m<sup>6</sup>A-containing mRNA (Dominissini et al. 2012; Wang et al. 2014a) and mediates mRNA decay (Wang et al. 2014a), thereby suggesting a role for m<sup>6</sup>A RNA as a negative regulator of gene expression. On the other hand, a transcriptome-wide m<sup>6</sup>A profiling method was developed to decipher the m<sup>6</sup>A RNA landscape (Dominissini et al. 2012; Meyer et al. 2012). Intriguingly, m<sup>6</sup>A sites in mammalian polyadenylated RNA are dominated by the conserved Pu[G > A]m<sup>6</sup>AC[A/C/U] motif that localizes near stop codons, in 3' untranslated regions (UTRs), within long internal exons, and at 5' UTRs (Dominissini et al. 2012; Meyer et al. 2012; Schwartz et al. 2013; Li et al. 2014; Luo et al. 2014), immediately raising the question of how this specificity is achieved. The m<sup>6</sup>A RNA landscape is initially sculptured by a methyltransferase complex, but for a long time, METTL3 (methyltransferase-like 3) was the only known SAM (S-adenosyl methionine)-binding subunit associated with mRNA methylation (Bokar et al. 1997). In 2014, a new mammalian methyltransferase, METTL14, was discovered to catalyze m<sup>6</sup>A methylation. Together with METTL3, these two proteins form a stable heterodimer complex that mediates cellular m<sup>6</sup>A deposition on mammalian mRNAs (Liu et al. 2014; Wang et al. 2014b). Recently, the mammalian splicing factor WTAP (Wilms' tumor 1-associating protein) was identified as the third auxiliary factor of the core methyltransferase complex that affects cellular m<sup>6</sup>A methylation (Liu et al. 2014; Ping et al. 2014). The identification and characterization of the complete mammalian m<sup>6</sup>A methylation machinery are the first steps toward deciphering the selectivity and biological functions of m<sup>6</sup>A deposition in eukaryotic mRNAs.

In this review, we mainly summarize recent progress in the study of m<sup>6</sup>A methylation in mRNA across different eukaryotes and discuss their newly discovered roles in post-transcriptional gene expression regulation. We first

describe the features of m<sup>6</sup>A on a global scale and briefly introduce the mammalian m<sup>6</sup>A writers, erasers, and readers that specifically install, remove, or bind to m<sup>6</sup>A at defined sequence motifs (Fig. 1). We then discuss the evolutionary conservation of the m<sup>6</sup>A methylation machinery across eukaryotic species that range from yeast, plants, and flies to mammals, highlighting the broad roles of methyltransferases and m<sup>6</sup>A in regulating cell status and embryonic development. Finally, we discuss the emerging functions of m<sup>6</sup>A in several mechanisms of post-transcriptional gene expression regulation with a special focus on the effects of m<sup>6</sup>A on differentiation and reprogramming of stem cells.

### Features of m<sup>6</sup>A on a global scale

Studies in the 1970s revealed that m<sup>6</sup>A modification in mRNA mainly occurs at Pu[G > A]m<sup>6</sup>AC[U > A > C] (Pu represents purine) and is estimated to be present at an average level of approximately three m<sup>6</sup>A residues per mRNA (Rottman et al. 1974; Narayan and Rottman 1988; Csepány et al. 1990; Narayan et al. 1994). Transcriptome-wide mapping of m<sup>6</sup>A is hindered by the following two facts: (1) m<sup>6</sup>A, akin to A, reverse-transcribes to a thymine (T), and (2) m<sup>6</sup>A is not susceptible to chemical modifications that might promote its detection. In 2012, two groups independently developed an antibody-based high-throughput sequencing method (Dominissini et al. 2012; Meyer et al. 2012) and for the first time profiled the transcriptome-wide m<sup>6</sup>A distribution. In each method, mammalian mRNA is properly fragmented and immunoprecipitated by an m<sup>6</sup>A-specific antibody. Libraries are prepared from immunoprecipitated and input control fragments, respectively, and subjected to high-throughput sequencing. In general, ~12,000 m<sup>6</sup>A sites in the transcripts



**Figure 1.** Illustration of the cellular pathways of m<sup>6</sup>A in nuclear RNAs. The m<sup>6</sup>A methyltransferases and demethylases dynamically control the m<sup>6</sup>A methylation landscape within the nucleus. The m<sup>6</sup>A reader proteins preferentially bind to the methylated RNA and mediate specific functions. In the nucleus, m<sup>6</sup>A may affect alternative splicing of pre-mRNA and mature mRNA storage and export. After mature RNAs are exported to the cytoplasm, cytoplasmic m<sup>6</sup>A reader YTHDF2 can bind to the m<sup>6</sup>A-containing mRNAs to mediate mRNA decay. Other cytoplasmic readers could modulate mRNA translation and storage.

of ~7000 coding genes and ~300 noncoding ones are identified in human cells. The resolution of the m<sup>6</sup>A peak site is ~100 nucleotides (nt), which was further improved by later optimization (Schwartz et al. 2013, 2014b; Chen et al. 2015a). However, transcriptome-wide m<sup>6</sup>A detection at single-base resolution remains a challenge.


To date, m<sup>6</sup>A RNA methylomes across many eukaryotes, including human (Dominissini et al. 2012; Meyer et al. 2012; Batista et al. 2014; Schwartz et al. 2014b), mouse (Dominissini et al. 2012; Meyer et al. 2012; Batista et al. 2014; Schwartz et al. 2014b; Wang et al. 2014b; Geula et al. 2015), yeast (Schwartz et al. 2013), and plant (Li et al. 2014; Luo et al. 2014), have been profiled. In general, global mapping reveals a conserved, widespread, and dynamic mRNA methylation in eukaryotes. Three salient features of the m<sup>6</sup>A methylome are evident. First, m<sup>6</sup>A sites are mainly confined to the consensus motif Pu[G > A] m<sup>6</sup>AC[U > A > C], which is consistent with earlier studies. Second, m<sup>6</sup>A marks are not equally distributed across the transcriptome; they are preferentially enriched in a subset of consensus sequences near stop codons, in 3' UTRs, and within long internal exons (Fig. 2). In particular, this topology is preserved upon endodermal differentiation of stem cells (Batista et al. 2014; Geula et al. 2015). Last, m<sup>6</sup>A-modified genes are well conserved between human and mouse embryonic stem cells (ESCs) and somatic cells (Batista et al. 2014). For instance, ~70% of human ESC genes are also m<sup>6</sup>A-modified in the orthologous mouse gene, with ~46% of the m<sup>6</sup>A peak sites in common. As expected, higher m<sup>6</sup>A peak intensities were detected in conserved sites compared with those that are not conserved. On the other hand, distinct m<sup>6</sup>A patterns can also be detected among different species or cells residing in different developmental stages (Meyer

et al. 2012; Schwartz et al. 2013; Batista et al. 2014; Geula et al. 2015). Certain m<sup>6</sup>A modifications are tissue-specific and dynamically alter in response to different stimuli, indicating the potential role of m<sup>6</sup>A in regulating diverse cellular processes.


### m<sup>6</sup>A writers in mammals

The m<sup>6</sup>A modification is installed by a multicomponent methyltransferase complex (Fig. 1), which has not been fully characterized. In a pioneer work reported in 1997 (Bokar et al. 1997), a 200-kDa methyltransferase complex was isolated from the HeLa nuclear extract, which exhibits methyltransferase activity. Only a 70-kDa protein, termed MT-A70 or METTL3, was identified as one SAM-binding unit within this 200-kDa methyltransferase complex. The knockdown of METTL3 led to apoptosis of human HeLa cells and a concomitant reduction in cellular m<sup>6</sup>A level (Bokar 2005). METTL3 and m<sup>6</sup>A appear to be strongly associated with development and gametogenesis, since the depletion of the METTL3 homologs in yeast (Agarwala et al. 2012), flies (Hongay and Orr-Weaver 2011), and plants (Zhong et al. 2008; Bodi et al. 2012) readily lead to developmental arrest or defects in gametogenesis.

A phylogenetic analysis of the MT-A70 (METTL3) family methyltransferase has suggested METTL14, which shares 43% identity with METTL3 but belongs to a different lineage, as a homolog of METTL3 (Fig. 3; Bujnicki et al. 2002). The highly conserved nature of METTL14 in mammals together with the fact that the METTL14 protein can be pulled down by METTL3 has prompted researchers to consider METTL14 as a putative candidate for m<sup>6</sup>A deposition on mRNA (Liu et al. 2014). Intriguingly, knockdown of METTL14 results in a more pronounced decrease of m<sup>6</sup>A in polyadenylated RNA compared with knockdown of METTL3 in both HeLa and human 293 FT cell lines (Liu et al. 2014). The recombinant METTL3 and METTL14 proteins can form a stable METTL3–METTL14 complex in the gel filtration experiment, and subsequent two-dimensional native/SDS-PAGE analysis has further demonstrated the formation of a heterodimer between these two proteins, with a stoichiometry of 1/1 (Liu et al. 2014). While the METTL14 protein itself exhibits higher methylation activity compared with METTL3 in vitro, the combination of both methyltransferases substantially enhances methylation efficiency, demonstrating a synergistic effect that is further confirmed by in vivo studies (Liu et al. 2014; Wang et al. 2014b). The METTL3–METTL14 heterodimer preferentially methylates RNA substrates containing the previously identified consensus sequence GGACU and exhibits a modest preference for the less structured RNA substrate in vitro. Furthermore, the methyltransferase complex was isolated from the native HeLa cell nuclear extract. The nuclear extract fraction that exhibits the highest methylation activity was found to be mostly enriched with METTL3 and METTL14 (Liu et al. 2014), thus clearly indicating that the heterodimer of METTL3–METTL14 forms the catalytic core of the mammalian m<sup>6</sup>A methyltransferase complex.



**Figure 2.** The normalized distribution (density) of m<sup>6</sup>A peaks along the mRNA transcripts in HeLa cells (*top* panel) and *Arabidopsis thaliana* (*bottom* panel), where each mRNA transcript is divided into the 5' UTR, coding sequences (CDS), and the 3' UTR.



**Figure 3.** Simplified phylogenetic analysis of the MT-A70 (METTL3) superfamily. Each subfamily is marked with different colors; its corresponding conserved signature motif at the catalytic site is listed for comparison.

WTAP has been identified as the third crucial component of the mammalian m<sup>6</sup>A methyltransferase complex (Fig. 1; Liu et al. 2014; Ping et al. 2014). WTAP was initially shown to act as a splicing factor that binds to Wilms' tumor 1 protein (Little et al. 2000) and plays a regulatory role in cell cycle progression and early embryo development (Horiuchi et al. 2006, 2013). The first evidence of WTAP as a third component of the methyltransferase complex came from the coimmunoprecipitation result, which showed that WTAP readily binds to the METTL3–METTL14 heterodimer inside cells, although the interactions between WTAP and the two methyltransferases are weaker compared with that between METTL3 and METTL14 (Liu et al. 2014). WTAP itself does not possess methylation activity, consistent with its lack of a conserved catalytic methylation domain, but interacts with the METTL3–METTL14 heterodimer to substantially affect cellular m<sup>6</sup>A deposition (Liu et al. 2014; Schwartz et al. 2014b). A subsequent study suggests that WTAP helps to coordinate the localization of the METTL3–METTL14 heterodimer into nuclear speckles, thereby facilitating m<sup>6</sup>A deposition (Ping et al. 2014).

Global analysis indicates that METTL3, METTL14, and WTAP share a large portion of common binding sites (~36%) on their RNA substrates and exhibit a binding consensus motif similar, if not identical, to that of m<sup>6</sup>A (Liu et al. 2014). A PAR-CLIP (photoactivatable ribonucle-

oside-enhanced cross-linking and immunoprecipitation) assay revealed that a large fraction of the binding sites fall into intergenic regions (~46%) and introns (~31%). This observation suggests that the core methyltransferase complex might work on precursor mRNAs (pre-mRNAs); however, whether and how m<sup>6</sup>A is installed is not yet known (Fig. 1). The m<sup>6</sup>A mark may play a regulatory role in alternative splicing pathways because alternative splicing can be directly affected by the presence of the m<sup>6</sup>A modification in the spliced region (Fig. 1; Geula et al. 2015). In addition, silencing of the methyltransferase complex leads to enhanced abundance of their m<sup>6</sup>A target transcripts, supporting the role of m<sup>6</sup>A as a negative regulator of gene expression (Batista et al. 2014; Liu et al. 2014; Schwartz et al. 2014b; Wang et al. 2014a,b; Geula et al. 2015).

The discovery of the core mammalian m<sup>6</sup>A methyltransferase complex comprised of METTL3, METTL14, and WTAP reveals several new insights. It is surprising and interesting that the core complex of the mRNA m<sup>6</sup>A methyltransferase contains two parallel active methyltransferases. Each is active and seems to impact different sets of transcripts. One potential explanation points to the selective regulation of different pathways and functions inside cells. Each component may be subjected to different post-translational modifications or binding of partner proteins for the tuning of specific



pathways through m<sup>6</sup>A methylation. Meanwhile, how WTAP, a splicing factor with a noticeable mouse phenotype (Horiuchi et al. 2006), participates in and facilitates m<sup>6</sup>A methylation remains to be unveiled. Intriguingly, WTAP orthologs in yeast and plants are also shown to interact with the corresponding METTL3 orthologs (Zhong et al. 2008; Agarwala et al. 2012), while its presence in yeast is directly associated with m<sup>6</sup>A methylation activity (Agarwala et al. 2012). In principle, WTAP could recruit additional auxiliary proteins or RNAs to coordinate methylation of selective RNA substrates. Careful identification of its binding proteins or RNAs may provide a hint in the future. Recent work identified several mammalian WTAP-interacting protein candidates, many of which reside in the RNA processing machinery and have reported roles in alternative splicing (Horiuchi et al. 2013). Whether and how WTAP regulates alternative splicing in an m<sup>6</sup>A-dependent manner have yet to be systematically explored.

### m<sup>6</sup>A erasers in mammals

FTO is the first identified demethylase that oxidatively reverses m<sup>6</sup>A to adenosine in mRNA (Jia et al. 2011). FTO is a member of the AlkB subfamily of Fe<sup>II</sup>/ $\alpha$ -ketoglutarate-dependent dioxygenases, which has eight other family members in humans (ALKBH1–ALKBH8) and catalyzes the oxidation of diverse biological substrates (Kurowski et al. 2003; Gerken et al. 2007; Fu et al. 2010; Zheng et al. 2014). FTO was initially thought to work on 3-methylthymidine (3mT) in ssDNA (Gerken et al. 2007) and 3-methyluracil (3mU) in ssRNA (Jia et al. 2008). In 2011, FTO was discovered to efficiently demethylate m<sup>6</sup>A in nuclear RNA (Jia et al. 2011). A subsequent study showed that FTO can oxidize m<sup>6</sup>A to two previously unknown intermediates—N<sup>6</sup>-hydroxymethyladenosine (hm<sup>6</sup>A) and N<sup>6</sup>-formyladenosine (f<sup>6</sup>A)—in a stepwise manner (Fu et al. 2013). Intriguingly, this process is similar to the oxidation of 5-methylcytosine (5mC) in genomic DNA to 5-hydroxymethylcytosine (5hmC) and then 5-formylcytosine (5fC) by the TET (ten eleven translocation) family proteins (Tahiliani et al. 2009; Ito et al. 2010, 2011), which also belong to the general family of Fe<sup>II</sup>/ $\alpha$ -ketoglutarate-dependent dioxygenases. TET proteins can further oxidize 5fC to 5-carboxylcytosine (5caC) (He et al. 2011; Ito et al. 2011; Zhang et al. 2012). While 5hmC, 5fC, and 5caC are stable cytosine derivatives, hm<sup>6</sup>A and f<sup>6</sup>A are short-lived intermediates with half-lives of ~3 h in aqueous solution under physiological conditions (Fu et al. 2013). The continuous oxidation of 5hmC by the TET family proteins is a critical step in the active DNA demethylation pathway in mammals (He et al. 2011; Pastor et al. 2013; Shen et al. 2014). It is not yet clear whether hm<sup>6</sup>A and f<sup>6</sup>A have specific biological functions.

Immunostaining revealed that the FTO protein mainly resides in the nucleus and partially colocalizes with nuclear speckles (Jia et al. 2011), suggesting a dynamic model of m<sup>6</sup>A demethylation on mRNA coupled with m<sup>6</sup>A deposition and RNA processing. A recent study found

that FTO can modulate alternative splicing of the important adipogenic factor RUNX1T1 by removing the m<sup>6</sup>A residues around the splice sites (Zhao et al. 2014). It is proposed that loss of m<sup>6</sup>A on *RUNX1T1* transcripts prevents the binding of the splicing factor SRSF2 protein and promotes the production of a shorter isoform, which in turn acts to induce preadipocyte differentiation. FTO is also found in the cytoplasm in several cell types, suggesting a possible role of FTO in modulating cytosolic mRNA processing (Gulati et al. 2013; Vujovic et al. 2013).

Shortly after the discovery of FTO, ALKBH5 was identified and characterized as a second mammalian m<sup>6</sup>A demethylase that displays distinct biological functions (Zheng et al. 2013). Like FTO, ALKBH5 preferentially binds ssRNAs due to the presence of a unique loop in ALKBH5 that confers single-stranded substrate selectivity (Aik et al. 2014; Xu et al. 2014a). Distinct from FTO, though, ALKBH5 directly reverses m<sup>6</sup>A to adenosine with no detected intermediates. ALKBH5 is primarily colocalized with nuclear speckles and affects mRNA export and RNA metabolism in a demethylation-dependent manner (Zheng et al. 2013). ALKBH5 knockout mice exhibit impaired male fertility, consistent with the highest expression level of ALKBH5 being in the testis (Zheng et al. 2013). In contrast, FTO is most highly expressed in mouse brains, and FTO-deficient mice mainly suffer from early mortality and reduced body mass (Gerken et al. 2007; Fischer et al. 2009). Taken together, the diverse functions regulated by these two demethylases suggest broad physiological roles of m<sup>6</sup>A.

Further research is needed to delineate the mechanisms by which demethylases act on specific mRNAs and lncRNAs. Advanced sequencing methods coupled with global analysis approaches will help to define the demethylomes of FTO and ALKBH5.

### m<sup>6</sup>A readers in mammals

While the transcriptome-wide RNA m<sup>6</sup>A landscape is sculpted by methyltransferases and demethylases in a dynamic and reversible manner, proteins that preferentially recognize m<sup>6</sup>A (termed m<sup>6</sup>A readers) bind to methylated RNA and confer downstream functions. Studies using methylated RNA probes to pull down binding proteins followed by mass spectrometry identification have identified several m<sup>6</sup>A reader candidates in mammalian cells (Dominissini et al. 2012). Among them, the YTH domain-containing family proteins (YTHDF1–3) were validated as m<sup>6</sup>A readers in cytoplasm, with binding affinities to methylated RNA ranging from ~180 nM to ~520 nM (Wang et al. 2014a). Subsequently, YTHDC1, another member of the YTH domain family, was identified as a mammalian m<sup>6</sup>A reader in the nucleus (Xu et al. 2014b). Mrb1 (methylated RNA-binding 1), a yeast protein with an YTH domain, was also shown to be an m<sup>6</sup>A reader (Schwartz et al. 2013). Crystal structure characterizations of the YTH domain containing a bound m<sup>6</sup>A further reveal a conserved hydrophobic pocket used for the binding of the methyl group of m<sup>6</sup>A as well as the preferential

recognition of the GG(m<sup>6</sup>A)C motif by certain reader proteins (Xu et al. 2014b).

The binding sites and physiological targets of these m<sup>6</sup>A reader proteins can be readily profiled using transcriptome-wide methods, such as PAR-CLIP. In fact, changing the cellular level of the specific reader proteins could give functional insight into the roles of the reader proteins as well as the fate of the corresponding substrate mRNA. YTHDF2 was shown to mediate mRNA decay (Fig. 1) by selectively binding to its transcript targets at a defined G(m<sup>6</sup>A)C consequence motif (Wang et al. 2014a). YTHDF2 binds to m<sup>6</sup>A via its C-terminal YTH domain and facilitates the relocalization of the cognate mRNA from the actively translating pool to mRNA decay sites through its N-terminal domain. However, biological functions of YTHDF1, YTHDF3, and YTHDC1 remain to be unveiled. A recent study showed that YTHDF1 promotes translation of m<sup>6</sup>A-containing transcripts (Wang et al. 2015), presenting a novel mechanism of translation promotion by m<sup>6</sup>A in mRNA.

A recent study also suggests heterogeneous nuclear ribonucleoproteins (hnRNPs) as potential “indirect” nuclear m<sup>6</sup>A readers. When m<sup>6</sup>A is installed in a stem-loop of RNA, it can alter the local RNA structure by destabilizing the base-pairing between the m<sup>6</sup>A consensus motif and the uridine track and thus facilitate the binding of HNRNPC to the uridine track in the loop (Liu et al. 2015). Depletion of m<sup>6</sup>A impairs the binding of HNRNPC and thereby affects the abundance and alternative splicing of its target RNAs. This study reveals another function of m<sup>6</sup>A; namely, by altering the structure of RNA (termed m<sup>6</sup>A switch), m<sup>6</sup>A facilitates the binding of a regulatory protein and thereby modulates gene expression and RNA maturation. Indeed, structural mapping of mRNA inside mammalian cells has revealed that the methylation regions of mRNA tend to lack secondary structures, highlighting the potential role of m<sup>6</sup>A in shaping RNA structures (Schwartz et al. 2013; Wan et al. 2014; Spitale et al. 2015).

### Conservation of m<sup>6</sup>A RNA methylation machinery and its related biological functions across eukaryotes

The identification and characterization of the m<sup>6</sup>A methylation machineries are the first steps toward elucidating the biological roles of m<sup>6</sup>A in mRNAs. Phylogenetic analysis revealed that the MT-A70 (METTL3) superfamily consists of four lineages of proteins with varied degrees of interrelatedness (Bujnicki et al. 2002). The simplified and updated version is shown in Figure 3. Lineages A, B, and C are unique to eukaryotes, while lineage D corresponds to a small group of bacterial DNA m<sup>6</sup>A methyltransferases associated with restriction/modification systems. Among eukaryotes, humans, mice, pufferfish, *Drosophila melanogaster*, and *Arabidopsis thaliana* each contain representatives of the A, B, and C lineages. For instance, humans have representative proteins METTL3, METTL14, and METTL4 that belong to the A–C subfamily, respectively. The budding yeast *Saccharomyces cerevisiae* specifies IME4 (inducer of meiosis 4)

and KAR4 (karyogamy protein) in the A and B lineages, respectively, while the fission yeast *Schizosaccharomyces pombe* seems to have only one member in lineage C. Conservation of the methylation signature motifs such as DPPW and EPPL (Fig. 3) in the MT-A70 superfamily members suggests a common ancestry. Genetic studies of methyltransferases in different organisms have been performed in order to understand functional roles of m<sup>6</sup>A methylation on mRNA (Table 1). Below we focus on reviewing methyltransferases in different organisms and their associated biological functions.

#### *m<sup>6</sup>A methylation machinery in yeast: the MIS [MUM2 (muddled meiosis 2)–IME4–SLZ1 (sporulation-specific leucine zipper 1)] complex mediates m<sup>6</sup>A RNA deposition during yeast meiosis*

Unlike mammals, m<sup>6</sup>A methylation in yeast *S. cerevisiae* is confined to meiosis; m<sup>6</sup>A starts to accumulate on mRNA at the onset of meiosis, peaks in premeiotic S and G2/prophase, and decreases as strains enter into the meiotic divisions. In fact, the modification is hardly detected in yeast undergoing mitotic growth (Clancy et al. 2002; Bodi et al. 2010; Agarwala et al. 2012). High-resolution mapping of m<sup>6</sup>A sites in meiotic yeast transcripts reveals that the methylation sites are primarily enriched in a consensus motif—RGAC (R = A/G), similar to the consensus motif in mammals—and are strongly biased toward the 3' end of the transcripts (Schwartz et al. 2013). IME4 (yeast homolog of mammalian METTL3) is identified as an essential component for m<sup>6</sup>A deposition on yeast mRNA and regulates meiotic progression via RNA methylation. Depletion of IME4 in yeast is not lethal but delays cellular entry into meiosis divisions and hinders sporulation (Shah and Clancy 1992; Hongay et al. 2006; Agarwala et al. 2012). A two-hybrid screen in yeast has identified a core m<sup>6</sup>A RNA methyltransferase complex (termed MIS) composed of IME4, MUM2 (yeast homolog of mammalian WTAP), and a third crucial component, SLZ1 (not conserved in mammals) (Table 1; Agarwala et al. 2012). Intriguingly, each component of the MIS complex is expressed in a meiosis-specific manner, consistent with meiosis-confined methylation (Agarwala et al. 2012; Schwartz et al. 2013). At the onset of meiosis, SLZ1 expression is transcriptionally activated by IME1, a master regulator of yeast meiosis (Schwartz et al. 2013). Upon the induction of meiosis, SLZ1 shuttles IME4 and MUM2 from the cytoplasm into the nucleolus. Notably, nucleolar entry of the MIS complex is essential for m<sup>6</sup>A deposition on yeast mRNA, and the global m<sup>6</sup>A level subsequently reaches its maximum at meiotic prophase. After that, down-regulation of m<sup>6</sup>A deposition is induced by activation of NDT80, a transcription factor required for exit from meiotic G2/prophase (Chu and Herskowitz 1998). As a result, the MIS complex exits from the nucleolus, and m<sup>6</sup>A abundance returns to the basal level as cells enter into the meiotic divisions. Interestingly, researchers have found that IME4 also regulates IME1, which implies a putative positive feedback loop between m<sup>6</sup>A deposition and IME1 expression (Schwartz et al. 2013).

**Table 1.** Evolutionary conservation of nuclear RNA m<sup>6</sup>A methylation machinery

Species	Methyltransferases	Auxiliary factors	Biological roles
<i>Saccharomyces cerevisiae</i>	IME4	MUM2, SLZ1	Required for meiosis and sporulation (Clancy et al. 2002). SLZ1 localizes the complex to the nucleolus for m <sup>6</sup> A methylation (Schwartz et al. 2013).
<i>Drosophila melanogaster</i>	IME4	FL(2)D	IME4 is essential for viability (Hongay and Orr-Weaver 2011). IME4 is required for Notch signaling during oogenesis (Hongay and Orr-Weaver 2011). FL(2)D is required for splicing of <i>Sxl</i> and <i>tra</i> pre-mRNAs that are responsible for sexual determination (Penalva et al. 2000).
<i>Arabidopsis thaliana</i>	MTA	FIP37	Required for embryonic development (Zhong et al. 2008). Required for normal growth patterns, apical dominance, and plant development (Bodi et al. 2012).
<i>Danio rerio</i>	METTL3, METTL14	WTAP	METTL3 and WTAP are required for normal embryogenesis (Ping et al. 2014).
Mammals	METTL3, METTL14	WTAP	METTL3 and METTL14 regulate stem cell differentiation and reprogramming (Batista et al. 2014; Wang et al. 2014b; Geula et al. 2015). METTL3 regulates circadian periods (Fustin et al. 2013). Depletion of METTL3 and METTL14 leads to apoptosis in cancer cells (Bokar 2005). WTAP localizes METTL3–METTL14 to nucleus speckles (Ping et al. 2014). WTAP regulates cell cycle, splicing, and embryonic development (Horiuchi et al. 2006, 2013; Ping et al. 2014).

### m<sup>6</sup>A methylation in *D. melanogaster*

*D. melanogaster* IME4 shows significant amino acid similarity to and a conserved catalytic domain with its eukaryotic homologs (Table 1; Fig. 3). Unlike in yeast, elimination of the full-length *D. melanogaster* IME4 in *Drosophila* is lethal (Hongay and Orr-Weaver 2011). Partial deletion of *D. melanogaster* IME4 is semilethal, with the rare viable adults showing significantly reduced fecundity. The catalytic domain of *D. melanogaster* IME4 is required for the rescue of this semilethality (Hongay and Orr-Weaver 2011), indicating a potential role for m<sup>6</sup>A RNA methylation in metazoan development. Further studies showed that *D. melanogaster* IME4 was primarily expressed in the gonads of adult flies. In females, *D. melanogaster* IME4 plays a crucial role in oogenesis; *D. melanogaster* ime4-deficient females exhibit compound egg chambers accompanied by significant defects in the Notch signaling pathway. The ancillary factor FL(2)D (female-lethal 2 D), the homolog of yeast MUM2 and mammalian WTAP, is conserved in *Drosophila*. This protein is required for the splicing regulation of *Sxl* (Sex lethal) and *tra* (transformer) pre-mRNAs, two critical gene transcripts associated with *Drosophila* sex determination and dosage compensation (Penalva et al. 2000; Ortega et al. 2003; Penn et al. 2008).

### m<sup>6</sup>A methylation in plants

m<sup>6</sup>A is a ubiquitous modification found in the mRNAs of various plants, including monocot plants maize (Nichols 1979), wheat (Kennedy and Lane 1979), oat (Haugland and Cline 1980), *A. thaliana* (Zhong et al. 2008; Luo et al. 2014), and rice (Li et al. 2014). MTA (encoded by

At4g10760), a METTL3 ortholog in *Arabidopsis*, has been identified as an active component of the m<sup>6</sup>A methyltransferase complex (Zhong et al. 2008). MTA interacts with FIP37 (encoded by At3g54170), an *Arabidopsis* homolog of mammalian WTAP and *Drosophila* FL(2)D, highlighting the highly conserved nature of the methyltransferase components across eukaryotes (Table 1). Intriguingly, MTA tends to be expressed in higher levels in dividing tissues, such as developing seeds, shoot meristems, and emerging lateral roots (Craigon et al. 2004; Zhong et al. 2008). Disruption of either MTA or FIP37 in *Arabidopsis* leads to developmental arrest of embryos at the globular stage (Vespa et al. 2004; Zhong et al. 2008), coupled to a loss of m<sup>6</sup>A from the mRNA in arrested seeds (Vespa et al. 2004; Zhong et al. 2008). Later in development, perturbation of MTA causes multiple growth defects, including reduced apical dominance, organ abnormality, and increased trichome branching (Bodi et al. 2012). Collectively, these results demonstrate that the methyltransferase and hence m<sup>6</sup>A methylation in mRNA play a crucial role in plant development. Very recently, transcriptome-wide m<sup>6</sup>A profiling was performed in two accessions of *Arabidopsis* (Luo et al. 2014)—Can-0 and Hen-16—as well as in the rice callus and leaf (Li et al. 2014). It is worth noting that *Arabidopsis* and rice are unique in their enrichment of m<sup>6</sup>A not only around the stop codon and within 3' UTRs—as observed in yeast and mammals—but also around the start codon (Fig. 2). As genes possessing m<sup>6</sup>A sites around the start codon are associated with photosynthesis and appear to be highly expressed in *Arabidopsis*, this suggests a potential direct role of m<sup>6</sup>A at the 5' UTR during translation (Luo et al. 2014). It will be interesting to determine whether this feature observed in plants is conserved in other organisms such as mammals.



### *m<sup>6</sup>A methylation machinery in vertebrates and mammals*

We previously discussed the m<sup>6</sup>A methylation machinery of mammals in our description of writer proteins. The core m<sup>6</sup>A methyltransferase complex METTL3–METTL14–WTAP is highly conserved from zebrafish to mammals. In zebrafish, both METTL3 and WTAP proteins are ubiquitously expressed during embryogenesis and specifically enriched in the brain 36 h after fertilization (Ping et al. 2014). Embryos injected with either METTL3 or WTAP antisense morpholinos (MOs) suffer from various developmental defects, including smaller heads, eyes, and brain ventricles and curved notochord. In comparison with embryos injected with single-gene-targeted MOs, simultaneous knockdown of these two genes leads to a more pronounced phenotype in embryonic development as well as more severe decreases in the m<sup>6</sup>A level, indicating the *in vivo* synergistic effect of the methyltransferase complex. How METTL14 affects m<sup>6</sup>A deposition and zebrafish tissue differentiation remains to be studied.

Methyltransferases METTL3 and METTL14 are also shown to mediate the m<sup>6</sup>A formation in mouse ESCs (mESCs) (Batista et al. 2014; Wang et al. 2014b; Geula et al. 2015). Recent work has identified m<sup>6</sup>A as a crucial regulator in the differentiation and reprogramming of stem cells, which are discussed next.

### **Biological consequences of m<sup>6</sup>A methylation of mRNA and the underlying mechanisms**

#### *m<sup>6</sup>A RNA methylation determines stem cell fate by regulating pluripotency transition toward differentiation*


ESCs are pluripotent stem cells derived from the inner cell mass (ICM) of a preimplantation embryo, exhibiting prolonged undifferentiated proliferation and stable developmental potential to form derivatives of all three embryonic germ layers (Thomson et al. 1998). The ESCs reside in a so-called “naïve” pluripotent state, while epiblast stem cells (EpiSC) that are derived from a post-implantation epiblast reside in a more differentiation-prepared, “primed” pluripotent state (Geula et al. 2015). The transition from naïve pluripotency to differentiation is tightly regulated by a plethora of pluripotency markers and developmental factors. Transcriptome-wide m<sup>6</sup>A profiling in mESCs and human ESCs showed that the majority of these core pluripotent genes (e.g., *Nanog*, *Sox2*, *Klf4*, *Myc*, *Jarid2*, and *Smad3*) and developmental regulators (e.g., *Foxa2* and *Sox17*) have m<sup>6</sup>A modifications on their transcripts, with most of them being targets of Mettl3 (Batista et al. 2014; Wang et al. 2014b; Geula et al. 2015). Meanwhile, siRNA screening also identified Mettl3 as an epigenetic repressor that specifically destabilizes the primed EpiSCs (Geula et al. 2015). Importantly, both of the two methyltransferases, Mettl3 and Mettl14, are shown to catalyze m<sup>6</sup>A RNA deposition in mESCs (Batista et al. 2014; Wang et al. 2014b; Geula et al. 2015). Wang et al. (2014b) reported that the partial depletion of Mettl3

or Mettl14 by shRNAs leads to decreased m<sup>6</sup>A levels and reduced self-renewal of mESCs. However, in more recent studies (Batista et al. 2014; Geula et al. 2015) complete Mettl3 knockout mESCs and epiblasts were generated that actually displayed increased self-renewal but substantially impaired differentiation into mature cardiomyocytes and neurons (Batista et al. 2014). When subcutaneously injected into immunodeficient mice, *Mettl3* knockout mESCs readily generate larger but poorly differentiated teratomas *in vivo*, further indicating that depletion of m<sup>6</sup>A in mESCs enhances self-renewal but hampers differentiation (Batista et al. 2014).

Recently, Geula et al. (2015) demonstrated that the m<sup>6</sup>A modification plays a key role in facilitating transition of mESCs from the naïve state to the primed state upon differentiation (Fig. 4). To resolve the role of m<sup>6</sup>A in the naïve pluripotent state, genetic ablation of *Mettl3* was performed in mESCs, and mating the *Mettl3*<sup>+/-</sup> heterozygote mice yielded the *Mettl3*<sup>-/-</sup> knockout blastocysts. Consistent with previous results of Batista et al. (2014), Mettl3-depleted mESCs showed an almost complete loss of m<sup>6</sup>A and preserved naïve pluripotency but failed to proceed into the primed EpiSC-like state. Like *Mettl3*<sup>-/-</sup> mESCs, *Mettl14*<sup>-/-</sup> knockout mESCs resisted progression out of the naïve state. Taken together, this evidence suggests that m<sup>6</sup>A ablation in naïve mESCs impairs the transition of naïve mESCs into the primed state and hence blocks the subsequent differentiation. In contrast, mouse EpiSCs (mEpiSCs) at a primed pluripotency state showed a distinct response to m<sup>6</sup>A depletion; namely, Mettl3 knockdown in mEpiSCs resulted in attenuated stability and an enhanced tendency to lineage priming, which finally led to fast differentiation and/or cell death.

The balance between naïve pluripotency and lineage priming is fine-tuned by the relative expression of naïve pluripotency markers and lineage commitment factors. Global analysis of methylomes of naïve ESCs and primed EBs showed that m<sup>6</sup>A modification was detected in 80% of the transcripts of naïve pluripotency genes (e.g., *Nanog*, *Klf4*, *Sox2*, and *Esrrb*) as well as multiple lineage commitment regulators (e.g., *Foxa2* and *Sox17*). In general, m<sup>6</sup>A deposition in mESCs decreases the expression of methylated transcripts and directly reduces their stability. For both types of regulators, loss of m<sup>6</sup>A results in increased abundance of transcripts and longer mRNA lifetime (Batista et al. 2014; Wang et al. 2014b; Geula et al. 2015), reminiscent of the role of YTHDF2 in mediating the degradation of methylated mRNA (Wang et al. 2014a). Thus, depletion of m<sup>6</sup>A acts to boost the expression of the dominant regulators (pluripotent-promoting or lineage commitment genes) at a given pluripotency state, thereby driving stem cell differentiation. In the ground naïve state, where pluripotency-promoting transcripts prevail, Mettl3 depletion further amplifies the already highly expressed naïve pluripotency genes but leads to only a marginal increase in lineage commitment transcripts, resulting in a so-called “hypernaïve” pluripotency phenotype (Fig. 4; Batista et al. 2014; Geula et al. 2015). In the primed state, where lineage commitment transcripts dominate, Mettl3 depletion primarily up-regulates lineage commitment





**Figure 4.** Methyltransferases set m<sup>6</sup>A marks on mRNAs to balance the expression levels of pluripotency genes and lineage commitment genes in naive and primed states of the ESCs. In the naive state, the expression level of the pluripotency genes is dominant over that of lineage commitment genes, while in the primed state, the trend exhibits the opposite. The m<sup>6</sup>A methyltransferase depletion in naive pluripotent cells further up-regulates already highly abundant naive pluripotency genes, while the lineage commitment genes remain at very low residual levels. As a result, cells stay in a “hypernaive” pluripotent state and fail to progress into the primed state. If the methyltransferase depletion occurs in the primed state, the expression level of the differentiation priming markers is further boosted, which pushes cells above the critical threshold toward differentiation, leading to fast differentiation and/or cell death.

factors while leading to a minimal increase of naive pluripotency markers, further tipping the balance toward lineage priming and differentiation (Fig. 4; Geula et al. 2015). Similar divergent effects were also found when m<sup>6</sup>A was depleted in different stages of cellular reprogramming toward naive pluripotency. During the reprogramming of primed mEpiSCs to naive mESCs, early inactivation of *Mettl3* compromises the pluripotency stability of primed cells and impairs their reversion, whereas late depletion of *Mettl3* significantly enhances the reprogramming efficiency of mEpiSCs (Geula et al. 2015).

It should be noted that knockout of *Mettl3* is embryonic-lethal (Geula et al. 2015). Post-implantation embryonic day 5.5 (E5.5)–E7.5 knockout embryos retained the widespread expression of pluripotent marker *Nanog* and failed to up-regulate early differentiation markers (e.g., *Foxa2* and *Brachyury*), which recapitulated the in vitro resistance to differentiation and ultimately resulted in embryonic lethality (Batista et al. 2014; Geula et al. 2015).

Collectively, these studies showed that m<sup>6</sup>A modification precisely modulates the differentiation and reprogramming of stem cells via regulation of the expression of dominant genes involved in corresponding processes. In addition to its role in RNA stability, m<sup>6</sup>A might regulate gene expression via other pathways, such as translation and alternative splicing (Geula et al. 2015). Interestingly, protein profiling showed that loss of m<sup>6</sup>A in mESCs enhances the overall protein production level; this trend is intensified for transcripts that bear more m<sup>6</sup>A peaks. Subsequent ribosomal profiling experiments revealed that the absence of m<sup>6</sup>A in mESCs and mouse EBs resulted in a modest yet significant increase in translation efficiency, which might also contribute to the maintenance of naive pluripotency state in *Mettl3* knockout mESCs (Geula et al. 2015). Alternative splicing is affected by the presence of m<sup>6</sup>A modification in the spliced region. Depletion of m<sup>6</sup>A significantly increases the frequency of two types of alternative splicing: skipped exons and retained introns

(Geula et al. 2015). The underlying mechanism is not fully understood.

In general, dynamic mRNA modifications appear to be tightly correlated to the differentiation and reprogramming of stem cells. In addition to m<sup>6</sup>A, recent studies have characterized the distribution of pseudouridine as another widespread and dynamic modification of mRNA (Carlile et al. 2014; Schwartz et al. 2014a). Intriguingly, mutations in dyskerin, an enzyme responsible for pseudouridine formation, lead to aberrant differentiation of hematopoietic stem cells, whereas the conditional expression of dyskerin with a catalytically active domain rescues the severe defects in differentiation (Bellodi et al. 2013). Recent work has also reported efficient generation of induced pluripotent stem cells (iPSCs) from human fibroblasts by using synthetic mRNA with certain modifications. Complete substitution of pseudouridine for uridine and 5mC for cytidine in synthetic mRNAs encoding reprogramming factors attenuated the interferon-mediated innate immune response and enhanced the protein expression yield, thereby remarkably increasing the reprogramming efficiency (Warren et al. 2010).

#### *m<sup>6</sup>A RNA methylation controls the circadian clock*

The mechanism of the mammalian circadian clock involves a negative transcription–translation feedback loop in which the transcription of the clock genes is suppressed by their own encoded proteins. The period of the circadian cycle is set according to this general principle. Around 10% of the transcriptome in livers is known to be rhythmic, but only about one-fifth is driven by de novo transcription, which indicates that mRNA processing could serve as a major circadian component. Recent work showed that many clock genes as well as clock output gene transcripts bear m<sup>6</sup>A modifications (Fustin et al. 2013). Inhibition of m<sup>6</sup>A formation by silencing *METTL3* causes an mRNA processing delay and circadian period

elongation. It appears that m<sup>6</sup>A depletion prolongs nuclear retention of mature mRNAs of the clock genes *Per2* and *Arntl*. This result reveals an important physiological function of m<sup>6</sup>A methylation in setting the pace of the circadian cycle and determining clock speed and stability.

## Perspectives

The last few years have witnessed breakthrough discoveries on biological functions of m<sup>6</sup>A in mRNA, but the field is still in its infancy. Methylation specificity stands out as one of several challenging questions that remain to be addressed. In mammals, m<sup>6</sup>A occurs in only ~15% of all methylation consensus Pu[G > A]m<sup>6</sup>AC[A/C/U] motifs, and these methylated sites are primarily enriched near the stop codon, at the 3' UTR, within long exons, and at the 5' UTR. How the methylation machinery selectively targets a subset of consensus motifs in the transcriptome remains to be understood. This specificity likely has functional implications on the methylated RNAs. The METTL3–METTL14 heterodimer exhibits higher activity to the GGACU sequence located in a random structure region compared with that residing in the stem or loop (Liu et al. 2014). In agreement with the biochemistry results, global analysis also shows that methylated sites are significantly less structured when compared with randomly selected counterparts from the same genes, possibly because these sites are more exposed and accessible to the methylation machinery (Schwartz et al. 2013). However, more complicated pathways/mechanisms must be involved to achieve target selectivity. A recent study indicated that microRNAs (miRNAs) could partially regulate m<sup>6</sup>A modification via a sequence-pairing mechanism (Chen et al. 2015b), whereby miRNA expression may modulate the binding of METTL3 to mRNA substrates. Further biochemical and cellular validations are required to confirm this model. Interestingly, another recent study revealed that the m<sup>6</sup>A mark on primary miRNA (pri-miRNA) plays critical roles in miRNA maturation (Alarcon et al. 2015). METTL3 methylates pri-miRNAs, which facilitates their recognition and processing by the RNA-binding protein DGCR8 in the initiation of miRNA biogenesis. Collectively, these studies suggest a potential regulatory network between the miRNA-based regulation and the m<sup>6</sup>A-dependent regulation as two main pathways that post-transcriptionally control gene expression (Alarcon et al. 2015; Berulava et al. 2015; Chen et al. 2015b).

The multicomponent mammalian methyltransferase complex still needs to be completely resolved because auxiliary components in the complex may play roles in recruiting the catalytic core to the particular locations of the cognate pre-mRNAs and/or tuning activities of the methyltransferases. Thus, careful characterizations of proteins that interact with METTL3/METTL14/WTAP within the nuclear speckles will shed further insights on the origination of the m<sup>6</sup>A specificity.

Transcriptome-wide mapping of m<sup>6</sup>A at single-base resolution will greatly facilitate our understanding of selective m<sup>6</sup>A installation by the methyltransferase complex.

With a base-resolution m<sup>6</sup>A map, single and clustered m<sup>6</sup>A sites can be differentiated from each other; m<sup>6</sup>A fractions on particular transcripts and nearby *cis* elements can be derived. Additionally, one can study the knockout cell lines to determine whether METTL3 and METTL14 control individual groups of transcripts or share the same targets. Most m<sup>6</sup>A-seq studies to date have profiled the steady-state polyadenylated RNA inside cells, with the majority of them being mature mRNA rather than highly labile pre-mRNA. Therefore, it is necessary to carry out m<sup>6</sup>A sequencing on pre-mRNA in order to thoroughly examine the prevalence and distribution of m<sup>6</sup>A within the intronic regions and estimate the percentage of mRNAs that could be methylated either cotranscriptionally or, potentially, post-transcriptionally.

Emerging results suggest that m<sup>6</sup>A serves as a dynamic mark on a large number of mRNAs and lncRNAs, which help cells rapidly respond and/or adapt to external signaling and stimuli. By virtue of the reversible nature of the m<sup>6</sup>A modification, the stability, localization, and translatability of a large group of mRNA transcripts and lncRNAs can be regulated by m<sup>6</sup>A reader proteins and thereby participate in a timely manner in various biological pathways. The methyltransferases, demethylases, and reader proteins can all direct the methylation-based signaling process. Development of small molecule inhibitors or gene therapy tools for targeting these proteins could lead to new ways of controlling gene expression and potential new therapies for human diseases.

Last, m<sup>6</sup>A in eukaryotic mRNA exhibits substantial contributions to post-transcriptional gene expression regulation. This same modification, N<sup>6</sup>-methyladenine (6mA or m<sup>6</sup>dA), in DNA has been known to play important roles in bacterial genomes. Very recently, three independent studies reported the presence and characterizations of 6mA/m<sup>6</sup>dA in three different eukaryotic genomes (green alga, worm, and fly) with proposed transcription regulation functions (Fu et al. 2015; Greer et al. 2015; Zhang et al. 2015). Indeed, the adenine methylation appears to be a common mechanism to control gene expression.

## Acknowledgments

This work was supported by the National Institutes of Health (GM071440 to C.H.). C.H. is an investigator of the Howard Hughes Medical Institute. S.F. Reichard contributed editing.

## References

- Agarwala SD, Blitzblau HG, Hochwagen A, Fink GR. 2012. RNA methylation by the MIS complex regulates a cell fate decision in yeast. *PLoS Genet* **8**: e1002732.
- Agris PF, Vendeix FAP, Graham WD. 2007. tRNA's wobble decoding of the genome: 40 years of modification. *J Mol Biol* **366**: 1–13.
- Aik W, Scotti JS, Choi H, Gong L, Demetriades M, Schofield CJ, McDonough MA. 2014. Structure of human RNA N<sup>6</sup>-methyladenine demethylase ALKBH5 provides insights into its mechanisms of nucleic acid recognition and demethylation. *Nucleic Acids Res* **42**: 4741–4754.

- Alarcon CR, Lee H, Goodarzi H, Halberg N, Tavazoie SF. 2015. N<sup>6</sup>-methyladenosine marks primary microRNAs for processing. *Nature* **519**: 482–485.
- Batista Pedro J, Molinie B, Wang J, Qu K, Zhang J, Li L, Bouley Donna M, Lujan E, Haddad B, Daneshvar K, et al. 2014. m<sup>6</sup>A RNA modification controls cell fate transition in mammalian embryonic stem cells. *Cell Stem Cell* **15**: 707–719.
- Beemon K, Keith J. 1977. Localization of N<sup>6</sup>-methyladenosine in the Rous sarcoma virus genome. *J Mol Biol* **113**: 165–179.
- Bellodi C, McMahon M, Contreras A, Juliano D, Kopmar N, Nakamura T, Maltby D, Burlingame A, Savage SA, Shimamura A, et al. 2013. H/ACA small RNA dysfunctions in disease reveal key roles for noncoding RNA modifications in hematopoietic stem cell differentiation. *Cell Rep* **3**: 1493–1502.
- Berulava T, Rahmann S, Rademacher K, Klein-Hitpass L, Horsthemke B. 2015. N<sup>6</sup>-adenosine methylation in miRNAs. *PLoS One* **10**: e0118438.
- Bhutani N, Burns DM, Blau HM. 2011. DNA demethylation dynamics. *Cell* **146**: 866–872.
- Bird A. 2001. Methylation talk between histones and DNA. *Science* **294**: 2113–2115.
- Bjork GR, Ericson JU, Gustafsson CED, Hagervall TG, Jonsson YH, Wikstrom PM. 1987. Transfer RNA Modification. *Annu Rev Biochem* **56**: 263–285.
- Bodi Z, Button JD, Grierson D, Fray RG. 2010. Yeast targets for mRNA methylation. *Nucleic Acids Res* **38**: 5327–5335.
- Bodi Z, Zhong S, Mehra S, Song J, Graham N, Li H, May S, Fray RG. 2012. Adenosine methylation in *Arabidopsis* mRNA is associated with the 3' end and reduced levels cause developmental defects. *Front Plant Sci* **3**: 48.
- Bokar JA. 2005. The biosynthesis and functional roles of methylated nucleosides in eukaryotic mRNA. In *Topics in Current Genetics* Vol. 12, *Fine-tuning of RNA functions by modification and editing* (ed. Grosjean H), pp. 141–177. Springer, Heidelberg.
- Bokar JA, Shambaugh ME, Polayes D, Matera AG, Rottman FM. 1997. Purification and cDNA cloning of the AdoMet-binding subunit of the human mRNA (N<sup>6</sup>-adenosine)-methyltransferase. *RNA* **3**: 1233–1247.
- Bujnicki JM, Feder M, Radlinska M, Blumenthal RM. 2002. Structure prediction and phylogenetic analysis of a functionally diverse family of proteins homologous to the MT-A70 subunit of the human mRNA:m<sup>6</sup>A methyltransferase. *J Mol Evol* **55**: 431–444.
- Cantara WA, Crain PF, Rozenski J, McCloskey JA, Harris KA, Zhang X, Vendeix FAP, Fabris D, Agris PF. 2011. The RNA modification database, RNAMDB: 2011 update. *Nucleic Acids Res* **39**: D195–D201.
- Carlile TM, Rojas-Duran MF, Zinshteyn B, Shin H, Bartoli KM, Gilbert WV. 2014. Pseudouridine profiling reveals regulated mRNA pseudouridylation in yeast and human cells. *Nature* **515**: 143–146.
- Chan CTY, Pang YLJ, Deng W, Babu IR, Dyavaiah M, Begley TJ, Dedon PC. 2012. Reprogramming of tRNA modifications controls the oxidative stress response by codon-biased translation of proteins. *Nat Commun* **3**: 937.
- Chen K, Lu Z, Wang X, Fu Y, Luo GZ, Liu N, Han D, Dominissini D, Dai Q, Pan T, et al. 2015a. High-resolution N<sup>6</sup>-methyladenosine (m<sup>6</sup>A) map using photo-crosslinking-assisted m<sup>6</sup>A sequencing. *Angew Chem Int Ed Engl* **54**: 1587–1590.
- Chen T, Hao YJ, Zhang Y, Li MM, Wang M, Han W, Wu Y, Lv Y, Hao J, Wang L, et al. 2015b. m<sup>6</sup>A RNA methylation is regulated by microRNAs and promotes reprogramming to pluripotency. *Cell Stem Cell* **16**: 289–301.
- Chu S, Herskowitz I. 1998. Gametogenesis in yeast is regulated by a transcriptional cascade dependent on Ndt80. *Mol Cell* **1**: 685–696.
- Clancy MJ, Shambaugh ME, Timpte CS, Bokar JA. 2002. Induction of sporulation in *Saccharomyces cerevisiae* leads to the formation of N<sup>6</sup>-methyladenosine in mRNA: a potential mechanism for the activity of the IME4 gene. *Nucleic Acids Res* **30**: 4509–4518.
- Craigon DJ, James N, Okyere J, Higgins J, Jotham J, May S. 2004. NASCArrays: a repository for microarray data generated by NASC's transcriptomics service. *Nucleic Acids Res* **32**: D575–D577.
- Csepány T, Lin A, Baldick CJ, Beemon K. 1990. Sequence specificity of mRNA N<sup>6</sup>-adenosine methyltransferase. *J Biol Chem* **265**: 20117–20122.
- Desrosiers R, Friderici K, Rottman F. 1974. Identification of methylated nucleosides in messenger RNA from Novikoff hepatoma cells. *Proc Natl Acad Sci* **71**: 3971–3975.
- Dina C, Meyre D, Gallina S, Durand E, Korner A, Jacobson P, Carlsson LMS, Kiess W, Vatin V, Lecoœur C, et al. 2007. Variation in FTO contributes to childhood obesity and severe adult obesity. *Nat Genet* **39**: 724–726.
- Dominissini D, Moshitch-Moshkovitz S, Schwartz S, Salmon-Divon M, Ungar L, Osenberg S, Cesarkas K, Jacob-Hirsch J, Amariglio N, Kupiec M, et al. 2012. Topology of the human and mouse m<sup>6</sup>A RNA methylomes revealed by m<sup>6</sup>A-seq. *Nature* **485**: 201–206.
- Fischer J, Koch L, Emmerling C, Vierkotten J, Peters T, Bruning JC, Ruther U. 2009. Inactivation of the FTO gene protects from obesity. *Nature* **458**: 894–898.
- Frayling TM, Timpson NJ, Weedon MN, Zeggini E, Freathy RM, Lindgren CM, Perry JRB, Elliott KS, Lango H, Rayner NW, et al. 2007. A common variant in the FTO gene is associated with body mass index and predisposes to childhood and adult obesity. *Science* **316**: 889–894.
- Fu Y, Dai Q, Zhang W, Ren J, Pan T, He C. 2010. The AlkB domain of mammalian ABH8 catalyzes hydroxylation of 5-methoxycarbonylmethyluridine at the wobble position of tRNA. *Angew Chem Int Ed Engl* **49**: 8885–8888.
- Fu Y, Jia G, Pang X, Wang RN, Wang X, Li CJ, Smemo S, Dai Q, Bailey KA, Nobrega MA, et al. 2013. FTO-mediated formation of N<sup>6</sup>-hydroxymethyladenosine and N<sup>6</sup>-formyladenosine in mammalian RNA. *Nat Commun* **4**: 1798.
- Fu Y, Dominissini D, Rechavi G, He C. 2014. Gene expression regulation mediated through reversible m<sup>6</sup>A RNA methylation. *Nat Rev Genet* **15**: 293–306.
- Fu Y, Luo G-Z, Chen K, Deng X, Yu M, Han D, Hao Z, Liu J, Lu X, Doré Louis C, et al. 2015. N<sup>6</sup>-methyldeoxyadenosine marks active transcription start sites in *Chlamydomonas*. *Cell* **161**: 879–892.
- Fustin JM, Doi M, Yamaguchi Y, Hida H, Nishimura S, Yoshida M, Isagawa T, Morioka MS, Takeya H, Manabe I, et al. 2013. RNA-methylation-dependent RNA processing controls the speed of the circadian clock. *Cell* **155**: 793–806.
- Gerken T, Girard CA, Tung Y-CL, Webby CJ, Saudek V, Hewitson KS, Yeo GSH, McDonough MA, Cunliffe S, McNeill LA, et al. 2007. The obesity-associated FTO gene encodes a 2-oxoglutarate-dependent nucleic acid demethylase. *Science* **318**: 1469–1472.
- Geula S, Moshitch-Moshkovitz S, Dominissini D, Mansour AA, Kol N, Salmon-Divon M, Herskowitz V, Peer E, Mor N, Manor YS, et al. 2015. m<sup>6</sup>A mRNA methylation facilitates resolution of naïve pluripotency toward differentiation. *Science* **347**: 1002–1006.



- Greer EL, Blanco MA, Gu L, Sendinc E, Liu J, Aristizábal-Corrales D, Hsu C-H, Aravind L, He C, Shi Y. 2015. DNA methylation on N6-adenine in *C. elegans*. *Cell* **161**: 868–878.
- Gu J, Patton JR, Shimba S, Reddy R. 1996. Localization of modified nucleotides in *Schizosaccharomyces pombe* spliceosomal small nuclear RNAs: modified nucleotides are clustered in functionally important regions. *RNA* **2**: 909–918.
- Gulati P, Cheung MK, Antrobus R, Church CD, Harding HP, Tung YC, Rimmington D, Ma M, Ron D, Lehner PJ, et al. 2013. Role for the obesity-related FTO gene in the cellular sensing of amino acids. *Proc Natl Acad Sci* **110**: 2557–2562.
- Haugland RA, Cline MG. 1980. Post-transcriptional modifications of oat coleoptile ribonucleic acids. *Eur J Biochem* **104**: 271–277.
- He C. 2010. RNA epigenetics? *Nat Chem Biol* **6**: 863–865.
- He Y-F, Li B-Z, Li Z, Liu P, Wang Y, Tang Q, Ding J, Jia Y, Chen Z, Li L, et al. 2011. Tet-mediated formation of 5-carboxylcytosine and its excision by TDG in mammalian DNA. *Science* **333**: 1303–1307.
- Hess ME, Hess S, Meyer KD, Verhagen LAW, Koch L, Bronneke HS, Dietrich MO, Jordan SD, Saletore Y, Elemento O, et al. 2013. The fat mass and obesity associated gene (Fto) regulates activity of the dopaminergic midbrain circuitry. *Nat Neurosci* **16**: 1042–1048.
- Hongay CF, Orr-Weaver TL. 2011. *Drosophila* inducer of meiosis 4 (IME4) is required for Notch signaling during oogenesis. *Proc Natl Acad Sci* **108**: 14855–14860.
- Hongay CF, Grisafi PL, Galitski T, Fink GR. 2006. Antisense transcription controls cell fate in *Saccharomyces cerevisiae*. *Cell* **127**: 735–745.
- Horiuchi K, Umetani M, Minami T, Okayama H, Takada S, Yamamoto M, Aburatani H, Reid PC, Housman DE, Hamakubo T, et al. 2006. Wilms' tumor 1-associating protein regulates G(2)/M transition through stabilization of cyclin A2 mRNA. *Proc Natl Acad Sci* **103**: 17278–17283.
- Horiuchi K, Kawamura T, Iwanari H, Ohashi R, Naito M, Kodama T, Hamakubo T. 2013. Identification of Wilms' tumor 1-associating protein complex and its role in alternative splicing and the cell cycle. *J Biol Chem* **288**: 33292–33302.
- Horowitz S, Horowitz A, Nilsen TW, Munns TW, Rottman FM. 1984. Mapping of N<sup>6</sup>-methyladenosine residues in bovine prolactin mRNA. *Proc Natl Acad Sci* **81**: 5667–5671.
- Ito S, D'Alessio AC, Taranova OV, Hong K, Sowers LC, Zhang Y. 2010. Role of Tet proteins in 5mC to 5hmC conversion, ES-cell self-renewal and inner cell mass specification. *Nature* **466**: 1129–1133.
- Ito S, Shen L, Dai Q, Wu SC, Collins LB, Swenberg JA, He C, Zhang Y. 2011. Tet proteins can convert 5-methylcytosine to 5-formylcytosine and 5-carboxylcytosine. *Science* **333**: 1300–1303.
- Jia G, Yang C-G, Yang S, Jian X, Yi C, Zhou Z, He C. 2008. Oxidative demethylation of 3-methylthymine and 3-methyluracil in single-stranded DNA and RNA by mouse and human FTO. *FEBS Lett* **582**: 3313–3319.
- Jia G, Fu Y, Zhao X, Dai Q, Zheng G, Yang Y, Yi C, Lindahl T, Pan T, Yang Y-G, et al. 2011. N<sup>6</sup>-methyladenosine in nuclear RNA is a major substrate of the obesity-associated FTO. *Nat Chem Biol* **7**: 885–887.
- Jones PA. 2012. Functions of DNA methylation: islands, start sites, gene bodies and beyond. *Nat Rev Genet* **13**: 484–492.
- Kennedy TD, Lane BG. 1979. Wheat embryo ribonucleates. XIII. Methyl-substituted nucleoside constituents and 5'-terminal dinucleotide sequences in bulk poly(A)-rich RNA from imbibing wheat embryos. *Can J Biochem* **57**: 927–931.
- Kirchner S, Ignatova Z. 2015. Emerging roles of tRNA in adaptive translation, signalling dynamics and disease. *Nat Rev Genet* **16**: 98–112.
- Kohli RM, Zhang Y. 2013. TET enzymes, TDG and the dynamics of DNA demethylation. *Nature* **502**: 472–479.
- Krug RM, Morgan MA, Shatkin AJ. 1976. Influenza viral mRNA contains internal N<sup>6</sup>-methyladenosine and 5'-terminal 7-methylguanosine in cap structures. *J Virol* **20**: 45–53.
- Kurowski M, Bhagwat A, Papaj G, Bujnicki J. 2003. Phylogenomic identification of five new human homologs of the DNA repair enzyme AlkB. *BMC Genomics* **4**: 48.
- Lavi S, Shatkin AJ. 1975. Methylated simian virus 40-specific RNA from nuclei and cytoplasm of infected BSC-1 cells. *Proc Natl Acad Sci* **72**: 2012–2016.
- Li Y, Wang X, Li C, Hu S, Yu J, Song S. 2014. Transcriptome-wide N<sup>6</sup>-methyladenosine profiling of rice callus and leaf reveals the presence of tissue-specific competitors involved in selective mRNA modification. *RNA Biol* **11**: 1180–1188.
- Little NA, Hastie ND, Davies RC. 2000. Identification of WTAP, a novel Wilms' tumour 1-associating protein. *Hum Mol Genet* **9**: 2231–2239.
- Liu J, Yue Y, Han D, Wang X, Fu Y, Zhang L, Jia G, Yu M, Lu Z, Deng X, et al. 2014. A METTL3–METTL14 complex mediates mammalian nuclear RNA N<sup>6</sup>-adenosine methylation. *Nat Chem Biol* **10**: 93–95.
- Liu N, Dai Q, Zheng G, He C, Parisien M, Pan T. 2015. N<sup>6</sup>-methyladenosine-dependent RNA structural switches regulate RNA–protein interactions. *Nature* **518**: 560–564.
- Loos RJ, Yeo GS. 2014. The bigger picture of FTO—the first GWAS-identified obesity gene. *Nat Rev Endocrinol* **10**: 51–61.
- Luo G-Z, MacQueen A, Zheng G, Duan H, Dore LC, Lu Z, Liu J, Chen K, Jia G, Bergelson J, et al. 2014. Unique features of the m<sup>6</sup>A methylome in *Arabidopsis thaliana*. *Nat Commun* **5**: 5630.
- Machnicka MA, Milanowska K, Osman Oglou O, Purta E, Kurkowska M, Olchowik A, Januszewski W, Kalinowski S, Dunin-Horkawicz S, Rother KM, et al. 2013. MODOMICS: a database of RNA modification pathways—2013 update. *Nucleic Acids Res* **41**: D262–D267.
- Maden BE. 1990. The numerous modified nucleotides in eukaryotic ribosomal RNA. *Prog Nucleic Acid Res Mol Biol* **39**: 241–303.
- Meyer KD, Jaffrey SR. 2014. The dynamic epitranscriptome: N<sup>6</sup>-methyladenosine and gene expression control. *Nat Rev Mol Cell Biol* **15**: 313–326.
- Meyer KD, Saletore Y, Zumbo P, Elemento O, Mason CE, Jaffrey SR. 2012. Comprehensive analysis of mRNA methylation reveals enrichment in 3' UTRs and near stop codons. *Cell* **149**: 1635–1646.
- Motorin Y, Helm M. 2011. RNA nucleotide methylation. *Wiley Interdiscip Rev RNA* **2**: 611–631.
- Narayan P, Rottman FM. 1988. An invitro system for accurate methylation of internal adenosine residues in messenger RNA. *Science* **242**: 1159–1162.
- Narayan P, Ludwiczak RL, Goodwin EC, Rottman FM. 1994. Context effects on N<sup>6</sup>-adenosine methylation sites in prolactin mRNA. *Nucleic Acids Res* **22**: 419–426.
- Nichols JL. 1979. N<sup>6</sup>-methyladenosine in maize poly(A)-containing RNA. *Plant Sci Lett* **15**: 357–361.
- Ortega A, Niksic M, Bachi A, Wilm M, Sánchez L, Hastie N, Valcárcel J. 2003. Biochemical function of female-lethal (2)D/Wilms' tumor suppressor-1-associated proteins in alternative pre-mRNA splicing. *J Biol Chem* **278**: 3040–3047.



- Pastor WA, Aravind L, Rao A. 2013. TETonic shift: biological roles of TET proteins in DNA demethylation and transcription. *Nat Rev Mol Cell Biol* **14**: 341–356.
- Penalva LO, Ruiz MF, Ortega A, Granadino B, Vicente L, Segarra C, Valcárcel J, Sánchez L. 2000. The *Drosophila* fl(2)d gene, required for female-specific splicing of Sxl and tra pre-mRNAs, encodes a novel nuclear protein with a HQ-rich domain. *Genetics* **155**: 129–139.
- Penn JK, Graham P, Deshpande G, Calhoun G, Chaouki AS, Salz HK, Schedl P. 2008. Functioning of the *Drosophila* Wilms'-tumor-1-associated protein homolog, Fl(2)d, in sex-lethal-dependent alternative splicing. *Genetics* **178**: 737–748.
- Perry RP, Kelley DE. 1974. Existence of methylated messenger RNA in mouse L cells. *Cell* **1**: 37–42.
- Piekna-Przybylska D, Decatur WA, Fournier MJ. 2008. The 3D rRNA modification maps database: with interactive tools for ribosome analysis. *Nucleic Acids Res* **36**: D178–D183.
- Ping XL, Sun BF, Wang L, Xiao W, Yang X, Wang W-J, Adhikari S, Shi Y, Lv Y, Chen Y-S, et al. 2014. Mammalian WTAP is a regulatory subunit of the RNA N<sup>6</sup>-methyladenosine methyltransferase. *Cell Res* **24**: 177–189.
- Rottman F, Shatkin AJ, Perry RP. 1974. Sequences containing methylated nucleotides at the 5' termini of messenger RNAs: possible implications for processing. *Cell* **3**: 197–199.
- Schibler U, Kelley DE, Perry RP. 1977. Comparison of methylated sequences in messenger RNA and heterogeneous nuclear RNA from mouse L cells. *J Mol Biol* **115**: 695–714.
- Schwartz S, Agarwala SD, Mumbach MR, Jovanovic M, Mertins P, Shishkin A, Tabach Y, Mikkelsen TS, Satija R, Ruvkun G, et al. 2013. High-resolution mapping reveals a conserved, widespread, dynamic mRNA methylation program in yeast meiosis. *Cell* **155**: 1409–1421.
- Schwartz S, Bernstein DA, Mumbach MR, Jovanovic M, Herbst RH, León-Ricardo BX, Engreitz JM, Guttman M, Satija R, Lander ES, et al. 2014a. Transcriptome-wide mapping reveals widespread dynamic-regulated pseudouridylation of ncRNA and mRNA. *Cell* **159**: 148–162.
- Schwartz S, Mumbach MR, Jovanovic M, Wang T, Maciag K, Bushkin GG, Mertins P, Ter-Ovanesyan D, Habib N, Cacciarelli D, et al. 2014b. Perturbation of m<sup>6</sup>A writers reveals two distinct classes of mRNA methylation at internal and 5' sites. *Cell Rep* **8**: 284–296.
- Shah JC, Clancy MJ. 1992. IME4, a gene that mediates MAT and nutritional control of meiosis in *Saccharomyces cerevisiae*. *Mol Cell Biol* **12**: 1078–1086.
- Shen L, Song C-X, He C, Zhang Y. 2014. Mechanism and function of oxidative reversal of DNA and RNA methylation. *Annu Rev Biochem* **83**: 585–614.
- Shimba S, Bokar JA, Rottman F, Reddy R. 1995. Accurate and efficient N-6-adenosine methylation in spliceosomal U6 small nuclear RNA by HeLa cell extract in vitro. *Nucleic Acids Res* **23**: 2421–2426.
- Spitale RC, Flynn RA, Zhang QC, Crisalli P, Lee B, Jung J-W, Kuchelmeister HY, Batista PJ, Torre EA, Kool ET, et al. 2015. Structural imprints in vivo decode RNA regulatory mechanisms. *Nature* **519**: 486–490.
- Strahl BD, Allis CD. 2000. The language of covalent histone modifications. *Nature* **403**: 41–45.
- Suzuki MM, Bird A. 2008. DNA methylation landscapes: provocative insights from epigenomics. *Nat Rev Genet* **9**: 465–476.
- Tahiliani M, Koh KP, Shen Y, Pastor WA, Bandukwala H, Brudno Y, Agarwal S, Iyer LM, Liu DR, Aravind L, et al. 2009. Conversion of 5-methylcytosine to 5-hydroxymethylcytosine in mammalian DNA by MLL partner TET1. *Science* **324**: 930–935.
- Thomson JA, Itskovitz-Eldor J, Shapiro SS, Waknitz MA, Swiergiel JJ, Marshall VS, Jones JM. 1998. Embryonic stem cell lines derived from human blastocysts. *Science* **282**: 1145–1147.
- Vespa L, Vachon G, Berger F, Perazza D, Faure J-D, Herzog M. 2004. The immunophilin-interacting protein AtFIP37 from *Arabidopsis* is essential for plant development and is involved in trichome endoreduplication. *Plant Physiol* **134**: 1283–1292.
- Vujovic P, Stamenkovic S, Jasnic N, Lalic I, Djurasevic SF, Cvijic G, Djordjevic J. 2013. Fasting induced cytoplasmic Fto expression in some neurons of rat hypothalamus. *PLoS One* **8**: e63694.
- Wan Y, Qu K, Zhang QC, Flynn RA, Manor O, Ouyang Z, Zhang J, Spitale RC, Snyder MP, Segal E, et al. 2014. Landscape and variation of RNA secondary structure across the human transcriptome. *Nature* **505**: 706–709.
- Wang X, Lu Z, Gomez A, Hon GC, Yue Y, Han D, Fu Y, Parisien M, Dai Q, Jia G, et al. 2014a. N<sup>6</sup>-methyladenosine-dependent regulation of messenger RNA stability. *Nature* **505**: 117–120.
- Wang Y, Li Y, Toth JI, Petroski MD, Zhang Z, Zhao JC. 2014b. N<sup>6</sup>-methyladenosine modification destabilizes developmental regulators in embryonic stem cells. *Nat Cell Biol* **16**: 191–198.
- Wang X, Zhao BS, Roundtree IA, Lu Z, Han D, Ma H, Weng X, Chen K, Shi H, He C. 2015. N<sup>6</sup>-methyladenosine modulates messenger RNA translation efficiency. *Cell* **161**: 1388–1399.
- Warren L, Manos PD, Ahfeldt T, Loh Y-H, Li H, Lau F, Ebina W, Mandal P, Smith ZD, Meissner A, et al. 2010. Highly efficient reprogramming to pluripotency and directed differentiation of human cells using synthetic modified mRNA. *Cell Stem Cell* **7**: 618–630.
- Wei CM, Moss B. 1977. Nucleotide sequences at the N<sup>6</sup>-methyladenosine sites of HeLa cell messenger ribonucleic acid. *Biochemistry* **16**: 1672–1676.
- Wei CM, Gershowitz A, Moss B. 1975. Methylated nucleotides block 5' terminus of HeLa cell messenger RNA. *Cell* **4**: 379–386.
- Xu C, Liu K, Tempel W, Demetriades M, Aik W, Schofield CJ, Min J. 2014a. Structures of human ALKBH5 demethylase reveal a unique binding mode for specific single stranded m<sup>6</sup>A RNA demethylation. *J Biol Chem* **289**: 17299–17311.
- Xu C, Wang X, Liu K, Roundtree IA, Tempel W, Li Y, Lu Z, He C, Min J. 2014b. Structural basis for selective binding of m<sup>6</sup>A RNA by the YTHDC1 YTH domain. *Nat Chem Biol* **10**: 927–929.
- Yi C, Pan T. 2011. Cellular dynamics of RNA modification. *Acc Chem Res* **44**: 1380–1388.
- Zhang L, Lu X, Lu J, Liang H, Dai Q, Xu G-L, Luo C, Jiang H, He C. 2012. Thymine DNA glycosylase specifically recognizes 5-carboxylcytosine-modified DNA. *Nat Chem Biol* **8**: 328–330.
- Zhang G, Huang H, Liu D, Cheng Y, Liu X, Zhang W, Yin R, Zhang D, Zhang P, Liu J, et al. 2015. N<sup>6</sup>-methyladenine DNA modification in *Drosophila*. *Cell* **161**: 893–906.
- Zhao X, Yang Y, Sun BF, Shi Y, Yang X, Xiao W, Hao YJ, Ping X-L, Chen YS, Wang WJ, et al. 2014. FTO-dependent demethylation of N<sup>6</sup>-methyladenosine regulates mRNA splicing and is required for adipogenesis. *Cell Res* **24**: 1403–1419.
- Zheng G, Dahl JA, Niu Y, Fedorcsak P, Huang C-M, Li CJ, Vågbø CB, Shi Y, Wang W-L, Song S-H, et al. 2013. ALKBH5 is a mammalian RNA demethylase that impacts RNA metabolism and mouse fertility. *Mol Cell* **49**: 18–29.
- Zheng G, Fu Y, He C. 2014. Nucleic acid oxidation in DNA damage repair and epigenetics. *Chem Rev* **114**: 4602–4620.
- Zhong S, Li H, Bodi Z, Button J, Vespa L, Herzog M, Fray RG. 2008. MTA is an *Arabidopsis* messenger RNA adenosine methylase and interacts with a homolog of a sex-specific splicing factor. *Plant Cell* **20**: 1278–1288.