

## Targeting the SARS-CoV-2 Main Protease to Repurpose Drugs for COVID-19

Erol C. Vatansever<sup>a, †</sup>, Kai Yang<sup>a, †</sup>, Kaci C. Kratch<sup>a, †</sup>, Aleksandra Drelich<sup>b</sup>, Chia-Chuan Cho<sup>a</sup>, Drake M. Mellot<sup>c</sup>, Shiqing Xu,<sup>a</sup> Chien-Te K. Tseng<sup>b</sup>, and Wenshe Ray Liu<sup>a,c,d,\*</sup>

<sup>a</sup>The Texas A&M Drug Discovery Laboratory, Department of Chemistry, Texas A&M University, College Station, TX 77843, USA

<sup>b</sup>Department of Microbiology and Immunology, University of Texas Medical Branch, Galveston, TX 77555, USA

<sup>b</sup>Department of Biochemistry and Biophysics, Texas A&M University, College Station, TX 77843, USA

<sup>c</sup>Department of Molecular and Cellular Medicine, College of Medicine, Texas A&M University, College Station, TX 77843, USA

<sup>†</sup>Contribute equally to the paper.

\*Correspondence should be addressed to Wenshe Ray Liu: [wliu@chem.tamu.edu](mailto:wliu@chem.tamu.edu).

Keywords: COVID-19, SARS-CoV-2, bepridil, pimozide, ebastine

## ABSTRACT

Guided by a computational docking analysis, about 30 FDA/EMA-approved small molecule medicines were characterized on their inhibition of the SARS-CoV-2 main protease ( $M^{\text{Pro}}$ ). Of these tested small molecule medicines, six displayed an  $IC_{50}$  value in inhibiting  $M^{\text{Pro}}$  below 100  $\mu\text{M}$ . Three medicines pimozone, ebastine, and bepridil are basic small molecules that are expected to exert a similar effect as hydroxychloroquine in raising endosomal pH for slowing down the SARS-CoV-2 entry into human cell hosts. Bepridil has been previously explored in a high dose as 100 mg/kg for treating diseases. Its high dose use will likely achieve dual functions in treating COVID-19 by both raising the endosomal pH to slow viral entry and inhibiting  $M^{\text{Pro}}$  in infected cells. Therefore, the current study urges serious considerations of using bepridil in COVID-19 clinical tests.

## INTRODUCTION

The current worldwide impact of the COVID-19 pandemic has been so profound that it is often compared to that of 1918 influenza pandemic.<sup>1,2</sup> While this manuscript was under preparation, the total global COVID-19 cases had surpassed 4 million among which close to 300,000 had succumbed to death.<sup>3</sup> A modelling study has predicted that this pandemic will continue to affect everyday life and the circumstances may require societies to follow social distancing until 2022.<sup>4</sup> Finding timely treatment options are of tremendous importance to alleviate catastrophic damages of COVID-19. However, the time window that is paramount to contain the disease is extremely challenging to a conventional drug discovery process that requires typically many years to finalize a drug and therefore might not achieve its goal before the pandemic ceases. In this January, we did a comparative biochemical analysis between severe acute respiratory syndrome-coronavirus 2 (SARS-CoV-2), the virus that has caused COVID-19, and SARS-CoV that led to an epidemic in China in 2003 and proposed that remdesivir was a viable choice for the treatment of COVID-19.<sup>5,6</sup> We were excited to see that remdesivir was finally approved for emergency use in the United States and for use in Japan for people with severe symptoms. With only one medicine in stock right now, the virus may easily evade it, leading us once again with no medicine to use. Given the rapid spread and the high fatality of COVID-19, finding alternative medicines is imperative. Drug repurposing stands out as an attractive option in the current situation. If an approved drug can be identified to treat COVID-19, it can be quickly proceeded to clinical trials and manufactured at a large scale using its existing GMP lines. Previously, encouraging results were obtained from repurposing small molecule medicines including teicoplanin, ivermectin, itraconazole, and nitazoxanide.<sup>7-10</sup> These antimicrobial agents were found effective against virus infections.<sup>11</sup> However, a common drawback of all these repurposed drugs is their low efficacy level. One way to circumvent this problem is to combine multiple existing medicines to accrue a synergistic effect. To be able to discover such combinations, breaking down the druggable targets of the SARS-CoV-2 to identify drugs that do not cross-act on each other's targets is a promising strategy. For example, a recent study showed that triple combination of interferon beta-1b, lopinavir-ritonavir, and ribavirin was safe and superior to lopinavir-ritonavir alone for treating COVID-19 patients.<sup>12</sup>

In our January paper, we recommended four SARS-CoV-2 essential proteins including Spike, RNA-dependent RNA polymerase, the main protease (M<sup>Pro</sup>), and papain-like protease as drug

targets for the development of novel anti-COVID-19 medicines. Among these four proteins, M<sup>Pro</sup> that is also called 3C-like protease (3CL<sup>Pro</sup>) provides the most facile opportunity for drug repurposing owing to the ease of its biochemical assay. M<sup>Pro</sup> is a cysteine protease that processes itself and then cleaves a number of nonstructural viral proteins from two polypeptide translates that are made from the virus RNA in the human cell host.<sup>13</sup> Its relatively large active site pocket and a highly nucleophilic, catalytic cysteine residue make it likely to be inhibited by a host of existing and investigational drugs. Previous work has disclosed some existing drugs that inhibit M<sup>Pro</sup>.<sup>14</sup> However, complete characterization of existing drugs on the inhibition of M<sup>Pro</sup> is not yet available. Since the release of the first M<sup>Pro</sup> crystal structure, many computational studies have been carried out to screen existing drugs in their inhibition of M<sup>Pro</sup> and many potent leads have been proposed.<sup>15-18</sup> However, most of these lead drugs have not yet been confirmed experimentally. To investigate whether some existing drugs can potently inhibit M<sup>Pro</sup>, we have docked a group of selected FDA/EMA-approved small molecule medicines to the active site of M<sup>Pro</sup> and selected about 30 hit drugs to characterize their inhibition on M<sup>Pro</sup> experimentally. Our results revealed that a number of FDA/EMA-approved small molecule medicines do have high potency in inhibiting M<sup>Pro</sup> and therefore encourage the possible use of these medicines in fighting COVID-19.

## RESULTS & DISCUSSION

Deng et al. released the first crystal structure of M<sup>Pro</sup> on Feb 5th, 2020.<sup>14</sup> We chose this structure (the pdb entry 6LU7) as the basis for our initial docking study. M<sup>Pro</sup> has a very large active site that consists of several smaller pockets for the recognition of amino acid residues in its protein substrates. Three pockets that bind the P1, P2, and P4 residues in a protein substrate potentially interact with aromatic and large hydrophobic moieties.<sup>19</sup> Although the P1' residue in a protein substrate is a small residue such as glycine or serine, the previous studies based on the same functional enzyme from SARS-CoV showed that an aromatic moiety can occupy the site that originally bind the P1' and P2' residues in a substrate.<sup>20</sup> Based on this analysis of the M<sup>Pro</sup> structure, we selected 55 FDA/EMA-approved small molecule medicines that have several aromatic or large hydrophobic moieties inter-connected and did a deep docking analysis of their binding to M<sup>Pro</sup>. Some of the small molecule medicines used in our docking study were previously reported in other computational studies.<sup>15-18</sup> Autodock was the program we adopted for the docking analysis.<sup>21</sup> The covalent ligand and non-bonded small molecules in the structure of 6LU7 was removed for the preparation of the protein structure for the docking. Four residues His41, Met49, Asn142, and Gln189 that have shown conformational variations in the SARS-CoV enzyme were set as flexible residues during the docking process. We carried out a genetic algorithm method with 100 runs to dock each small molecule medicine to the enzyme. We collected the lowest binding energy from the total 100 runs for each small molecule medicine and summarized them in Table 1. Among all 55 small molecule drugs that we used in the docking study, 29 showed a binding energy lower than -8.3 kcal/mol. We chose these molecules to do further experimental characterizations.

To express M<sup>Pro</sup> for experimental characterizations of 29 selected small molecule medicines, we fused the M<sup>Pro</sup> gene to a superfolder green fluorescent protein (sfGFP) gene and a 6xHis tag at its 5' and 3' ends respectively in a pBAD-sfGFP plasmid that we used previously in the lab. SfGFP is known to stabilize proteins when it is genetically fused to them. We designed a TEV protease cleavage site between sfGFP and M<sup>Pro</sup> for the TEV-catalyzed proteolytic release of M<sup>Pro</sup> from sfGFP after we expressed and purified the fusion protein. We placed the 6xHis tag right after the M<sup>Pro</sup> C-terminus. The addition of this tag was for straightforward purification with Ni-NTA resins. We expected that the TEV protease cleavage of sfGFP would activate M<sup>Pro</sup> to cleave the C-terminal 6xHis tag so that a finally intact M<sup>Pro</sup> protein would be obtained. We carried out the expression in

TOP10 cells. To our surprise, after expression there was a minimal amount of the fusion protein that we were able to purify. The analysis of the cell lysate showed clearly the cleavage of a substantial amount of M<sup>Pro</sup> from sfGFP. Since we were not able to enrich the cleaved M<sup>Pro</sup> using Ni-NTA resins, the C-terminal 6xHis tag was apparently cleaved as well. TEV protease is a cysteine protease that cleaves after the Gln residue in the sequence Glu-Asn-Leu-Tyr-Phe-Gln-(Gly/Ser).<sup>22</sup> M<sup>Pro</sup> is known to cleave the sequence Thr-Val-Leu-Gln-(Gly/Ser).<sup>23</sup> The two cleavage sites share a same P1 residue. It was evident in our expression work that M<sup>Pro</sup> is able to efficiently cleave the TEV protease cutting site to mature inside *E. coli* cells. It is likely that M<sup>Pro</sup> has a substrate promiscuity higher than what we have learnt from the SARS-CoV enzyme. This aspect needs to be investigated further to help understand the virus replication inside host cells. To purify the cleaved and matured M<sup>Pro</sup>, we used ammonium sulfate to precipitate it from the cell lysate and then used the ion exchange and size exclusion chromatography to isolate it to more than 95% purity. We designed and synthesized a fluorogenic coumarin-based hexapeptide substrate (Sub1) and a FRET-based decapeptide substrate (Sub2) and acquired a commercial FRET-based tetradecapeptide substrate (Sub3) (Figure 1A). The test of enzyme activities on the three substrates indicated that the enzyme had low activity toward Sub1 under our assay conditions (Figure 1B) and its activity on Sub3 was higher than that on Sub2 (Figure 1C). Sub3 was subsequently used in all following inhibition analysis. To identify an optimal enzyme concentration for use in our inhibition analysis, we tested activities of different concentrations of M<sup>Pro</sup> on 10  $\mu$ M Sub3, the detected catalytic rate of the Sub3 cleavage was not proportional to the enzyme concentration (Figure 1D). When the enzyme concentration decreased from 50 nM to 10 nM, the Sub3 cleavage rate dropped roughly proportionally to the square of the concentration decrease, characteristics of second-order kinetics. This observation supports previous claims that the enzyme needs to dimerize in order to be active. In all the following assays, 50 nM M<sup>Pro</sup> and 10  $\mu$ M Sub3 were used throughout.

We purchased all 29 small molecule medicines from commercial providers without further purification and characterization. Rupintrivir is a previously developed 3C protease inhibitor.<sup>24</sup> It has a key lactone side chain in the P1 residue that has a demonstrated role in tight binding to 3C and 3CL proteases. Since it has been an investigational antiviral medicine, we purchased it as well with a hope that it could be a potent inhibitor of M<sup>Pro</sup>. We dissolved most purchased small molecule

medicines in DMSO to make 5 mM stock solutions and proceeded to use these stock solutions to test inhibition on M<sup>Pro</sup>. Except itraconazole that has low solubility in DMSO, all tested small molecule medicines were diluted to a 1 mM final concentration in the inhibition assay conditions. We maintained 20% DMSO in the final assay condition to prevent small molecule medicines from precipitation. The activity of M<sup>Pro</sup> in 20% DMSO was a little lower than that in a regular buffer but satisfied our assay requirement. An M<sup>Pro</sup> activity assay in the absence of a small molecule medicine was set up as a comparison. Triplicate repeats were carried out for all tested small molecules and the control. The results presented in Figure 2 displayed two easily discernable characteristics. First, about half of the tested compounds showed strong inhibition of M<sup>Pro</sup> at the 1 mM concentration level (itraconazole was at 0.14 mM due to its low solubility in DMSO), supporting the practical use of a docking method in guiding the drug repurposing research of COVID-19. Second, several small molecule medicines including fexofenadine, indinavir, pirenzepine, reboxetine, and doxapram clearly activated M<sup>Pro</sup> (> 15%). This was to the contrary of what the docking program predicted. This observation strongly suggests that frontline clinicians need to exhibit caution in repurposing medicines for COVID-19 patients before they are thoroughly investigated on influencing the SARS-CoV-2 biology. A not-well-understood drug might deteriorate the already devastating symptoms in COVID-19 patients. Although it is not the focus of the current study, the observation that M<sup>Pro</sup> can be activated by existing drugs needs to be further investigated.

We selected 17 small molecule medicines and rupintrivir that displayed strong inhibition of M<sup>Pro</sup> to conduct further characterizations of their IC<sub>50</sub> values in inhibiting M<sup>Pro</sup> by varying the small molecule concentration from 1 μM to 10 mM. Results collectively presented in Figure 3 identifies that of the 18 tested compounds, 7 had an IC<sub>50</sub> value below 100 μM. These include pimozone, ebastine, rupintrivir, bepridil, sertaconazole, rimonabant, and oxiconazole. Pimozone, ebastine, and bepridil were the three most potent FDA/EMA-approved medicines with IC<sub>50</sub> values as 42 ± 2, 57 ± 12 and 72 ± 12 μM, respectively. Although rupintrivir is a covalent inhibitor that was developed specifically for 3C and 3CL proteases, its IC<sub>50</sub> value (68 ± 7 μM) is higher than that of pimozone and ebastine. The relatively low activity of rupintrivir in inhibiting M<sup>Pro</sup> might be due to the change of the amide bond between the P2 and P3 residues to a methylene ketone. This conversion served to increase the serum stability of rupintrivir, but has likely eliminated a key hydrogen bonding

interaction with  $M^{\text{Pro}}$ .<sup>14</sup> The repurposing of HIV medicines for the treatment of COVID-19, particularly those targeting HIV1 protease, has been area of much attention. In fact, the cocktail of lopinavir and ritonavir was previously tested in China for the treatment of COVID-19.<sup>25</sup> The  $IC_{50}$  value of lopinavir in inhibiting  $M^{\text{Pro}}$  is about 500  $\mu\text{M}$ , which possibly explains why this anti-HIV viral cocktail demonstrated no significant benefit for treating patients. Nelfinavir was previously shown having high potency in inhibiting the entry of SARS-CoV-2 into mammalian cell hosts. A cell-based study in inhibiting the SARS-CoV-2 entry indicated a 1  $\mu\text{M}$   $EC_{50}$  value.<sup>26,27</sup> However, our  $IC_{50}$  determination against  $M^{\text{Pro}}$  resulted in a value of  $234 \pm 5 \mu\text{M}$ , highlighting that nelfinavir likely inhibits another key SARS-CoV-2 enzyme or protein or interferes with key cellular processes that are required for the SARS-CoV-2 entry into host cells. These possibilities need to be studied further. Structurally the two most potent medicines pimozone and ebastine share a same diphenylmethyl moiety. A spatially similar structure moiety N-phenyl-N-benzylamine exists in bepridil. Our docking results suggested a same binding mode for this similar structure moiety in all three drugs (Figure 5). The two aromatic rings occupy the enzyme pockets that associate with the P2 and P4 residues in a substrate. This observation is in line with a crystallographic study that showed two aromatic rings with a single methylene linker bound to the active site of the SARS-CoV enzyme.<sup>28</sup> We believe that the inclusion of the diphenylmethyl moiety in structure-activity relationship studies of  $M^{\text{Pro}}$ -targeting ligands will likely contribute to the identification of both potent and high cell-permeable  $M^{\text{Pro}}$  inhibitors. Figure 4 also revealed large variations in Hill coefficients of  $IC_{50}$  curves for different small molecule medicines ( $IC_{50}$  values and Hill coefficients are summarized in Table 2). Duloxetine and zopiclone gave the two highest Hill coefficients with a gradual  $M^{\text{Pro}}$  activity decrease over an increasing inhibitor concentration. On the contrary, saquinavir and lopinavir yielded lowest Hill coefficients with highly steep  $IC_{50}$  curves. There are three possible explanations for the large discrepancy in Hill coefficients. It could be attributed to different solubility of tested compounds. It is possible that a high DMSO percentage and a relatively high inhibitor concentration created phase transition for some inhibitors.<sup>29</sup> A high Hill coefficient may also be due to different ligand to enzyme ratios when tested compounds bind to  $M^{\text{Pro}}$ . An additionally possible reason is the co-existence of the  $M^{\text{Pro}}$  dimer and monomer in the assay conditions. A previous report showed a  $K_d$  value of the  $M^{\text{Pro}}$  dimerization as 2.5  $\mu\text{M}$ .<sup>19</sup> In theory, the catalytically active dimer species was present at a very low concentration in our assay conditions, leaving the catalytically inactive monomer species as the major form of  $M^{\text{Pro}}$ . In this



situation, the inhibitors that preferentially bind to the M<sup>Pro</sup> dimer and the inhibitors that have a higher affinity to the M<sup>Pro</sup> monomer might yield different Hill coefficients.

From cell biology point of view, our three lead compounds share some similarities with some proposed COVID-19 treatment options. There are reports on the investigation of using hydroxychloroquine to treat COVID-19 patients.<sup>30,31</sup> A likely mechanism of action for hydroxychloroquine is its ability to raise endosomal pH that impacts significantly activities of endosomal proteases that may be required to process the virus membrane proteins.<sup>32,33</sup> Our top three hits pimozone, ebastine, and bepridil are all basic small molecules that can potentiate a similar effect.<sup>34</sup> Among the three drugs, bepridil can be very interesting because it was previously used at a dose of 12 mg/kg for the treatment of Ebola virus infections.<sup>35</sup> In another study, bepridil was administered at a dose of 100 mg/kg for managing atrial arrhythmias, indicating that it has a very low level of cellular toxicity.<sup>36</sup> Moreover, a previous study showed that bepridil can increase the pH of acidic endosomes.<sup>37</sup> Administration of a high dose of bepridil is expected to have dual functions to slow down the virus replication in host cells by both inhibiting M<sup>Pro</sup> and raising the pH of endosomes. Therefore, we urge clinical tests of bepridil in the treatment of COVID-19. The same use of pimozone and ebastine needs to be explored as well.

## CONCLUSION

Guided by a computational docking analysis, we experimentally characterized about 30 FDA/EMA-approved drugs on their inhibition of the essential M<sup>Pro</sup> enzyme of the COVID-19 pathogen SARS-CoV-2. From the study, we identified six FDA/EMA-approved drugs that can potentially inhibit M<sup>Pro</sup> with IC<sub>50</sub> value lower than 100 μM. Three lead drugs pimozide, ebastine, and bepridil are basic molecules. They are expected to exert a similar effect as hydroxychloroquine in raising endosomal pH as well as provide inhibition to M<sup>Pro</sup> for slowing down the SARS-CoV-2 entry and replication in human cell hosts. Given that bepridil has been previously explored to treat Ebola infected patients, we urge a serious consideration of its clinical tests in treating COVID-19. The anti-COVID-19 use of pimozide and ebastine may be explored clinically as well. Our current study indicates that there is a large amount of FDA/EMA-approved drug space open for exploration that could hold promise for repurposing existing drugs to target COVID-19. Performing screening studies on different SARS-CoV-2 protein targets are necessary to uncover existing medicines that may be combined for cocktail treatments of COVID-19. More explorations in this direction are imperative.

## MATERIALS AND METHODS

### Chemicals

We purchased econazole nitrate, duloxetine hydrochloride, doxapram hydrochloride monohydrate, clemastine fumarate salt, sertaconazole nitrate, isavuconazole, rupintrivir, and zopiclone from Sigma-Aldrich, pimavanserin, trihexyphenidyl hydrochloride, reboxetine mesylate, sertindole, bepridil hydrochloride, darunavir, nelfinavir mesylate, indinavir sulfate, lopinavir, tipranavir, saquinavir, pirenzepine hydrochloride, oxiconazole nitrate, pimozide, and rimonabant from Cayman Chemicals, ebastine and itraconazole from Alfa Aesar, metixene hydrochloride hydrate and lemborexant from MedChemExpress, fexofenadine hydrochloride from TCI Chemicals, ketoconazole from Acros Organics, clotiapine from Enzo Life Sciences, and oxyphencyclimine from Boc Sciences. We acquired Sub3 with the sequence as DABCYL-Lys-Thr-Ser-Ala-Val-Leu-Gln-Ser-Gly-Phe-Arg-Lys-Met-Glu-EDANS from Bachem Inc.

### Docking

Autodock 4 was used for all docking analysis. For each small molecule, the genetic algorithm-based calculation was carried out for 100 runs with each run having a maximal number of evaluations as 2,500,000.

### M<sup>Pro</sup> Expression and Purification

We constructed the plasmid pBAD-sfGFP-M<sup>Pro</sup> from pBAD-sfGFP. The M<sup>Pro</sup> gene was inserted between DNA sequences that coded sfGFP and 6xHis. The overall sfGFP-M<sup>Pro</sup>-6xHis fusion gene was under control of a pBAD promoter. The antibiotic selection marker was ampicillin. To express sfGFP-M<sup>Pro</sup>-6xHis, *E. coli* TOP10 cells were transformed with pBAD-sfGFP-M<sup>Pro</sup>. A single colony was picked and grew in 5 mL LB medium with 100 µg/mL ampicillin overnight. In the next day, we inoculated this starting culture into 5 L 2xYT medium with 100 µg/mL ampicillin in 5 separate flasks at 37 °C. When the OD reached to 0.6, we added L-arabinose (working concentration as 0.2%) to each flask to induce protein expression at 37 °C for 4 h. Then, the cells were pelleted at 4000 rpm at 4 °C, washed with cold PBS and stored in -80 °C until purification. To purify the expressed protein, we re-suspended frozen cells in a 125 mL buffer containing Tris pH 7.5, 2.5 mM DTT, and 1.25 mg lysozyme. We sonicated resuspended cells

using a Branson 250W sonicator with 1 second on, 4 second off, and a total 5 min 60% power output in two rounds. After sonication, we spun down the cellular debris at 16000 rpm for 30 min at 4 °C. We collected the supernatant and recorded the volume. The whole cell lysate analysis showed almost all of the fusion protein was hydrolyzed to two separate proteins sfGFP and M<sup>Pro</sup>. We were able to obtain an insignificant amount of M<sup>Pro</sup> when Ni-NTA resins were used for purification. Therefore, we did ammonium sulfate precipitation of the whole cell lysate method. This was done by the addition of a saturated ammonium sulfate solution at 0 °C. We collected the fraction between 30% and 40% of ammonium sulfate. We dissolved the collected fraction in buffer A (20 mM Tris, 10 mM NaCl, and 1 mM DTT at pH 8.0) and dialyzed the obtained solution against the same buffer to remove ammonium sulfate. Then, we subjected this solution to anion exchange column chromatography using Q sepharose resins. We eluted proteins from the Q sepharose column by applying a gradient with increasing the concentration of buffer B (20 mM Tris, 1 M NaCl, and 1 mM DTT at pH 8.0). We concentrated the eluted fractions that contained M<sup>Pro</sup> and subject the concentrated solution to size exclusion chromatography using a HiPrep 16/60 Sephacryl S-100 HR column with a mobile phase containing 10 mM sodium phosphate, 10 mM NaCl, 0.5 mM EDTA and 1 mM DTT at pH 7.8. The final yield of the purified enzyme was 1 mg/L with respect to the original expression medium volume. We determined the concentration of the finally purified M<sup>Pro</sup> using the Pierce™ 660nm protein assay and aliquoted 10 μM M<sup>Pro</sup> in the size exclusion chromatography buffer for storage in -80 °C.

### **The synthesis of Sub1**

We loaded the first amino acid (0.5 mmol, 2 equiv.) manually on chlorotriyl chloride resin (0.52 mmol/g loading) on a 0.25 mmol scale by the addition of DIPEA (3 equiv.). After addition of the first amino acid, automated Fmoc-based solid phases synthesis was performed using a Liberty Blue automated peptide synthesizer. Deprotection of the Fmoc group was carried out with 20% piperidine/DMF. Coupling was done with a Fmoc-protected amino acid (0.75 mmol, 3.0 equiv.) and the coupling reagent HATU (0.9 mmol, 3.6 equiv.) and DIEA in NMP (1 mmol, 4.0 equiv.). The final amino acid was capped by the addition of 25% acetic anhydride (v/v) in DMF and DIEA (0.2mmol, 2.0 equiv.). Coumarin coupling was performed in anhydrous THF using T3P in EtOAc (50% w/v) (3.0 equiv.), DIEA (3 equiv.) and 7-amino-4methyl-coumarin (0.8 equiv.) and mixed for 16 h. The solvent was removed and the peptide was dissolved in DCM and washed

with H<sub>2</sub>O (4x) followed by HCl (2x) and brine (1x). The organic layer was dried with Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated. Global deprotection was then carried out using triisopropylsilane (5%) and trifluoroacetic acid (30%) v/v in DCM and mixed for 2-3 h to result in the crude substrate. The peptide was then purified by semi-preparative HPLC and the fractions containing pure product were pooled, concentrated, and analyzed by LC-MS for purity.

## **The synthesis of Sub2**

We performed automated Fmoc-based solid phase synthesis on a Liberty Blue automated peptide synthesizer. Synthesis was conducted on a 0.1 mmol scale with Fmoc Rink amide MBHA resin (0.52 mmol/g loading) and 3 equiv. of protected amino acids. Deprotection of the Fmoc group was carried out with 20% piperidine/DMF. Coupling was done using the desired Fmoc-protected amino acid (0.2 mmol, 2.0 equiv.), coupling reagent Oxyma (0.4 mmol, 4.0 equiv.) and DIC (0.4 mmol, 4.0 equiv.). After the final amino acid had been coupled on, the resin was washed three times with DMF and DCM. Cleavage from the resin was performed using trifluoroacetic acid (95%), triisopropylsilane (2.5%) and water (2.5%) with agitation for 4 h. The peptide was drained into cold methyl tert-butyl ether where it precipitated out. We centrifuged the precipitate, decanted mother liquor, dissolved the pellet in DMF and then purified the peptide by LCMS.

## **Screening assay**

5 mM stock solutions of medicines were prepared in DMSO. The final screening assay conditions contained 50 nM M<sup>Pro</sup>, 10 μM Sub3, and 1 mM medicine. We diluted enzyme stock and substrate stock solutions using a buffer containing 10 mM sodium phosphate, 10 mM NaCl, and 0.5 mM EDTA at pH 7.8 for reaching desired final concentrations. We ran the assay in triplicates. First, we added 30 μL of a 167 nM M<sup>Pro</sup> solution to each well in a 96-well plate and then provided 20 μL of 5 mM stock solutions of medicines in DMSO. After a brief shaking, we incubated the plate at 37°C for 30 min. Then we added 50 μL of a 20 μM Sub3 solution to initiate the activity analysis. The EDANS fluorescence with excitation at 336 nm and emission at 455 nm from the cleaved substrate was detected. We determined the fluorescence increasing slopes at the initial 5 min and normalized them with respect to the control that had no inhibitor provided.

## **Inhibition analysis**

The final inhibition assay conditions contained 50 nM M<sup>Pro</sup>, 10  $\mu$ M Sub3, and a varying concentration of an inhibitor. Similar to screening assay, we diluted enzyme stock and substrate stock solutions using a buffer containing 10 mM sodium phosphate, 10 mM NaCl, and 0.5 mM EDTA at pH 7.8 for reaching desired final concentrations. We ran the assay in triplicates. For the inhibition analysis, we added 30  $\mu$ L of a 167 nM M<sup>Pro</sup> solution to each well in a 96-well plate and then provided 20  $\mu$ L of inhibitor solutions with varying concentrations in DMSO. After a brief shaking, we incubated the plate at 37°C for 30 min. Then we added 50  $\mu$ L of a 20  $\mu$ M Sub3 solution to initiate the activity analysis. We monitored the fluorescence signal and processed the initial slopes in the same way described in screening assay part. We used GraphPad 8.0 to analyze the data and used the [Inhibitor] vs. response - Variable slope (four parameters) fitting to determine the values of both IC<sub>50</sub> and Hill coefficient.

## REFERENCES

- (1) Gates, B. Responding to Covid-19 - A Once-in-a-Century Pandemic?, *N Engl J Med* **2020**, 382, 1677-1679.
- (2) Morens, D. M.; Daszak, P.; Taubenberger, J. K. Escaping Pandora's Box - Another Novel Coronavirus, *N Engl J Med* **2020**, 382, 1293-1295.
- (3) World Health Organization 2020, May 13.
- (4) Kissler, S. M.; Tedijanto, C.; Goldstein, E.; Grad, Y. H.; Lipsitch, M. Projecting the transmission dynamics of SARS-CoV-2 through the postpandemic period, *Science* **2020**, 10.1126/science.abb5793.
- (5) Morse, J. S.; Lalonde, T.; Xu, S.; Liu, W. R. Learning from the Past: Possible Urgent Prevention and Treatment Options for Severe Acute Respiratory Infections Caused by 2019-nCoV, *ChemBiochem* **2020**, 21, 730-738.
- (6) Morse, J. S.; Lalonde, T.; Xu, S.; Liu, W. R. Learning from the Past: Possible Urgent Prevention and Treatment Options for Severe Acute Respiratory Infections Caused by 2019-nCoV, *ChemRxiv* **2020**, 10.26434/chemrxiv.11728983.v1.
- (7) Zhou, N.; Pan, T.; Zhang, J.; Li, Q.; Zhang, X.; Bai, C.; Huang, F.; Peng, T.; Zhang, J.; Liu, C.; Tao, L.; Zhang, H. Glycopeptide Antibiotics Potently Inhibit Cathepsin L in the Late Endosome/Lysosome and Block the Entry of Ebola Virus, Middle East Respiratory Syndrome Coronavirus (MERS-CoV), and Severe Acute Respiratory Syndrome Coronavirus (SARS-CoV), *J Biol Chem* **2016**, 291, 9218-9232.
- (8) Strating, J. R.; van der Linden, L.; Albulescu, L.; Bigay, J.; Arita, M.; Delang, L.; Leyssen, P.; van der Schaar, H. M.; Lanke, K. H.; Thibaut, H. J.; Ulferts, R.; Drin, G.; Schlinck, N.; Wubbolts, R. W.; Sever, N.; Head, S. A.; Liu, J. O.; Beachy, P. A.; De Matteis, M. A.; Shair, M. D.; Olkkonen, V. M.; Neyts, J.; van Kuppeveld, F. J. Itraconazole inhibits enterovirus replication by targeting the oxysterol-binding protein, *Cell Rep* **2015**, 10, 600-615.
- (9) Mastrangelo, E.; Pezzullo, M.; De Burghgraeve, T.; Kaptein, S.; Pastorino, B.; Dallmeier, K.; de Lamballerie, X.; Neyts, J.; Hanson, A. M.; Frick, D. N.; Bolognesi, M.; Milani, M. Ivermectin is a potent inhibitor of flavivirus replication specifically targeting NS3 helicase activity: new prospects for an old drug, *J Antimicrob Chemother* **2012**, 67, 1884-1894.
- (10) Rossignol, J. F. Nitazoxanide: a first-in-class broad-spectrum antiviral agent, *Antiviral Res* **2014**, 110, 94-103.

- (11) Mercorelli, B.; Palu, G.; Loregian, A. Drug Repurposing for Viral Infectious Diseases: How Far Are We?, *Trends Microbiol* **2018**, *26*, 865-876.
- (12) Hung, I. F.; Lung, K. C.; Tso, E. Y.; Liu, R.; Chung, T. W.; Chu, M. Y.; Ng, Y. Y.; Lo, J.; Chan, J.; Tam, A. R.; Shum, H. P.; Chan, V.; Wu, A. K.; Sin, K. M.; Leung, W. S.; Law, W. L.; Lung, D. C.; Sin, S.; Yeung, P.; Yip, C. C.; Zhang, R. R.; Fung, A. Y.; Yan, E. Y.; Leung, K. H.; Ip, J. D.; Chu, A. W.; Chan, W. M.; Ng, A. C.; Lee, R.; Fung, K.; Yeung, A.; Wu, T. C.; Chan, J. W.; Yan, W. W.; Chan, W. M.; Chan, J. F.; Lie, A. K.; Tsang, O. T.; Cheng, V. C.; Que, T. L.; Lau, C. S.; Chan, K. H.; To, K. K.; Yuen, K. Y. Triple combination of interferon beta-1b, lopinavir-ritonavir, and ribavirin in the treatment of patients admitted to hospital with COVID-19: an open-label, randomised, phase 2 trial, *Lancet* **2020**.
- (13) Baez-Santos, Y. M.; St John, S. E.; Mesecar, A. D. The SARS-coronavirus papain-like protease: structure, function and inhibition by designed antiviral compounds, *Antiviral Res* **2015**, *115*, 21-38.
- (14) Jin, Z.; Du, X.; Xu, Y.; Deng, Y.; Liu, M.; Zhao, Y.; Zhang, B.; Li, X.; Zhang, L.; Peng, C.; Duan, Y.; Yu, J.; Wang, L.; Yang, K.; Liu, F.; Jiang, R.; Yang, X.; You, T.; Liu, X.; Yang, X.; Bai, F.; Liu, H.; Liu, X.; Guddat, L. W.; Xu, W.; Xiao, G.; Qin, C.; Shi, Z.; Jiang, H.; Rao, Z.; Yang, H. Structure of M(pro) from COVID-19 virus and discovery of its inhibitors, *Nature* **2020**, *10.1038/s41586-020-2223-y*.
- (15) Nguyen, D. D.; Gao, K.; Chen, J.; Wang, R.; Wei, G.-W. Potentially highly potent drugs for 2019-nCoV, *bioRxiv* **2020**, 2020.2002.2005.936013.
- (16) Xu, Z.; Peng, C.; Shi, Y.; Zhu, Z.; Mu, K.; Wang, X.; Zhu, W. Nelfinavir was predicted to be a potential inhibitor of 2019-nCov main protease by an integrative approach combining homology modelling, molecular docking and binding free energy calculation, *bioRxiv* **2020**, 2020.2001.2027.921627.
- (17) Wu, C.; Liu, Y.; Yang, Y.; Zhang, P.; Zhong, W.; Wang, Y.; Wang, Q.; Xu, Y.; Li, M.; Li, X.; Zheng, M.; Chen, L.; Li, H. Analysis of therapeutic targets for SARS-CoV-2 and discovery of potential drugs by computational methods, *Acta Pharm Sin B* **2020**.
- (18) Ayman, F.; Ping, W.; Mahmoud, A.; Hesham, S. Identification of FDA Approved Drugs Targeting COVID-19 Virus by Structure-Based Drug Repositioning, *ChemRxiv* **2020**, *10.26434/chemrxiv.12003930.v3*.



- (19) Zhang, L.; Lin, D.; Sun, X.; Curth, U.; Drosten, C.; Sauerhering, L.; Becker, S.; Rox, K.; Hilgenfeld, R. Crystal structure of SARS-CoV-2 main protease provides a basis for design of improved alpha-ketoamide inhibitors, *Science* **2020**, *368*, 409-412.
- (20) Pillaiyar, T.; Manickam, M.; Namasivayam, V.; Hayashi, Y.; Jung, S. H. An Overview of Severe Acute Respiratory Syndrome-Coronavirus (SARS-CoV) 3CL Protease Inhibitors: Peptidomimetics and Small Molecule Chemotherapy, *J Med Chem* **2016**, *59*, 6595-6628.
- (21) Morris, G. M.; Huey, R.; Lindstrom, W.; Sanner, M. F.; Belew, R. K.; Goodsell, D. S.; Olson, A. J. AutoDock4 and AutoDockTools4: Automated docking with selective receptor flexibility, *J Comput Chem* **2009**, *30*, 2785-2791.
- (22) Phan, J.; Zdanov, A.; Evdokimov, A. G.; Tropea, J. E.; Peters, H. K., 3rd; Kapust, R. B.; Li, M.; Wlodawer, A.; Waugh, D. S. Structural basis for the substrate specificity of tobacco etch virus protease, *J Biol Chem* **2002**, *277*, 50564-50572.
- (23) Rut, W.; Groborz, K.; Zhang, L.; Sun, X.; Zmudzinski, M.; Hilgenfeld, R.; Drag, M. Substrate specificity profiling of SARS-CoV-2 M<sup>pro</sup> protease provides basis for anti-COVID-19 drug design, *bioRxiv* **2020**, 2020.2003.2007.981928.
- (24) Dragovich, P. S.; Prins, T. J.; Zhou, R.; Webber, S. E.; Marakovits, J. T.; Fuhrman, S. A.; Patick, A. K.; Matthews, D. A.; Lee, C. A.; Ford, C. E.; Burke, B. J.; Rejto, P. A.; Hendrickson, T. F.; Tuntland, T.; Brown, E. L.; Meador, J. W., 3rd; Ferre, R. A.; Harr, J. E.; Kosa, M. B.; Worland, S. T. Structure-based design, synthesis, and biological evaluation of irreversible human rhinovirus 3C protease inhibitors. 4. Incorporation of P1 lactam moieties as L-glutamine replacements, *J Med Chem* **1999**, *42*, 1213-1224.
- (25) Cao, B.; Wang, Y.; Wen, D.; Liu, W.; Wang, J.; Fan, G.; Ruan, L.; Song, B.; Cai, Y.; Wei, M.; Li, X.; Xia, J.; Chen, N.; Xiang, J.; Yu, T.; Bai, T.; Xie, X.; Zhang, L.; Li, C.; Yuan, Y.; Chen, H.; Li, H.; Huang, H.; Tu, S.; Gong, F.; Liu, Y.; Wei, Y.; Dong, C.; Zhou, F.; Gu, X.; Xu, J.; Liu, Z.; Zhang, Y.; Li, H.; Shang, L.; Wang, K.; Li, K.; Zhou, X.; Dong, X.; Qu, Z.; Lu, S.; Hu, X.; Ruan, S.; Luo, S.; Wu, J.; Peng, L.; Cheng, F.; Pan, L.; Zou, J.; Jia, C.; Wang, J.; Liu, X.; Wang, S.; Wu, X.; Ge, Q.; He, J.; Zhan, H.; Qiu, F.; Guo, L.; Huang, C.; Jaki, T.; Hayden, F. G.; Horby, P. W.; Zhang, D.; Wang, C. A Trial of Lopinavir-Ritonavir in Adults Hospitalized with Severe Covid-19, *N Engl J Med* **2020**, *382*, 1787-1799.
- (26) Musarrat, F.; Chouljenko, V.; Dahal, A.; Nabi, R.; Chouljenko, T.; Jois, S. D.; Kousoulas, K. G. The anti-HIV Drug Nelfinavir Mesylate (Viracept) is a Potent Inhibitor of Cell Fusion

Caused by the SARS-CoV-2 Spike (S) Glycoprotein Warranting further Evaluation as an Antiviral against COVID-19 infections, *J Med Virol* **2020**, *10.1002/jmv.25985*.

- (27) Yamamoto, N.; Matsuyama, S.; Hoshino, T.; Yamamoto, N. Nelfinavir inhibits replication of severe acute respiratory syndrome coronavirus 2 in vitro, *BioRxiv* **2020**, *10.1101/2020.04.06.026476*.
- (28) Williams, A. L.; Yem, D. W.; McGinnis, E.; Williams, L. S. Control of arginine biosynthesis in *Escherichia coli*: inhibition of arginyl-transfer ribonucleic acid synthetase activity, *J Bacteriol* **1973**, *115*, 228-234.
- (29) Shoichet, B. K. Interpreting steep dose-response curves in early inhibitor discovery, *J Med Chem* **2006**, *49*, 7274-7277.
- (30) Chen, Z.; Hu, J.; Zhang, Z.; Jiang, S.; Han, S.; Yan, D.; Zhuang, R.; Hu, B.; Zhang, Z. Efficacy of hydroxychloroquine in patients with COVID-19: results of a randomized clinical trial, *medRxiv* **2020**, *2020.2003.2022.20040758*.
- (31) CHEN Jun, L. D., LIU Li, LIU Ping, XU Qingnian, XIA Lu, LING Yun, HUANG Dan, SONG Shuli, ZHANG Dandan, QIAN Zhiping, LI Tao, SHEN Yinzhong, LU Hongzhou A pilot study of hydroxychloroquine in treatment of patients with moderate COVID-19, *J Zhejiang Univ (Med Sci)* **2020**, *49*, 215-219.
- (32) Savarino, A.; Boelaert, J. R.; Cassone, A.; Majori, G.; Cauda, R. Effects of chloroquine on viral infections: an old drug against today's diseases?, *Lancet Infectious Diseases* **2003**, *3*, 722-727.
- (33) Geleris, J.; Sun, Y.; Platt, J.; Zucker, J.; Baldwin, M.; Hripcsak, G.; Labella, A.; Manson, D.; Kubin, C.; Barr, R. G.; Sobieszczyk, M. E.; Schluger, N. W. Observational Study of Hydroxychloroquine in Hospitalized Patients with Covid-19, *N Engl J Med* **2020**, *10.1056/NEJMoa2012410*.
- (34) Ohkuma, S.; Poole, B. Cytoplasmic vacuolation of mouse peritoneal macrophages and the uptake into lysosomes of weakly basic substances, *J Cell Biol* **1981**, *90*, 656-664.
- (35) DeWald, L. E.; Dyall, J.; Sword, J. M.; Torzewski, L.; Zhou, H.; Postnikova, E.; Kollins, E.; Alexander, I.; Gross, R.; Cong, Y.; Gerhardt, D. M.; Johnson, R. F.; Olinger, G. G., Jr.; Holbrook, M. R.; Hensley, L. E.; Jahrling, P. B. The Calcium Channel Blocker Bepridil Demonstrates Efficacy in the Murine Model of Marburg Virus Disease, *J Infect Dis* **2018**, *218*, S588-S591.

- (36) Miura, S.; Sumiyoshi, M.; Tsuchiya, H.; Maruyama, M.; Seigen, I.; Okai, I.; Masaki, Y.; Okazaki, S.; Inoue, K.; Fujiwara, Y.; Komatsu, K.; Hayashi, H.; Sekita, G.; Tokano, T.; Nakazato, Y.; Daida, H. The use of serum bepridil concentration as a safe rhythm control strategy in patients with atrial tachyarrhythmias, *Journal of Arrhythmia* **2012**, *28*, 187-191.
- (37) Mitterreiter, S.; Page, R. M.; Kamp, F.; Hopson, J.; Winkler, E.; Ha, H. R.; Hamid, R.; Herms, J.; Mayer, T. U.; Nelson, D. J.; Steiner, H.; Stahl, T.; Zeitschel, U.; Rossner, S.; Haass, C.; Lichtenthaler, S. F. Bepridil and amiodarone simultaneously target the Alzheimer's disease beta- and gamma-secretase via distinct mechanisms, *J Neurosci* **2010**, *30*, 8974-8983.

## **ACKNOWLEDGEMENTS**

We thank Profs. Thomas Meek and Kevin Burgess for allowing us to use their instruments. This work was supported in part by National Institutes of Health (grants R01GM127575 and R01GM121584) and Welch Foundation (A-1715).


## **AUTHOR CONTRIBUTIONS**

W.R.L. conceived the project. E.C.V., K.Y., K.C.K., C.-C.C., A.D., D.M.M., S.X., C.-T.K.T., and W.R.L. designed experiments. E.C.V., K.Y., K.C.K., C.-C.C., A.D., D.M.M. performed the experiments. E.C.V., K.Y. and W.R.L. wrote the manuscript. All authors approved the final manuscript before submission.


## **COMPETING FINANCIAL INTERESTS**

The authors declare no competing financial interests.


## FIGURES




**Figure 1:** Structures of 29 FDA/EMA-approved medicines and rupintrivir whose  $IC_{50}$  values in inhibiting  $M^{Pro}$  were determined in the study.



**Figure 2:** Activity of  $\text{M}^{\text{Pro}}$ . (A) The structures of three substrates. (B) Activities of 50 nM and 1  $\mu\text{M}$   $\text{M}^{\text{Pro}}$  on 10  $\mu\text{M}$  Sub1. (C) Activity of 50 nM  $\text{M}^{\text{Pro}}$  on 10  $\mu\text{M}$  Sub2 and Sub3. The fluorescence signals are normalized for easy comparison. (D) Activity of different concentrations of  $\text{M}^{\text{Pro}}$  on 10  $\mu\text{M}$  Sub3.




**Figure 3:** Initial screening of M<sup>pro</sup> inhibition by 29 FDA/EMA-approved medicines and rupintrivir. 1 mM (0.14 mM for Itraconazole due to its low solubility in DMSO) was used for each inhibitor to perform the inhibition assay. Fluorescence intensity was normalized with respect to the control that had no small molecule provided. Triplicate experiments were performed for each compound, and the value was presented as mean  $\pm$  standard error (SE).



**Figure 4:** IC<sub>50</sub> assays for 18 small molecule medicines on their inhibition of M<sup>Pro</sup>. Triplicate experiments were performed for each compound, and the IC<sub>50</sub> value was presented as mean ± standard error (SE). GraphPad Prism 8.0 was used to perform data analysis.





**Figure 5:** Pimozide (A), ebastine (B), bepridil (C), and their overlay (D) in the active site of M<sup>Pro</sup>. The protein surface topography in A, B, and C is presented to show the concaved active site.

**Table 1:** Docking results of small molecule medicines (Compounds whose IC<sub>50</sub> values were tested are asterisked.)

| Name             | $\Delta G_{\text{binding}}$<br>(kcal/mol) | Name              | $\Delta G_{\text{binding}}$<br>(kcal/mol) |
|------------------|---|-------------------|---|
| Rimonabant*      | -11.23                                    | Bepridil*         | -8.31                                     |
| Tipranavir*      | -10.74                                    | Isoconazole       | -8.15                                     |
| Ebastine*        | -10.62                                    | Econazole         | -8.14                                     |
| Saquinavir*      | -10.37                                    | Eluxadoline       | -8.12                                     |
| Zopiclone*       | -10.10                                    | (R)-Butoconazole  | -8.11                                     |
| Pimozide*        | -10.01                                    | (S)-Butoconazole  | -8.10                                     |
| Pirenzepine*     | -9.94                                     | Atazanavir        | -8.08                                     |
| Nelfinavir*      | -9.67                                     | Cetirizine        | -8.01                                     |
| Doxapram*        | -9.55                                     | Efinaconazole     | -8.01                                     |
| Oxiconazole*     | -9.18                                     | Amprenavir        | -7.99                                     |
| Indinavir*       | -9.13                                     | Hydroxyzine       | -7.99                                     |
| Sertindole*      | -9.04                                     | (R)-Tioconazole   | -7.98                                     |
| Metixene*        | -9.01                                     | (R)-Carbinoxamine | -7.96                                     |
| Fexofenadine*    | -8.95                                     | Armodafinil       | -7.90                                     |
| Lopinavir*       | -8.91                                     | Desipramine       | -7.84                                     |
| Sertaconazole*   | -8.87                                     | Ritonavir         | -7.74                                     |
| Reboxetine*      | -8.86                                     | Atomoxetine       | -7.73                                     |
| Ketoconazole*    | -8.85                                     | Sulconazole       | -7.69                                     |
| Duloxetine*      | -8.79                                     | Clotrimazole      | -7.67                                     |
| Isavuconazole*   | -8.77                                     | Dipyridamole      | -7.67                                     |
| Lemborexant*     | -8.75                                     | Phentolamine      | -7.61                                     |
| Oxyphenyclimine* | -8.74                                     | (S)-Tioconazole   | -7.48                                     |
| Darunavir*       | -8.72                                     | Doxylamine        | -7.33                                     |
| Trihexphenidyl*  | -8.72                                     | (S)-Carbinoxamine | -7.21                                     |
| Pimavanserin*    | -8.69                                     | Antazoline        | -6.86                                     |
| Clotiapine*      | -8.57                                     | Voriconazole      | -6.76                                     |
| Itraconazole*    | -8.44                                     | Fluconazole       | -6.41                                     |
| Clemastine*      | -8.36                                     |                   |   |

**Table 2:** IC<sub>50</sub> and Hill coefficient values of 18 identified inhibitors

| Name            | IC <sub>50</sub> (μM) | Hill Slope  |
|-----------------|-----------------------|-------------|
| Pimozide        | 42 ± 2                | 3.1 ± 0.4   |
| Ebastine        | 57 ± 12               | 1.5 ± 0.2   |
| Rupintrivir     | 68 ± 7                | 1.4 ± 0.2   |
| Bepidil         | 72 ± 3                | 2.9 ± 1.0   |
| Sertaconazole   | 76 ± 2                | 3.5 ± 0.2   |
| Rimonabant      | 85 ± 3                | 5.0 ± 0.4   |
| Oxiconazole     | 99 ± 6                | 3.8 ± 0.4   |
| Itraconazole    | 111 ± 35              | 1.6 ± 0.2   |
| Tipranavir      | 180 ± 20              | 1.4 ± 0.2   |
| Nelfinavir      | 234 ± 15              | 5.4 ± 1.0   |
| Zopiclone       | 349 ± 77              | 1.2 ± 0.2   |
| Trihexyphenidyl | 370 ± 53              | 8.9 ± 6.4   |
| Saquinavir      | 411 ± 6               | 26.8 ± 2.6  |
| Isavuconazole   | 438 ± 11              | 5.2 ± 0.7   |
| Lopinavir       | 486 ± 2               | 29.9 ± 2.4  |
| Clemastine      | 497 ± 148             | 11.2 ± 7.3  |
| Metixene        | 635 ± 43              | 8.7 ± 5     |
| Duloxetine      | 3047 ± 634            | 0.93 ± 0.07 |