

Rotigotine Effects on Early Morning Motor Function and Sleep in Parkinson's Disease: A Double-Blind, Randomized, Placebo-Controlled Study (RECOVER)

Claudia Trenkwalder, MD,^{1*} Bryan Kies, FCNeurol (SA),² Monika Rudzinska, MD,³ Jennifer Fine, FCP (SA) Neurology,⁴ Janos Nikl, MD,⁵ Krystyna Honczarenko, MD,⁶ Peter Dioszeghy, MD,⁷ Dennis Hill, MD,⁸ Tim Anderson, FRACP,⁹ Vilho Myllyla, MD,¹⁰ Jan Kassubek, MD,¹¹ Malcolm Steiger, FRCP,¹² Marco Zucconi, MD,¹³ Eduardo Tolosa, MD,¹⁴ Werner Poewe, MD,¹⁵ Erwin Surmann, MSc,¹⁶ John Whitesides, PhD,¹⁷ Babak Boroojerdi, MD,¹⁶ Kallol Ray Chaudhuri, DSc¹⁸ and the RECOVER Study Group

¹Department of Clinical Neurophysiology, University of Goettingen and Paracelsus-Elena Klinik, Kassel, Germany

²Neurology Division, University of Cape Town, South Africa

³Department of Neurology, Jagiellonian University Medical College, Krakow, Poland

⁴Constantiaberg MediClinic, Cape Town, South Africa

⁵Department of Neurology, Hospital of Zala County, Zalaegerszeg, Hungary

⁶Clinic of Neurology, Pomeranian Medical University, Szczecin, Poland

⁷Department of Neurology, County Hospital, Nyiregyhaza, Hungary

⁸Central Carolina Neurology and Sleep, Salisbury, North Carolina, USA

⁹Department of Medicine, University of Otago, Christchurch, and Van Der Veer Institute, Christchurch, New Zealand

¹⁰Department of Neurology, Oulu University Hospital, Oulu, Finland

¹¹Neurologische Klinik, Universität Ulm, Ulm, Germany

¹²The Walton Centre of Neurology and Neurosurgery, Liverpool, United Kingdom

¹³Centro per i Disturbi del Sonno, Istituto Scientifico H San Raffaele, Milan, Italy

¹⁴Neurology Service, Centro de Investigación Biomédica en Red sobre Enfermedades Neurodegenerativas (CIBERNED), Hospital Clínic, IDIBAPS, Universitat de Barcelona, Spain

¹⁵Department of Neurology, Medical University Innsbruck, Innsbruck, Austria

¹⁶UCB Pharma GmbH, Monheim, Germany

¹⁷Schwarz Biosciences Inc, a member of the UCB Group, Raleigh, NC, USA

¹⁸NPF Centre of Excellence, Kings College Hospital and University Hospital Lewisham, Kings College and Institute of Psychiatry, London, United Kingdom

ABSTRACT: In a multinational, double-blind, placebo-controlled trial (NCT00474058), 287 subjects with Parkinson's disease (PD) and unsatisfactory early-morning motor symptom control were randomized 2:1 to receive rotigotine (2–16 mg/24 hr [n = 190]) or placebo (n = 97). Treatment was titrated to optimal dose over 1–8 weeks with subsequent dose maintenance for 4 weeks. Early-morning motor function and nocturnal sleep disturbance were assessed as coprimary efficacy endpoints using the Unified Parkinson's Disease Rating Scale (UPDRS) Part III (Motor Examination) measured in the early morning prior to any medication intake and

the modified Parkinson's Disease Sleep Scale (PDSS-2) (mean change from baseline to end of maintenance [EOM], last observation carried forward). At EOM, mean UPDRS Part III score had decreased by –7.0 points with rotigotine (from a baseline of 29.6 [standard deviation (SD) 12.3] and by –3.9 points with placebo (baseline 32.0 [13.3]). Mean PDSS-2 total score had decreased by –5.9 points with rotigotine (from a baseline of 19.3 [SD 9.3]) and by –1.9 points with placebo (baseline 20.5 [10.4]). This represented a significantly greater improvement with rotigotine compared with placebo on both the UPDRS Part III (treatment difference:

Additional Supporting Information may be found in the online version of this article.

*Correspondence to: Claudia Trenkwalder, Department of Clinical Neurophysiology, University of Goettingen, and Paracelsus Elena-Klinik, Center of Parkinsonism and Movement Disorders, Kassel, Klinikstrasse 16, Kassel, Germany.; ctrenkwalder@gmx.de

Relevant conflicts of interest: All investigators received grant payments from Schwarz Biosciences GmbH, a member of the UCB group of companies, for enrolling patients into the study.

Members of the "RECOVER Study Group" are listed as an Appendix.

Full financial disclosures and author roles may be found in the online version of this article.

Received: 22 June 2010; **Revised:** 6 August 2010; **Accepted:** 27 August 2010

Published online 18 November 2010 in Wiley Online Library (wileyonlinelibrary.com). DOI: 10.1002/mds.23441

−3.55 [95% confidence interval (CI) −5.37, −1.73]; $P = 0.0002$) and PDSS-2 (treatment difference: −4.26 [95% CI −6.08, −2.45]; $P < 0.0001$). The most frequently reported adverse events were nausea (placebo, 9%; rotigotine, 21%), application site reactions (placebo, 4%; rotigotine, 15%), and dizziness (placebo, 6%; rotigotine 10%). Twenty-four-hour transdermal delivery of

rotigotine to PD patients with early-morning motor dysfunction resulted in significant benefits in control of both motor function and nocturnal sleep disturbances. © 2010 *Movement Disorder Society*

Key Words: dopamine agonist; rotigotine; transdermal; motor function; sleep; quality of life

Introduction

Parkinson's disease (PD) is characterized by motor symptoms including nocturnal and early morning dystonia and akinesia and is also associated with nonmotor symptoms, such as sleep disorders.¹ These sleep disorders may be secondary to the patient's underlying disease pathology but may, in part, be a consequence of motor complications and other nonmotor symptoms (such as nocturnal pain, insomnia, and vivid dreams), or medication use. Although the importance of identifying and treating sleep disorders in PD is well recognized, few published trials have prospectively studied their management.^{2–5}

Effective management of nocturnal and early morning PD symptoms may be achieved with either a long-acting treatment or by continuous administration of a short-acting treatment.^{2,3,6} Rotigotine is a non-ergoline dopamine agonist that is applied once daily using a transdermal patch, providing 24 hours of continuous drug delivery, and is generally well tolerated.⁷ Significant treatment benefits of rotigotine have been observed in both early and advanced PD and include improvements in activities of daily living and motor symptoms, clinically relevant reductions in "off" time, and reductions in concomitant L-dopa dose.⁸ Rotigotine's potential to improve early morning motor function and sleep-associated problems in PD was demonstrated recently by two large controlled studies⁹ that showed a reduction in the proportion of patients awakening in an "off" state.

This study compared the effects of rotigotine and placebo on early morning motor function and nocturnal sleep disturbances in subjects with PD. (It should be noted that while dopamine agonists have a variable effect on sleep architecture, this was not addressed in this study.) Effects on nocturnal symptoms such as limb restlessness, tremor, and cramps and on other nonmotor symptoms such as pain and mood were also investigated. To our knowledge, this is the first large, controlled trial in PD to use sleep outcomes as a copriary outcome measure.

Methods

Design

The RECOVER (Randomized Evaluation of the 24-hour Coverage: Efficacy of Rotigotine) study

(NCT00474058) was a Phase 3b, double-blind, placebo-controlled, randomized, parallel-group, two-arm trial undertaken in 49 centers in 12 countries. Subjects were recruited between May 2, 2007 and November 17, 2008. Maximum study duration per subject was 22 weeks, with a maximum duration of treatment of 12 weeks.

Subjects

Men and women (aged ≥ 18 years) with PD (Hoehn and Yahr Stage I-IV; both fluctuators and nonfluctuators) and unsatisfactory control of early morning motor symptoms as determined by the investigator (with the latter intended to reflect clinical practice), were eligible for inclusion. PD was defined by the presence of bradykinesia and at least one of the following: resting tremor, rigidity, or impairment of postural reflexes. Subjects not taking L-dopa were eligible for study inclusion as were those taking immediate-release L-dopa provided they had been on a stable dose for the 28 days prior to baseline (see Supporting Information text for study exclusion criteria).

Antiemetics without central dopaminergic activity were permitted to treat nausea and vomiting that occurred during study drug use. Anticholinergic agents, monoamine oxidase-B inhibitors, NMDA antagonists, entacapone, sedatives, hypnotics, selective serotonin reuptake inhibitors, anxiolytics, and other CNS medications were permitted if dose was stable for the 28 days prior to baseline and likely to remain so for the duration of the study. Controlled-release L-dopa, other centrally acting dopaminergic agents, monoamine oxidase-A inhibitors, tolcapone, budipine, or neuroleptics (except olanzapine, ziprasidone, aripiprazole, clozapine, or quetiapine) were prohibited from 28 days prior to baseline and during the study.

The study was conducted in accordance with Good Clinical Practice and the Declaration of Helsinki. The study protocol and amendments were approved by a national, regional, or Independent Ethics Committee or Institutional Review Board. All subjects provided written, informed consent before study participation.

Protocol

Screening took place up to 4 weeks before the baseline evaluation on Day 1 when subjects were randomized 2:1 to receive rotigotine or placebo, stratified by

site, using a computerized randomization schedule. Starting on Day 1 treatment was administered once daily in the morning using a 24-hour transdermal patch with identical-looking placebo patches to ensure blinding. Treatment was titrated to optimal dose (that at which investigator and subject felt that early morning motor impairment was adequately controlled) over 1–8 weeks, starting at 2 mg/24 hr and increasing in weekly increments of 2 mg/24 hr up to a maximum of 16 mg/24 hr. The dose was maintained at the optimal or maximal dose for a 4-week period (maintenance period) during which dose reduction (and alteration of concomitant L-dopa dose, if applicable) was not permitted. During the titration period, rotigotine dose could be back-titrated once if adverse events (AEs) occurred that were thought to be the result of excessive dopaminergic stimulation (see Supporting Information text). Subjects requiring back-titration immediately proceeded to the maintenance period.

Clinic visits took place at screening and baseline; every 2 weeks during dose titration; start and end of maintenance; and 30 days after treatment ended. Subjects were hospitalized for 2 nights at baseline (the first an adaptation night) and again at end of maintenance. Subjects who withdrew prematurely were asked to return to the clinic for a withdrawal visit. Efficacy assessments were performed after the first or second night of hospitalization depending on the measure (see Supporting Information text) and at the end of maintenance or the withdrawal visit.

Outcome Measures

Coprimary Efficacy Measures

Motor function was assessed in the early morning, before new patch application or PD medication intake, using the Unified Parkinson's Disease Rating Scale (UPDRS) Part III (Motor Examination)¹⁰ while in the practically defined “off” condition. Sleep and nocturnal disability were assessed using a modified version of the original 15-item Parkinson's Disease Sleep Scale (PDSS-2; Fig. 1)¹¹ on which individual items are scored from 0 to 4, with the total score ranging from 0 to 60 where higher scores indicate greater impairment.

Additional Efficacy Measures

The 15 individual items of the PDSS-2 were examined as post hoc efficacy measures. Three PDSS-2 domain scores were also calculated post hoc, by summing individual item scores in groups of five (for a maximum score of 20): “disturbed sleep” (items 1–3, 8, and 14); “motor symptoms at night” (items 4–6, 12, and 13); “PD symptoms at night” (items 7, 9–11, and 15).¹¹

Secondary efficacy measures were the Nocturnal Akinesia, Dystonia and Cramps Score (NADCS),¹² and number of nocturias. Exploratory outcome meas-

ures were the Parkinson's Disease Non-Motor Symptom scale (NMS)¹³; the Beck Depression Inventory (BDI-II)¹⁴; an 11-point Likert pain scale; the short-form Parkinson's Disease Questionnaire (PDQ-8)¹⁵; and the UPDRS Part II (activities of daily living) and UPDRS Part IV (complications of therapy) domains¹⁰ (see Supporting Information text).

Safety and Tolerability Measures

Safety and tolerability were assessed throughout the study and up to 30 days after treatment discontinuation by monitoring the frequency and severity of AEs (including clinically relevant laboratory parameter abnormalities) and any changes in vital signs, physical and neurological findings, and ECGs. Emergence of impulse control disorders was monitored using the modified Minnesota Impulsive Disorder Interview (mMIDI).¹⁶

Statistical Analyses

A sample size of 336 subjects was required to detect a between-group difference of 3.5 points (standard deviation [SD] 9.3) in change from baseline UPDRS Part III score with a power of 90% at the 0.05 significance level using a two-sided *t* test. Unavoidable changes in the manufacturing process for rotigotine during the course of the trial led to recruitment being stopped to avoid enrolled subjects switching to patches from the new manufacturing process; this resulted in recruitment of fewer subjects than planned. Since the power remained >80% at the 0.05 significance level, the reduced sample size was deemed to have no impact on the validity of the trial.

Safety analyses were performed on all randomized subjects who received at least one dose of study drug. Efficacy analyses were performed on the full analysis set (FAS)—all randomized subjects who received at least one dose of study drug and had baseline and at least one post-baseline assessment for both coprimary efficacy measures (mean change in UPDRS Part III and PDSS-2 scores from baseline to end of maintenance [last observation carried forward (LOCF); i.e., modified intent-to-treat analysis]). The two coprimary endpoints were tested in a hierarchical sequential fashion, such that if the testing procedure demonstrated significance at the 5% level on UPDRS Part III, the test for the PDSS-2 score was performed (see Supporting Information text). This enabled evaluation of the motor and non-motor symptoms of PD without adjustments for multiple comparisons. To estimate treatment differences, analyses of covariance were performed with treatment and pooled site as factors and baseline score as the covariate.

Results

Subject Disposition

Of 333 subjects screened, 287 were randomized and 246 (86%) completed the study (Fig. 2). The most

Please rate the severity of the following based on your experiences during the past week (7 days). Please make a cross in the answer box

	Very often <small>(This means 6 to 7 days a week)</small>	Often <small>(This means 4 to 5 days a week)</small>	Sometimes <small>(This means 2 to 3 days a week)</small>	Occasionally <small>(This means 1 day a week)</small>	Never
1) Overall, did you sleep well during the last week?	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4
2) Did you have difficulty falling asleep each night?	<input type="checkbox"/> 4	<input type="checkbox"/> 3	<input type="checkbox"/> 2	<input type="checkbox"/> 1	<input type="checkbox"/> 0
3) Did you have difficulty staying asleep?	<input type="checkbox"/> 4	<input type="checkbox"/> 3	<input type="checkbox"/> 2	<input type="checkbox"/> 1	<input type="checkbox"/> 0
4) Did you have restlessness of legs or arms at nights causing disruption of sleep?	<input type="checkbox"/> 4	<input type="checkbox"/> 3	<input type="checkbox"/> 2	<input type="checkbox"/> 1	<input type="checkbox"/> 0
5) Was your sleep disturbed due to an urge to move your legs or arms?	<input type="checkbox"/> 4	<input type="checkbox"/> 3	<input type="checkbox"/> 2	<input type="checkbox"/> 1	<input type="checkbox"/> 0
6) Did you suffer from distressing dreams at night?	<input type="checkbox"/> 4	<input type="checkbox"/> 3	<input type="checkbox"/> 2	<input type="checkbox"/> 1	<input type="checkbox"/> 0
7) Did you suffer from distressing hallucinations at night (seeing or hearing things that you are told do not exist)?	<input type="checkbox"/> 4	<input type="checkbox"/> 3	<input type="checkbox"/> 2	<input type="checkbox"/> 1	<input type="checkbox"/> 0
8) Did you get up at night to pass urine?	<input type="checkbox"/> 4	<input type="checkbox"/> 3	<input type="checkbox"/> 2	<input type="checkbox"/> 1	<input type="checkbox"/> 0
9) Did you feel uncomfortable at night because you were unable to turn around in bed or move due to immobility?	<input type="checkbox"/> 4	<input type="checkbox"/> 3	<input type="checkbox"/> 2	<input type="checkbox"/> 1	<input type="checkbox"/> 0
10) Did you feel pain in your arms or legs which woke you up from sleep at night?	<input type="checkbox"/> 4	<input type="checkbox"/> 3	<input type="checkbox"/> 2	<input type="checkbox"/> 1	<input type="checkbox"/> 0
11) Did you have muscle cramps in your arms or legs which woke you up whilst sleeping at night?	<input type="checkbox"/> 4	<input type="checkbox"/> 3	<input type="checkbox"/> 2	<input type="checkbox"/> 1	<input type="checkbox"/> 0
12) Did you wake early in the morning with painful posturing of arms and legs?	<input type="checkbox"/> 4	<input type="checkbox"/> 3	<input type="checkbox"/> 2	<input type="checkbox"/> 1	<input type="checkbox"/> 0
13) On waking, did you experience tremor?	<input type="checkbox"/> 4	<input type="checkbox"/> 3	<input type="checkbox"/> 2	<input type="checkbox"/> 1	<input type="checkbox"/> 0
14) Did you feel tired and sleepy after waking in the morning?	<input type="checkbox"/> 4	<input type="checkbox"/> 3	<input type="checkbox"/> 2	<input type="checkbox"/> 1	<input type="checkbox"/> 0
15) Did you wake up at night due to snoring or difficulties with breathing?	<input type="checkbox"/> 4	<input type="checkbox"/> 3	<input type="checkbox"/> 2	<input type="checkbox"/> 1	<input type="checkbox"/> 0

FIG. 1. Modified Parkinson's Disease Sleep Scale (PDSS-2). ©Trenkwalder and Chaudhuri, 2010.

common reason for discontinuation was withdrawn consent. All 287 randomized subjects were evaluated for safety, while 267 subjects were included in the FAS and evaluated for efficacy.


Baseline Characteristics

Baseline demographic and clinical characteristics, including UPDRS Part III and PDSS-2 total scores, were similar between treatment groups (Table 1). Most subjects (91%) had comorbid conditions at study entry, the most common of which were hypertension (44%), constipation (15%), depression (14%),

and hypercholesterolemia (14%). Details of concomitant medication use are provided in the Supporting Information.

Coprimary Efficacy Outcomes

Mean UPDRS Part III scores measured in the early morning and PDSS-2 total scores showed significantly greater improvement with rotigotine than placebo from baseline to end of maintenance (least squares [LS] mean [95% confidence interval (CI)] treatment differences of -3.55 [$-5.37, -1.73$] ($P = 0.0002$) in UPDRS Part III and -4.26 [$-6.08, -2.45$] ($P <$



*1 subject randomized to placebo received one dose of rotigotine during dose de-escalation, and is counted in the rotigotine group for the safety evaluation.

FIG. 2. Subject disposition.

0.0001) in PDSS-2 [Fig. 3A]). These treatment differences were similar for L-dopa treated patients (-3.14 [$-5.22, -1.05$] in UPDRS Part III and -4.11 [$-6.18, -2.05$] in PDSS-2) and de novo patients (-4.84 [$-8.32, -1.36$] in UPDRS Part III and -4.54 [$-7.94, -1.15$] in PDSS-2).

Additional Efficacy Outcomes

All three PDSS-2 domain scores showed significantly greater improvement with rotigotine than placebo from baseline to end of treatment (Fig. 3B). All 15 individual PDSS-2 items except “distressing hallucinations” showed improvement in the rotigotine group; with significant improvements on 10 items, particularly “difficulty falling asleep,” “urge to move arms or legs,” and “uncomfortable and immobile” (Fig. 3B). Likewise, greater improvement was seen with rotigotine than placebo on the NADCS (LS mean treatment difference -0.41 [$-0.79, -0.04$]; $P = 0.030$) (Table 2). The number of nocturias in each group changed by a mean of -0.3 from baseline, and no significant difference was seen between groups (Table 2; LS mean treatment difference -0.02 [$-0.29, 0.25$]; $P = 0.88$).

The mean NMS total score showed greater improvement with rotigotine than placebo from baseline to end of treatment (LS mean [95% CI] treatment difference -6.65 [$-11.99, -1.31$]; $P = 0.015$) (Table 2) with significant differences on the individual scores for sleep/fatigue and mood/cognition (LS mean treatment differences of -2.03 [$-3.31, -0.75$]; $P = 0.002$ and -3.40 [$-5.22, -1.58$]; $P = 0.0003$, respectively). Nu-

merical differences in favor of rotigotine were seen for attention/memory, gastrointestinal tract, urinary, and miscellaneous (Table 2).

Greater improvements with rotigotine than placebo from baseline to end of treatment were seen on the BDI-II (LS mean treatment difference -2.01 [$-3.55, -0.47$]; $P = 0.011$); Likert pain scale (-0.77 [$-1.28, -0.25$]; $P = 0.004$); PDQ-8 (-5.74 [$-8.74, -2.75$]; $P = 0.0002$), and UPDRS Part II (-1.49 [$-2.32, -0.65$]; $P = 0.0005$) (Table 2). UPDRS Part IV individual item scores showed very little change from baseline to end of treatment in either group (see table in Supporting Information text), with no significant between-group differences.

Analysis of all outcome measures using observed cases, and a per protocol analysis of the two copri-mary outcome measures yielded similar results to those reported above using LOCF.

Safety and Tolerability Outcomes

The mean duration of drug exposure was 73 days in the placebo group and 71 days in the rotigotine group. During the titration phase, 7 (7%) placebo-treated and 40 (16%) rotigotine-treated subjects decreased their dose by 2 mg/24 hour (or placebo equivalent) due to tolerability concerns. At the start of maintenance, 67 (35%) rotigotine-treated subjects were taking the maximal dose of 16 mg/24 hour. Overall, 80% of subjects were compliant (placebo, 76%; rotigotine 83%).

The most frequently reported AEs during treatment are shown in Table 3. Two rotigotine-treated subjects

TABLE 1. Baseline demographic and clinical characteristics (safety population^a)

	Placebo (n = 96)	Rotigotine (n = 191)
Age, mean (SD); range	64.4 (10.6); 37–86	64.8 (9.3); 37–85
Gender, n (%)		
Male	61 (64)	123 (64)
Female	35 (36)	68 (36)
BMI, mean (SD) kg/m ² ; range	26.6 (4.5); 16–41	26.6 (4.1); 16–43
Race, n (%)		
White or Caucasian	85 (89)	177 (93)
Black	1 (1)	2 (1)
Asian	1 (1)	1 (<1)
Other	9 (9)	11 (6)
L-DOPA use, n (%)		
No	17 (18)	36 (19)
Yes	79 (82)	155 (81)
Time since first diagnosis, mean (SD) years; range	4.9 (4.6); 0–26	4.6 (4.2); 0–23
Disease severity at baseline, UPDRS Part III sum score categories, n (%)		
0–9	2 (2)	5 (3)
10–19	12 (13)	38 (20)
20–29	32 (33)	60 (31)
30–39	22 (23)	49 (26)
≥40	28 (29)	38 (20)
UPDRS Part III mean (SD) score ^b	32.0 (13.3)	29.6 (12.3)
PDSS-2 total mean (SD) score; range ^b	20.5 (10.4); 3–49	19.3 (9.3); 1–49

^a1 subject randomized to placebo received one dose of rotigotine during dose de-escalation and is counted in the rotigotine group for the safety population.

^bRotigotine, n = 190.

experienced a sleep attack. One rotigotine-treated subject had positive findings of compulsive sexual behavior on the structured psychiatric interview, which was not reported as an impulse control disorder AE (this subject experienced one mild AE of appetite disorder/decreased appetite). A further 9 subjects (placebo, 2 [2%]; rotigotine, 7 [4%]) had a positive result on at least one mMIDI module.

Most reported AEs were mild or moderate in intensity (placebo, 96%; rotigotine, 97%). In both treatment groups 6% of subjects discontinued due to any AE. Five (3%) rotigotine-treated subjects discontinued due to application/instillation site reactions, none of which was considered serious. There was no clear relationship between rotigotine dose and application site reaction rates. Serious AEs were reported by 5 (5%) placebo-treated subjects and 10 (5%) rotigotine-treated subjects; only visual hallucination in 1 rotigotine-treated subject led to study withdrawal. Two subjects, both in the placebo group, died during the study (one committed suicide and the other died of pneumonia aspiration). There were no clinically relevant changes in vital signs or ECG findings, and few laboratory test values of clinical relevance.


Discussion

RECOVER is the first large-scale, double-blind, randomized trial to investigate early morning motor

function and sleep as coprimary outcome measures in PD. In this study, 24-hour transdermal rotigotine treatment was associated with significant benefits versus placebo in the management of early morning motor impairment and nocturnal sleep disturbances. Rotigotine was also associated with improvements in night-time disabilities (such as limb restlessness, immobility, pain, and cramps), and possibly dopaminergic nonmotor daytime symptoms (such as fatigue and mood) as well.

Other dopamine agonists and continuous L-dopa/DDCI infusion have been used successfully for treating early morning motor disabilities.^{3,17,18} The treatment difference of 3.55 points on the UPDRS Part III seen in this 4-week study is consistent with treatment differences of 3.91 and 3.82 points on the UPDRS Part III at the end of 11 weeks of treatment with rotigotine at doses of 6 mg/24 hr and 8 mg/24 hr in an earlier dose-ranging study.¹⁹ Likewise, as for rotigotine in the current study, improvements from baseline of approximately 7 points on the UPDRS Part III have been described over periods of 6–24 weeks for cabergoline,³ L-dopa continuous infusion,¹⁷ and ropinirole controlled release.¹⁸

The clinimetrics of the PDSS are well established²¹ and validated in several studies across the world in independent populations.^{22–25} In this study, a modified version of the PDSS—the PDSS-2¹¹—was used; this modified scale was developed to better reflect treatment effects on nocturnal disabilities and has been



FAS-LOCF: Full Analysis Set-Last Observation Carried Forward; LS: least squares.
 *P < 0.05; **P < 0.01; ***P < 0.001 for rotigotine-placebo treatment difference (ANCOVA; exploratory analyses).
^aPatients rate their severity on each of 15 PDSS-2 questionnaire items (in lower case) according to their frequency during the previous week (very often, often, sometimes, occasionally or never). PDSS-2 domain scores (shown in upper case; DISTURBED SLEEP, MOTOR SYMPTOMS AT NIGHT, and PD SYMPTOMS AT NIGHT) were obtained by summing the relevant group of 5 individual item scores.

FIG. 3. Mean change from baseline to end of maintenance in (A) UPDRS Part III scores and PDSS-2 total scores and (B) PDSS-2 domain and individual item LS mean rotigotine-placebo treatment differences (FAS-LOCF).

validated using patients' and caregivers' observations on nocturnal problems. The significant improvement in sleep seen on the PDSS-2 is consistent with results from a previous prospective open-label trial of rotigotine that used the original PDSS.²⁰ Sleep disturbances in PD patients may be caused by a myriad of factors, including nocturnal motor problems, mood or pain,

and dopaminergic AEs; likewise pain in PD may be the consequence of other PD symptoms, such as akinesia and rigidity. Musculoskeletal problems too might cause nocturnal pain. Improving nocturnal rigidity may benefit sleep and pain directly while better sleep continuity could be expected to improve pain in PD generally. Hence, in this study, the improvement in

TABLE 2. Mean change from baseline to end of treatment in the secondary outcomes, NADCS and number of nocturias, and the exploratory outcomes, NMS total and individual domain scores, BDI-II, Likert pain scale, PDQ-8, UPDRS Part II (FAS-observed cases)

	Mean (SD) baseline score		Mean (SD) change	
	Placebo	Rotigotine	Placebo	Rotigotine
NADCS	2.7 (2.1) (n = 89)	2.9 (2.2) (n = 178)	-0.7 (2.1) (n = 89)	-1.2 (1.8) ^a (n = 178)
Number of nocturias	2.0 (1.8) (n = 89)	1.9 (1.4) (n = 176)	-0.3 (1.6) (n = 89)	-0.3 (1.3) (n = 176)
NMS total score	41.3 (33.5) (n = 87)	41.1 (34.5) (n = 173)	-3.9 (25.5) (n = 86)	-10.3 (21.2) ^a (n = 172)
Individual NMS domain				
Cardiovascular	1.1 (1.9) (n = 89)	1.0 (1.9) (n = 178)	-0.2 (1.7) (n = 88)	0.0 (1.9) (n = 178)
Sleep/fatigue	9.1 (8.6) (n = 89)	9.5 (9.4) (n = 176)	-1.5 (6.2) (n = 88)	-3.7 (6.8) ^b (n = 176)
Mood/cognition	7.3 (10.0) (n = 89)	7.1 (9.3) (n = 178)	0.2 (11.1) (n = 88)	-3.0 (7.5) ^c (n = 178)
Perception/hallucinations	0.4 (1.4) (n = 89)	0.5 (1.7) (n = 178)	0.0 (1.3) (n = 88)	0.1 (2.4) (n = 178)
Attention/memory	3.9 (5.0) (n = 89)	4.5 (6.4) (n = 178)	0.0 (4.8) (n = 88)	-0.3 (4.6) (n = 178)
Gastrointestinal tract	4.2 (5.7) (n = 89)	3.6 (4.4) (n = 178)	0.0 (3.9) (n = 88)	-0.6 (2.6) (n = 178)
Urinary	6.7 (7.5) (n = 89)	6.6 (7.5) (n = 176)	-0.7 (5.0) (n = 88)	-1.1 (4.9) (n = 176)
Sexual function	3.9 (6.3) (n = 87)	3.7 (5.7) (n = 177)	-0.4 (4.3) (n = 86)	-0.2 (4.9) (n = 177)
Miscellaneous	5.1 (5.6) (n = 89)	4.8 (6.4) (n = 178)	-1.0 (4.2) (n = 88)	-1.3 (3.5) (n = 177)
BDI-II	12.6 (8.8) (n = 89)	12.3 (8.0) (n = 178)	-0.8 (7.6) (n = 89)	-2.7 (5.7) ^a (n = 177)
Likert pain scale	2.6 (2.5) (n = 89)	2.8 (2.4) (n = 178)	-0.1 (2.3) (n = 88)	-0.9 (2.2) ^b (n = 178)
PDQ-8	31.1 (17.0) (n = 89)	30.8 (18.2) (n = 177)	-1.2 (13.7) (n = 89)	-6.9 (11.9) ^c (n = 176)
UPDRS Part II	13.5 (6.3) (n = 89)	12.7 (5.6) (n = 178)	-1.3 (3.4) (n = 89)	-2.6 (3.6) ^c (n = 178)

^aP < 0.05; ^bP < 0.01; ^cP < 0.001 for rotigotine-placebo treatment difference (ANCOVA; Exploratory Analyses).

BDI-II, Beck depression inventory; PDQ-8, short-form Parkinson’s Disease questionnaire; NADCS, Nocturnal Akinesia, Dystonia and Cramps Score; NMS, Parkinson’s Disease Nonmotor Symptom Assessment Scale; UPDRS, Unified Parkinson’s Disease Rating Scale; FAS, Full Analysis Set.

sleep produced by rotigotine may be the result of the improvement in nocturnal motor symptoms (restlessness of arms or legs, urge to move arms or legs, painful posturing in the morning, and tremor on awakening). Likewise, improvement on the individual PDSS-2 item—“painful posturing in the morning”—may reflect the overall early morning improvement in akinesia and dystonia observed on the UPDRS Part III: if akinesia and dystonia improve, pain is relieved and in turn an overall early morning benefit is seen. The reported improvements in depressive symptoms

(BDI-II and NMS domain 2) and pain (Likert scale and PDSS-2 item 10) might also have contributed to the better sleep outcomes with rotigotine treatment. In contrast, number of nocturias showed little change during the study and is unlikely to have had any impact on sleep outcomes.

Subjective patient assessments in open-label investigations have suggested that long-acting dopaminergic drugs and formulations may improve sleep dysfunction in PD patients, possibly through improvement of sleep maintenance insomnia.^{2-4,6,17,26} Indeed, continuous delivery of rotigotine through the night could explain the reductions in nocturnal PD symptoms, and might also have counteracted any sleep problems caused by loss of L-dopa activity. Such observations have been reported previously with cabergoline and overnight apomorphine infusion.^{4,27}

As treatment of the nonmotor symptoms of PD is a key unmet need in management of the disease as identified in recently published guidelines,²⁸ an additional outcome in the current study was to investigate the nonmotor effects of rotigotine examined in pilot studies.²⁹ Rotigotine treatment effects in the NMS domains of sleep/fatigue and mood/cognition support the findings from the PDSS-2 and the BDI-II. The treatment benefit seen in the mood/cognition domain is most likely due to an improvement in mood, as the BDI-II showed a significant effect of rotigotine on depressive symptoms.

TABLE 3. Most frequently reported^a treatment-emergent AEs during overall treatment (safety population^b)

Adverse event	Number (%) subjects	
	Placebo (n = 96)	Rotigotine (n = 191)
At least 1 treatment-emergent AE ^c	54 (56)	137 (72)
Nausea	9 (9)	41 (21)
Application and instillation site reactions	4 (4)	29 (15)
Dizziness	6 (6)	20 (10)
Dyskinesia	4 (4)	15 (8)
Headache	5 (5)	13 (7)

^aAt least 5% in either group.

^bOne subject randomized to placebo received one dose of rotigotine during dose de-escalation and is counted in the rotigotine group for the safety evaluation.

^cSubjects could report more than one type of AE.

Rotigotine was generally well tolerated in this study. The reported dopaminergic AEs are comparable with those seen with other nonergoline dopamine agonists.³⁰ The occurrence of skin reactions showed no clear dose relationship and is related to use of the transdermal patch rather than to an allergic reaction to rotigotine. Few subjects showed emergence of impulsive behavior, with only one rotigotine-treated subject having a positive result on the structured psychiatric interview. Two rotigotine-treated subjects reported a sleep attack.

A significant strength of this study is the collection, under standardized conditions, of controlled data describing nocturnal disturbances in PD, including disease-specific problems such as nocturnal akinesia, restlessness, and nightmares. While this and other trials that examine early morning akinesia use data based on the previous night's sleep in a hospital environment, our study also used the PDSS-2, which collects data on the past week of sleep in the patient's own home. However, the lack of sleep laboratory data to measure sleep parameters and their changes objectively is a limitation. In addition, the inclusion of patients based on unsatisfactory control of early morning motor symptoms in the opinion of the investigator may have led to variation in the study population—that the majority (82%) of patients were taking L-dopa at study entry indicates the inclusion of a minority of de novo patients, which may have produced some heterogeneity in the observed treatment response.

Sleep disturbance; nocturnal limb restlessness; cramps, pain, and immobility; and impairment of early morning motor function, mood, and health-related quality of life are common and important nocturnal, early-morning, and daytime problems for PD patients. This study demonstrated significant treatment benefits with rotigotine on each of these outcomes. Indeed, this study is the first to confirm improvements in sleep-related PD problems with a dopaminergic treatment in a multicenter, double-blind, randomized, placebo-controlled trial using a patient-reported outcome as a coprimary outcome measure. Moreover, it has demonstrated the clinical benefits of continuous rotigotine delivery, as efficacy was maintained throughout the night and into the next morning, significantly improving the start of the new day, one of the most troubling times for PD patients. However, whether the effects of rotigotine on the nonmotor symptoms of PD are the result of its effects on motor symptoms, or whether there are additional effects of rotigotine which affect nonmotor qualities such as sleep fragmentation, dreaming, respiration and mood, which are unrelated to an improved motor condition, is still unclear and further objective study of sleep outcomes in PD patients may be warranted.

Appendix

RECOVER Study Group

Marco Onofri, MD, Università G. D'Annunzio di Chieti-Pescara, Italy; Michael William Hayes FRACP, Concord Hospital, Concord, NSW, Australia; Mustafa Saad Siddiqui, MD, Wake Forest University School of Medicine, Winston-Salem, NC, USA; Jens Carsten Möller, MD, Philipps-University, Marburg, Germany; and Jose Maria Gómez-Arguelles, MD, Sanatorio Nuestra Señora del Rosario, Madrid, Spain.

Acknowledgments: The authors acknowledge the following study investigators for their contributions to data acquisition in the RECOVER trial: Poland: Andrzej Tutaj, Zbigniew Stelmasiak, Jerzy Kotowicz, Urszula Fiszer, Walenty Nyka; South Africa: Jonathan Carr, Stanley Lipschitz, Chris Francois Retief; Hungary: Gábor Jacob, László Csiba, Béla Clemens; USA: Alberto Vasquez, Robert Hutchman, Robert Shorr, Joseph Friedman, Todd J Swick; Germany: Reinhard Puzich, Reinhard Ehret, Alexander Storch, Thomas Schwerdtfeger, Johannes Schwarz; UK: Bernard Boothman, Peter Heywood; Italy: Leonardo Scarzella; New Zealand: Stuart Mossman; Australia: Katya E Kotschet, James Temlett; Finland: Esko Kinnunen; Spain: Cristina Ruiz-Huete.

This study was supported by Schwarz Biosciences GmbH, a member of the UCB Group of Companies, delegated by Schwarz Pharma Ltd, Ireland. The authors acknowledge the contribution of Miriam Schmid (Project Manager, Schwarz Biosciences GmbH), who managed the set-up, global conduct, and reporting of the RECOVER trial and Ging-Ging Li (Global Publications Manager, Movement & Sleep Disorders, UCB Pharma SA) for publication coordination. The authors also acknowledge Rebecca Sutch, PhD and Aileen Young, PhD, Evidence Scientific Solutions, Horsham, UK, for writing and editorial assistance, which was contracted by UCB Pharma SA. No undisclosed groups or persons have had a primary role in manuscript preparation.

The PDQ-39 Questionnaire was written by Professor Crispin Jenkinson, Professor Ray Fitzpatrick, and Ms Viv Peto in 1993 who have asserted their moral rights in it and the copyright in which is owned by Isis Innovation Limited. The Questionnaire was first published in 1995 in the journal *Quality of Life Research*. No part of this questionnaire may be reproduced without the prior written consent of Isis Innovation Limited.

References

- Poewe W. Non-motor symptoms in Parkinson's disease. *Eur J Neurol* 2008;15(Suppl 1):14–20.
- Wolters EC, Tesselaar HJ. International (NL-UK) double-blind study of Sinemet CR and standard Sinemet (25/100) in 170 patients with fluctuating Parkinson's disease. *J Neurol* 1996;243:235–240.
- Högl B, Rothdach A, Wetter TC, Trenkwalder C. The effect of cabergoline on sleep, periodic leg movements in sleep, and early morning motor function in patients with Parkinson's disease. *Neuropsychopharmacology* 2003;28:1866–1870.
- Romigi A, Stanzione P, Marciani MG, et al. Effect of cabergoline added to levodopa treatment on sleep-wake cycle in idiopathic Parkinson's disease: an open label 24-hour polysomnographic study. *J Neural Transm* 2006;113:1909–1913.
- Rektorova I, Balaz M, Svatova J, et al. Effects of ropinirole on nonmotor symptoms of Parkinson disease: a prospective multicenter study. *Clin Neuropharmacol* 2008;31:261–266.
- Grandas F, Martínez-Martín P, Linazasoro G. Quality of life in patients with Parkinson's disease who transfer from standard levodopa to Sinemet CR: the STAR Study. The STAR Multicenter Study Group. *J Neurol* 1998;245(Suppl 1):S31–S33.
- Rascol O, Perez-Lloret S. Rotigotine transdermal delivery for the treatment of Parkinson's disease. *Expert Opin Pharmacother* 2009;10:677–691.
- Pham DQ, Nogid A. Rotigotine transdermal system for the treatment of Parkinson's disease. *Clin Ther* 2008;30:813–824.
- Pahwa R, Poewe W, Lyons KE, Boroojerdi B. Changes in early morning motor status following adjunctive treatment of advanced Parkinson's disease with rotigotine transdermal system in two

- large, placebo-controlled trials (PREFER and CLEOPATRA-PD). *Mov Disord* 2009;24:S280.
10. Martínez-Martín P, Gil-Nagel A, Gracia LM, Gómez JB, Martínez-Sarriés J, Bermejo F. Unified Parkinson's Disease Rating Scale characteristics and structure. The Cooperative Multicentric Group. *Mov Disord* 1994;9:76–83.
 11. Trenkwalder C, Kohlen R, Högl B, et al. Parkinson disease sleep scale—Validation of the Revised Version PDSS-2. *Mov Disord* 2010 (in press).
 12. Stocchi F, Barbato L, Nordera G, Berardelli A, Ruggieri S. Sleep disorders in Parkinson's disease. *J Neurol* 1998;245(Suppl 1):S15–8.
 13. Martínez-Martín P, Rodríguez-Blázquez C, Abe K, et al. International study on the psychometric attributes of the Non-Motor Symptoms Scale in Parkinson disease. *Neurology* 2009;73:1584–1591.
 14. Visser M, Leentjens AF, Marinus J, Stiggelbout AM, van Hilten JJ. Reliability and validity of the Beck depression inventory in patients with Parkinson's disease. *Mov Disord* 2006;21:668–672.
 15. Peto V, Jenkinson C, Fitzpatrick R. PDQ-39: a review of the development, validation and application of a Parkinson's disease quality of life questionnaire and its associated measures. *J Neurol* 1998;245(Suppl 1):S10–4.
 16. Christenson GA, Faber RJ, de Zwaan M, et al. Compulsive buying: descriptive characteristics and psychiatric comorbidity. *J Clin Psychiatry* 1994;55:5–11.
 17. Honig H, Antonini A, Martínez-Martín P, et al. Intrajejunal levodopa infusion in Parkinson's disease: a pilot multicenter study of effects on nonmotor symptoms and quality of life. *Mov Disord* 2009;24:1468–1474.
 18. Pahwa R, Stacy MA, Factor SA, et al. Ropinirole 24-hour prolonged release: randomized, controlled study in advanced Parkinson disease. *Neurology* 2007;68:1108–1115.
 19. Parkinson Study Group. A controlled trial of rotigotine monotherapy in early Parkinson's disease. *Arch Neurol* 2003;60:1721–1728.
 20. Giladi N, Mair KJ, Badenhorst F, et al. Effects of rotigotine transdermal patch on early morning motor function, sleep quality, and daytime sleepiness in patients with idiopathic Parkinson's disease. Results of a multicenter, multinational trial. *Eur J Neurol* 2006;13(Suppl. 2):74[P1127].
 21. Chaudhuri KR, Pal S, DiMarco A, et al. The Parkinson's disease sleep scale: a new instrument for assessing sleep and nocturnal disability in Parkinson's disease. *J Neurol Neurosurg Psychiatry* 2002;73:629–635.
 22. Martínez-Martín P, Salvador C, Menéndez-Guisasola L, et al. Parkinson's disease sleep scale: validation study of a Spanish version. *Mov Disord* 2004;19:1226–1232.
 23. Margis R, Donis K, Schönwald SV, et al. Psychometric properties of the Parkinson's disease sleep scale—Brazilian version. *Parkinsonism Relat Disord* 2009;15:495–499.
 24. Uemura Y, Nomura T, Inoue Y, et al. Validation of the Parkinson's disease sleep scale in Japanese patients: a comparison study using the Pittsburgh Sleep Quality Index, the Epworth Sleepiness Scale and Polysomnography. *J Neurol Sci* 2009;287:36–40.
 25. Wang G, Cheng Q, Zeng J, et al. Sleep disorders in Chinese patients with Parkinson's disease: validation study of a Chinese version of Parkinson's disease sleep scale. *J Neurol Sci* 2008;271:153–157.
 26. Eggert K, Schrader C, Hahn M, et al. Continuous jejunal levodopa infusion in patients with advanced parkinson disease: practical aspects and outcome of motor and non-motor complications. *Clin Neuropharmacol* 2008;31:151–166.
 27. Katzenschlager R, Hughes A, Evans A, et al. Continuous subcutaneous apomorphine therapy improves dyskinesias in Parkinson's disease: a prospective study using single-dose challenges. *Mov Disord* 2005;20:151–157.
 28. Zesiewicz TA, Sullivan KL, Arnulf I, et al. Practice parameter: Treatment of nonmotor symptoms of Parkinson disease. Report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology* 2010;74:924–931.
 29. Chaudhuri KR, Schapira AH. Non-motor symptoms of Parkinson's disease: dopaminergic pathophysiology and treatment. *Lancet Neurol* 2009;8:464–474.
 30. Etminan M, Gill S, Samii A. Comparison of the risk of adverse events with pramipexole and ropinirole in patients with Parkinson's disease: a meta-analysis. *Drug Saf* 2003;26:439–444.