

EDITORIAL

BIG FIVE strategies for survival following out-of-hospital cardiac arrest

Bernd W. Böttiger, Lance B. Becker, Karl B. Kern, Freddy Lippert, Andrew Lockey, Giuseppe Ristagno, Federico Semeraro and Sabine Wingen

European Journal of Anaesthesiology 2020, 37:955–958

Sudden cardiac arrest (SCA) in the out-of-hospital setting is the third leading cause of death in industrialised nations, and facts suggest that many of these deaths are potentially avoidable. In Europe and the USA alone, 700 000 people die each year due to SCA despite the fact that emergency medical services (EMS) initiate cardiopulmonary resuscitation (CPR).^{1,2} The same applies to most other parts of the world. Thus, SCA is currently one of the most important healthcare issues, not only because of the opportunities to avoid many of these premature deaths but also because of the huge implications for patients, relatives, healthcare systems and national economies.³

International consensus on our current CPR concepts, procedures and techniques is very well developed following many years of experimental and clinical research. A systematic review of randomised controlled trials has concluded, however, that overall survival has not improved despite 3 decades of initiatives.⁴ Several recent large-scale multicentre trials have failed to show further marked improvements in SCA survival with drugs,^{5–7} airway management techniques and devices,^{8–10} and other technical CPR equipment.^{11–13} It is currently not anticipated that the international guidelines and recommendations on CPR will come up with any new or ‘magic’ CPR strategy, drug or device to further increase survival in the near future.

In contrast, several large-scale studies have shown that it is relatively easy and extremely cost-effective to increase the number of survivors with good neurological outcome following SCA, not with drugs and devices, but with

robust data on cardiac arrest incidence and survival, political interventions and conceptual system changes aimed at strengthening each ring of the ‘chain of survival’.^{14–16} These studies have focused on cultural changes and nationwide campaigns and interventions, which we have summarised as the ‘BIG FIVE strategies for survival’ following SCA. With international implementation of these ‘BIG FIVE for survival’, we believe that it is possible to save several hundreds of thousands of lives after SCA every year worldwide. The ‘BIG FIVE strategies for survival’ following SCA are detailed below and depicted in Fig. 1.

(1) *Community programmes to increase bystander CPR.*


Local, national and international campaigns to increase lay CPR rates (KIDS SAVE LIVES/school-children education in resuscitation,^{14,17–19} ‘World Restart a Heart (WRAH)’ initiative,^{2,20} short CPR courses for adults, media and press campaigns, etc.).¹⁴ Because the brain can normally survive only for 3 to 5 min without any damage, and EMS often arrive later, one major focus to increase survival is the instigation of bystander CPR. This ‘bridging’ of the victim by bystander intervention until EMS arrival will slow down the clock of cerebral hypoxia and preserve the brain in this ‘time window for lay resuscitation’.^{14,17,21,22} In a minority of countries, bystander CPR rates are above 60%. In most countries worldwide, however, bystander CPR rates are below 20 or 30%.^{14,23} It has been shown that bystander CPR is significantly associated with higher survival rates, improved neurological outcome, better quality of life and an increase in return-to-work for SCA patients.^{14,15} This has been demonstrated in

From the Department of Anaesthesiology and Intensive Care Medicine, University Hospital, Medical Faculty of Cologne, Köln, Germany (BWB, SW), Department of Emergency Medicine, Donald and Barbara Zucker School of Medicine at Hofstra/Northwell, New York, New York (LBB), University of Arizona Sarver Heart Center, Tucson, Arizona, USA (KBK), Emergency Medical Services Copenhagen, University of Copenhagen, Copenhagen (FL), Emergency Department, Calderdale Royal Hospital, Halifax, UK (AL), Dipartimento di Fisiopatologia Medico-Chirurgica dei Trapianti, University of Milan, Milan (GR) and Department of Anaesthesia, Intensive Care and Emergency Medical Services, Ospedale Maggiore, Bologna, Italy (FS)

Correspondence to Bernd W. Böttiger, MD, ML, DEAA, FESC, FERC, FAHA, Director Science and Research, European Resuscitation Council (ERC); Chairman, German Resuscitation Council (GRC); Professor and Head of the Department of Anaesthesiology and Intensive Care Medicine, University Hospital of Cologne, Kerpener Straße 62, Köln 50937, Germany

Tel: +49 221 478 82054/+49 221 478 82052; fax: +49 221 478 87811; e-mail: bernd.boettiger@uk-koeln.de

0265-0215 Copyright © 2020 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the European Society of Anaesthesiology. DOI:10.1097/EJA.0000000000001247 This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Fig. 1. The 'BIG FIVE for survival' and their potential impact on survival following out-of-hospital cardiac arrest

ALS, advanced life support; CAC, cardiac arrest centres; CPR, cardiopulmonary resuscitation; EMS, emergency medical services.

Denmark where, over a period of 10 years, a national campaign has increased bystander CPR rates from around 20 to 45%. This was associated with a three-fold increase in survival and better neurological outcome following out-of-hospital cardiac arrest.^{14,15} Thus, increasing bystander CPR rates from lower levels to 50% and more is associated with a three-fold increase in survival following SCA.

(2) Dispatcher-assisted or telephone CPR.

In most emergency calls, the caller does not recognise that the victim is in cardiac arrest and does not start CPR spontaneously. Therefore, prompt recognition of cardiac arrest by the dispatcher who can then motivate the caller to start CPR is important.²⁴ It has been demonstrated clearly that instructions for chest compressions given by the dispatcher via phone are feasible and most effective. Telephone CPR may be combined with support from specific protocols, computer applications and techniques that allow the dispatcher to receive more information from the scene and the victim. The number needed to treat for telephone CPR has been calculated to be around

seven.^{24,25} Therefore, telephone CPR is associated with an up to two-fold increase in survival following SCA.²⁴

(3) First responder programmes to start CPR and use public access defibrillators.

Trained and/or untrained persons and independent medical personnel from nearby can be alerted in the case of SCA by the dispatch centre in parallel with the EMS.^{26–28} Several studies have shown a significant increase in the rate of CPR provided before EMS arrival and a potential increase in overall survival (OS).^{26,29} Early defibrillation using public access defibrillators delivered by lay or professional first responders has been shown to correlate with increased survival after out-of-hospital cardiac arrest, with reported median survival rates by lay responders of 53% (range 26 to 72) in one systematic review.³⁰ First responders have a high potential and are particularly helpful when bystander CPR rates are low and/or response times for EMS are long. According to the available studies, implementation of first responder programmes can thus be associated with an estimated

0.2 to 2-fold increase in survival, depending on and determined by the underlying culture and system characteristics.

(4) High-quality CPR.

Taking care of SCA patients by an EMS staffed with well-trained advanced life support paramedics and physicians in the out-of-hospital setting is associated with a two-fold increase in short-term and long-term survival.^{31,32} This has been demonstrated in several single and multicentre trials, comparisons and meta-analyses all around the world.^{31,32} Few other system configurations with high density levels of first responders and extremely short response times have achieved similar levels of outcomes worldwide.^{31,32}

(5) Specialised postresuscitation care.

In 60 to 80% of all SCA patients, acute coronary syndrome and/or acute myocardial infarction are the underlying causes of deterioration.^{33,34} All registry data and several prospective studies have demonstrated that treating the underlying cause of SCA by immediate acute percutaneous coronary intervention (PCI) within 60 or 90 min in a specialised cardiac arrest centre with 24/7 PCI availability is associated with a doubling in survival.^{35–39} SCA patients with coronary problems may need PCI at least as fast as patients with acute coronary syndrome and without cardiac arrest or shock. Even transport of SCA patients with ongoing CPR to a PCI facility with subsequent intervention may be associated with good outcomes.³⁸ Extracorporeal membrane oxygenation with transporting devices and subsequent PCI can also be indicated in selected patients, but clear outcome data are missing.⁴⁰ Adequate temperature management, optimised haemodynamic and ventilatory support, prognostication and other individualised interventions in specific circumstances, such as treatment of tension pneumothorax in traumatic cardiac arrest and specific interventions in patients suffering from acute pulmonary embolism etc., are further important quality and outcome indicators of specialised centres.⁴¹ Thus, ultrafast and straightforward management of SCA patients in specifically staffed and equipped hospitals, so-called cardiac arrest centres, seems to further improve survival by around two-fold.^{38,41}

Successful treatment of SCA patients to increase survival rates and neurological recovery has definitely moved the focus to the out-of-hospital setting, as by far the biggest impact on the chain of survival is within the first links. Implementing the BIG FIVE will, with current evidence, markedly improve the outcome of SCA patients worldwide.

Moreover, public awareness, motivating, educating and involving lay people – and school children in particular – has a major social impact, promotes empathy and is establishing a general culture of assisting the community.

A critical foundation for all these life-saving strategies is for regions and nations to create a robust cardiac arrest strategy and a registry or database that allows accurate determination of cardiac arrest incidence and survival rates.^{4–16,23} National cardiac arrest registries promote continuous quality improvement efforts, allow for identification of areas of strengths and weaknesses in the chain of survival, promote public health initiatives and will allow for identification of future opportunities. Nations with robust cardiac arrest data often enjoy significantly improved survival rates over relatively short periods of time.^{14–16,23} The experiences from Denmark and elsewhere around the world^{2,14–16} can and should serve as a blueprint to increase survival following SCA in all countries. Worldwide, we propose that these ‘BIG FIVE for survival’ strategies are the most important impact factors for increasing overall survival with good neurological recovery after SCA as well as improving the overall national health and global economics in industrialised countries.

Acknowledgements relating to this article

Assistance with the Editorial: none.

Financial support and sponsorship: none.

Conflicts of interest: BWB is European Resuscitation Council (ERC) Board Director Science and Research; Chairman of the German Resuscitation Council (GRC); Member of the Advanced Life Support (ALS) Task Force of the International Liaison Committee on Resuscitation (ILCOR). LBB has received institutional grants/research support from the National Institutes of Health (NIH) and he is a long-standing volunteer member of the American Heart Association (currently serving on several committees) which has a financial interest in the outcome of resuscitation studies being conducted. The American Heart Association sells training materials worldwide on resuscitation techniques. KBK is the current Chairman of the American Heart Association’s Emergency Cardiovascular Care committee and a delegate to ILCOR. FL has received unrestricted research grants from the Laerdal Foundation and the Danish TrygFonden. AL is Vice President of the Resuscitation Council UK, Member of the ERC Educational Group, Domain Leader and Member of the Education, Implementation and Teams Task Force of ILCOR. GR is Domain Leader for ‘drugs and fluids’ of ILCOR; Member of the ERC Science and Education (SEC) on Basic Life Support (BLS); Member of the BLS Task Force of ILCOR. FS is ERC SEC BLS Co-Chair; Past President of the Italian Resuscitation Council (IRC); Member of the ‘Digital Communication Working Group’ of ILCOR. SW is Executive personal assistant of the Board of the German Resuscitation Council. Some authors have received support from industry for studies and presentations, although these are not considered to be conflicts of interest regarding the topics discussed in this article.

Comment from the Editor: none.

References

- 1 Mozaffarian D, Benjamin EJ, Go AS, *et al.*, Writing Group Members; American Heart Association Statistics Committee; Stroke Statistics Subcommittee. Executive summary: heart disease and stroke statistics – 2016 update: a report from the American Heart Association. *Circulation* 2016; **133**:447–454.

- 2 Böttiger BW, Lockey A, Aickin R, et al. 'All citizens of the world can save a life' – The World Restart a Heart (WRAH) initiative starts in 2018. *Resuscitation* 2018; **128**:188–190.
- 3 Ornato JP, Becker LB, Weisfeldt ML, et al. Cardiac arrest and resuscitation: an opportunity to align research prioritization and public health need. *Circulation* 2010; **122**:1876–1879.
- 4 Sasson C, Rogers MA, Dahl J, et al. Predictors of survival from out-of-hospital cardiac arrest: a systematic review and meta-analysis. *Circ Cardiovasc Qual Outcomes* 2010; **3**:63–81.
- 5 Böttiger BW, Arntz HR, Chamberlain DA, et al., TROICA Trial Investigators; European Resuscitation Council Study Group. Thrombolysis during resuscitation for out-of-hospital cardiac arrest. *N Engl J Med* 2008; **359**:2651–2662.
- 6 Kudenchuk PJ, Brown SP, Daya M, et al., Resuscitation Outcomes Consortium Investigators. Amiodarone, lidocaine, or placebo in out-of-hospital cardiac arrest. *N Engl J Med* 2016; **374**:1711–1722.
- 7 Perkins GD, Ji C, Deakin CD, et al., PARAMEDIC2 Collaborators. A randomized trial of epinephrine in out-of-hospital cardiac arrest. *N Engl J Med* 2018; **379**:711–721.
- 8 Wang HE, Schmicker RH, Daya MR, et al. Effect of a strategy of initial laryngeal tube insertion vs endotracheal intubation on 72-h survival in adults with out-of-hospital cardiac arrest: a randomized clinical trial. *JAMA* 2018; **320**:769–778.
- 9 Jabre P, Penalzoza A, Pinero D, et al. Effect of bag-mask ventilation vs endotracheal intubation during cardiopulmonary resuscitation on neurological outcome after out-of-hospital cardiorespiratory arrest: a randomized clinical trial. *JAMA* 2018; **319**:779–787.
- 10 Bengier JR, Kirby K, Black S, et al. Effect of a strategy of a supraglottic airway device vs tracheal intubation during out-of-hospital cardiac arrest on functional outcome: the AIRWAYS-2 randomized clinical trial. *JAMA* 2018; **320**:779–791.
- 11 Wik L, Olsen JA, Persse D, et al. Manual vs. integrated automatic load-distributing band CPR with equal survival after out of hospital cardiac arrest. The randomized CIRC trial. *Resuscitation* 2014; **85**:741–748.
- 12 Rubertsson S, Lindgren E, Smekal D, et al. Mechanical chest compressions and simultaneous defibrillation vs conventional cardiopulmonary resuscitation in out-of-hospital cardiac arrest: the LINC randomized trial. *JAMA* 2014; **311**:53–61.
- 13 Perkins GD, Lall R, Quinn T, et al., PARAMEDIC Trial Collaborators. Mechanical versus manual chest compression for out-of-hospital cardiac arrest (PARAMEDIC): a pragmatic, cluster randomised controlled trial. *Lancet* 2015; **385**:947–955.
- 14 Wissenberg M, Lippert FK, Folke F, et al. Association of national initiatives to improve cardiac arrest management with rates of bystander intervention and patient survival after out-of-hospital cardiac arrest. *JAMA* 2013; **310**:1377–1384.
- 15 Kragholm K, Wissenberg M, Mortensen RN, et al. Bystander efforts and 1-year outcomes in out-of-hospital cardiac arrest. *N Engl J Med* 2017; **376**:1737–1747.
- 16 Kragholm K, Wissenberg M, Mortensen RN, et al. Return to work in out-of-hospital cardiac arrest survivors: a nationwide register-based follow-up study. *Circulation* 2015; **131**:1682–1690.
- 17 Böttiger BW, Bossaert LL, Castrén M, et al., Board of European Resuscitation Council (ERC). Kids save lives – ERC position statement on school children education in CPR: 'Hands that help – training children is training for life'. *Resuscitation* 2016; **105**:A1–A3.
- 18 Böttiger BW, Semeraro F, Altemeyer KH, et al. Kids save lives: school children education in resuscitation for Europe and the world. *Eur J Anaesthesiol* 2017; **34**:792–796.
- 19 Wingen S, Schroeder DC, Ecker H, et al. Self-confidence and level of knowledge after cardiopulmonary resuscitation training in 14 to 18-year-old schoolchildren: a randomised-interventional controlled study in secondary schools in Germany. *Eur J Anaesthesiol* 2018; **35**:519–526.
- 20 Böttiger BW, Lockey A. World restart a heart initiative: all citizens of the world can save a life. *Lancet* 2018; **392**:1305.
- 21 Breckwoldt J, Schloesser S, Arntz HR. Perceptions of collapse and assessment of cardiac arrest by bystanders of out-of-hospital cardiac arrest (OOHCA). *Resuscitation* 2009; **80**:1108–1113.
- 22 Böttiger BW, Dirks B, Just U, et al. The 10 fundamental principles of lay resuscitation: recommendations by the German Resuscitation Council. *Eur J Anaesthesiol* 2018; **35**:721–723.
- 23 Gräsner JT, Lefering R, Koster RW, et al. EuReCa ONE-27 Nations, ONE Europe, ONE Registry: a prospective one month analysis of out-of-hospital cardiac arrest outcomes in 27 countries in Europe. *Resuscitation* 2016; **105**:188–195.
- 24 Viereck S, Møller TP, Ersbøll AK, et al. Recognising out-of-hospital cardiac arrest during emergency calls increases bystander cardiopulmonary resuscitation and survival. *Resuscitation* 2017; **115**:141–147.
- 25 Hallstrom A, Cobb L, Johnson E, et al. Cardiopulmonary resuscitation by chest compression alone or with mouth-to-mouth ventilation. *N Engl J Med* 2000; **342**:1546–1553.
- 26 Ringh M, Rosenqvist M, Hollenberg J, et al. Mobile-phone dispatch of laypersons for CPR in out-of-hospital cardiac arrest. *N Engl J Med* 2015; **372**:2316–2325.
- 27 Zijlstra JA, Stieglis R, Riedijk F, et al. Local lay rescuers with AEDs, alerted by text messages, contribute to early defibrillation in a Dutch out-of-hospital cardiac arrest dispatch system. *Resuscitation* 2014; **85**:1444–1449.
- 28 Smith CM, Wilson MH, Ghorbangholi A, et al. The use of trained volunteers in the response to out-of-hospital cardiac arrest – the GoodSAM experience. *Resuscitation* 2017; **121**:123–126.
- 29 Hüpfel M, Selig HF, Nagele P. Chest-compression-only versus standard cardiopulmonary resuscitation: a meta-analysis. *Lancet* 2010; **376**:1552–1557.
- 30 Bækgaard JS, Viereck S, Møller TP, et al. The effects of public access defibrillation on survival after out-of-hospital cardiac arrest: a systematic review of observational studies. *Circulation* 2017; **136**:954–965.
- 31 Böttiger BW, Bernhard M, Knapp J, et al. Influence of EMS-physician presence on survival after out-of-hospital cardiopulmonary resuscitation: systematic review and meta-analysis. *Crit Care* 2016; **20**:4.
- 32 Fischer M, Krep H, Wierich D, et al. Comparison of the emergency medical services systems of Birmingham and Bonn: process efficacy and cost effectiveness. *Anesthesiol Intensivmed Notfallmed Schmerzther* 2003; **38**:630–642.
- 33 Grunau B, Kawano T, Scheuermeyer F, et al. Early advanced life support attendance is associated with improved survival and neurologic outcomes after nontraumatic out-of-hospital cardiac arrest in a tiered prehospital response system. *Resuscitation* 2018; **135**:137–144.
- 34 Spaulding CM, Joly LM, Rosenberg A, et al. Immediate coronary angiography in survivors of out-of-hospital cardiac arrest. *N Engl J Med* 1997; **336**:1629–1633.
- 35 Anyfantakis ZA, Baron G, Aubry P, et al. Acute coronary angiographic findings in survivors of out-of-hospital cardiac arrest. *Am Heart J* 2009; **157**:312–318.
- 36 Dumas F, White L, Stubbs BA, et al. Long-term prognosis following resuscitation from out of hospital cardiac arrest: role of percutaneous coronary intervention and therapeutic hypothermia. *J Am Coll Cardiol* 2012; **60**:21–27.
- 37 Camuglia AC, Randhawa VK, Lavi S, et al. Cardiac catheterization is associated with superior outcomes for survivors of out of hospital cardiac arrest: review and meta-analysis. *Resuscitation* 2014; **85**:1533–1540.
- 38 Scholz KH, Maier SKG, Maier LS, et al. Impact of treatment delay on mortality in ST-segment elevation myocardial infarction (STEMI) patients presenting with and without haemodynamic instability: results from the German prospective, multicentre FITT-STEMI trial. *Eur Heart J* 2018; **39**:1065–1074.
- 39 Søholm H, Kjaergaard J, Bro-Jeppesen J, et al. Prognostic implications of level-of-care at tertiary heart centers compared with other hospitals after resuscitation from out-of-hospital cardiac arrest. *Circ Cardiovasc Qual Outcomes* 2015; **8**:268–276.
- 40 Debaty G, Babaz V, Durand M, et al. Prognostic factors for extracorporeal cardiopulmonary resuscitation recipients following out-of-hospital refractory cardiac arrest. A systematic review and meta-analysis. *Resuscitation* 2017; **112**:1–10.
- 41 Nichol G, Aufderheide TP, Eigel B, et al. Regional systems of care for out-of-hospital cardiac arrest: a policy statement from the American Heart Association. *Circulation* 2010; **121**:709–729.